

Phanerozoic marine biodiversity dynamics in light of the incompleteness of the fossil record

Peter J. Lu*[†], Motohiro Yogo*[‡], and Charles R. Marshall[¶]

*Department of Physics, Jefferson Laboratory, 17 Oxford Street, [‡]Department of Economics, and [¶]Museum of Comparative Zoology, Department of Organismic and Evolutionary Biology, and Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA 02138

Communicated by Steven M. Stanley, University of Hawaii at Manoa, Honolulu, HI, December 22, 2005 (received for review April 21, 2005)

Long-term evolutionary dynamics have been approached through quantitative analysis of the fossil record, but without explicitly taking its incompleteness into account. Here we explore the temporal covariance structure of per-genus origination and extinction rates for global marine fossil genera throughout the Phanerozoic, both before and after corrections for the incompleteness of the fossil record. Using uncorrected data based on Sepkoski's compendium, we find significant autocovariance within origination and extinction rates, as well as covariance between extinction and origination, not one, but two, intervals later, corroborating evidence for the unexplained temporal gap found by past studies. However, these effects vanish when the data are corrected for the incompleteness of the fossil record. Instead, we observe significant covariance only between extinction and origination in the immediately following intervals. The gap in the response of the biosphere to extinction in the uncorrected fossil record thus appears to be an artifact of the incompleteness of the fossil record, specifically due to episodic variation in the probability that taxa will be preserved, on time scales comparable to the temporal resolution of Sepkoski's data. Our results also indicate that at that temporal resolution (the stage/substage of duration ≈ 5 million years), changes in origination and extinction do not persist for longer than one interval, except that elevated origination rates immediately after extinction may last for more than a single interval. Thus, although certain individual cases may deviate from the overall pattern, we find that in general the biosphere's response to perturbation is immediate geologically and usually short-lived.

extinction | Jaanusson effect | origination | Signor-Lipps effect | vector autoregression

Sepkoski's compendia of marine fossil families (1) and genera (2, 3) have been central to the analysis of the long-term patterns of origination, extinction, and overall diversity of marine animals throughout the Phanerozoic. Many analyses have looked for temporal correlations between rates of origination, extinction, and total biodiversity in stages and substages of the Phanerozoic (4–9). Of special interest has been the discovery of the delayed recovery of diversity after times of unusually high extinction (5, 6), as well as the observation that times of high extinction and origination tend to persist for more than a substage (5–7). These results suggest either that the causes of elevated rates of origination or extinction persist for many millions of years, and/or that the biosphere has some intrinsic limits in how fast it can recover from extinction (6).

However, as all of these studies have recognized, Sepkoski's raw data do not include any corrections for the incompleteness of the fossil record. Significantly, there is growing evidence that this incompleteness, and its temporal variation, can have significant effects on perceived patterns of origination, extinction, and biodiversity in the fossil record (10–12). While some analyses have been adjusted in an attempt to account for incompleteness (e.g., ref. 6), none has performed a basic time-series analysis on a data set that has used geological and paleontological first principles to quantitatively correct the observed rates of origi-

nation and extinction derived from Sepkoski's compendium of marine genera for the effects of incompleteness. Foote has recently presented such a data set (13), and its differences from Sepkoski's raw data are striking. Specifically, Foote has noted that the measured peaks of origination and extinction in the corrected data are far more volatile than in Sepkoski's uncorrected data (representative time series are presented in Fig. 1), and he cautioned that between-stage correlations in rates of origination and extinction may indeed be artifacts created by the incompleteness of the fossil record.

Analytic Approach

Here we examine how the previously reported correlations of between-stage rates of origination and extinction (5, 6, 8) are affected when Foote's corrected data are subjected to time-series analysis. Specifically, we use vector autoregression (VAR) analysis (14) to examine the temporal covariance structure between per-genus origination and extinction rates of Phanerozoic marine genera, for both Sepkoski's raw data and the data corrected by Foote for the incompleteness of the fossil record (13).

For each i th interval of duration t_i in the time-series data (Fig. 1), we calculated the covariances between per-genus rates of origination O_i and extinction E_i with the originations O_{i-n} and extinctions E_{i-n} in the n previous intervals, denoted VAR(n). Combining originations and extinctions into the two-dimensional vector $\mathbf{x}_i = (O_i, E_i)$, the VAR(1) model is $\mathbf{x}_i = \Phi_1 \mathbf{x}_{i-1} + \varepsilon_i$, where Φ_1 is the 2×2 matrix of normalized covariance coefficients relating to origination and extinction one interval previous, ε_i represents serially uncorrelated random noise, and intercepts have been omitted to simplify notation. The VAR(2) model extends the analysis to two lags, and is defined as $\mathbf{x}_i = \Phi_1 \mathbf{x}_{i-1} + \Phi_2 \mathbf{x}_{i-2} + \varepsilon_i$, explicitly:

$$\begin{bmatrix} O_i \\ E_i \end{bmatrix} = \begin{bmatrix} \Phi_1^{oo} & \Phi_1^{eo} \\ \Phi_1^{oe} & \Phi_1^{ee} \end{bmatrix} \begin{bmatrix} O_{i-1} \\ E_{i-1} \end{bmatrix} + \begin{bmatrix} \Phi_2^{oo} & \Phi_2^{eo} \\ \Phi_2^{oe} & \Phi_2^{ee} \end{bmatrix} \begin{bmatrix} O_{i-2} \\ E_{i-2} \end{bmatrix} + \varepsilon_i. \quad [1]$$

Previous analyses (4, 6, 8) have determined autocorrelations and cross-correlations, separately normalized to the range from -1 to $+1$, which precludes direct numerical comparison between the magnitudes of an autocorrelation within a time series and a cross-correlation with another. Here the multidimensional VAR technique (14) simultaneously determines the relative contributions of the two, but also uses a different normalization: clearly, autocovariance and covariance coefficients cannot both be constrained to the same finite range if their relative magnitudes are not bounded. Instead, the VAR coefficients are normalized by the inverse of the variance, which in our two-dimensional system constrains the autocovariance coefficients (diagonal Φ matrix

Conflict of interest statement: No conflicts declared.

Abbreviation: VAR, vector autoregression.

[†]To whom correspondence should be addressed. E-mail: plu@fas.harvard.edu.

[‡]Present address: Rolling Green Golf Club, 280 North State Road, Springfield, PA 19064.

© 2006 by The National Academy of Sciences of the USA




Fig. 1. Per-genus rates of (a) origination O_i and (b) extinction E_i as a function of time. Uncorrected data (Upper) and data corrected for the incompleteness of the fossil record (Lower) are presented. Diversity has been counted by using the boundary-crosser method, assuming pulsed turnover (13). Different geological periods are indicated in standard colors; average rates are indicated by dotted lines.

elements) to the range from -1 to $+1$, the same as for the more familiar autocorrelation (the autocovariance of a time series of one-dimensional scalars). An origination autocovariance coefficient $\Phi_1^{oo} = 0.5 \pm 0.2$ implies that, holding extinction at its average value, if origination in the previous time interval is 1% above average, origination will be $0.5 \pm 0.2\%$ above average in the present interval. By contrast, the covariance coefficients (off-diagonal Φ matrix elements) are determined relative to the normalized autocovariance coefficients; they are therefore numerically unconstrained, and in particular can have values greater than $+1$. Although these covariance coefficients cannot be easily mapped to cross-correlation coefficients, quantitative interpretation is nonetheless straightforward. For example, a value of the covariance coefficient $\Phi_1^{oe} = 1.5 \pm 0.5$ implies that, holding extinction at its average value, if origination in the previous time interval is 1% above average, extinction will be $1.5 \pm 0.5\%$ above average in the present interval.

Data

We analyzed several published tabulations of per-genus rates of origination and extinction of Phanerozoic genera. The uncorrected raw binned data come from the compendium of fossil genera published by Sepkoski (3), and have been used in previous studies (6–8). We used the 101 time intervals running from the Cambrian (543 million years ago) to the present, and excluded genera appearing in only a single interval (“singletons”) to minimize Lagerstätten and monographic effects. Per-genus rates of origination (extinction) were defined as the fraction of total genera that appeared (disappeared) in an interval.

A second major tabulation is Foote’s boundary-crossing data (13), which were broken into 77 intervals corresponding mainly to geological stages. With the boundary-crosser approach, we have only point estimates of origination and extinction rates.

Compiling the rates of origination and extinction depends on where in the intervening interval the taxa actually originated or went extinct. Foote used two end-member approaches for dealing with this uncertainty: (i) he assumed that taxa originated and went extinct at a uniform rate throughout the interval (the model of continuous turnover); or, (ii) he assumed that all originations occur at the beginning, and all extinctions, at the end, of the interval (the model of pulsed turnover). The uncorrected data presented in Fig. 1 are tabulated assuming this pulsed model (figure 2 of ref. 13), which currently seems the more likely approach (15).

Neither tabulation method, however, accounts for the incompleteness of the fossil record. Foote has used preservation and rock volume effects to estimate true origination and extinction per-taxon rates, in both continuous and pulsed turnover scenarios (figures 3 and 4 of ref. 13), which we term “corrected” data. While it is inherently challenging to verify any model that corrects for missing data, to date Foote is the only one to have made such a correction, and it highly desirable to determine what effect these corrections have on the perceived diversity dynamics derived from analysis of the uncorrected data. Analyzing the corrected data is especially important because the incompleteness of the fossil record distorts our view of the history of life; for example, observed times of extinction always predate true times of extinction, unless fossil misidentifications or correlation errors occur. The observed temporal ranges of taxa are not only incomplete but also biased.

Results

As a baseline to compare with previous studies, we first analyzed the uncorrected data in both binned and pulsed boundary-crosser tabulations. We then analyzed the corrected data in the preferred pulsed model, and for comparison we also analyzed the corrected data by assuming continuous turnover. Our results are

Table 1. VAR(2) coefficients for all data sets, reported as point estimate \pm heteroskedasticity robust standard errors

Covariance	Lag	Coefficient	Uncorrected binned	Uncorrected pulsed	Corrected continuous	Corrected pulsed
Origination autocovariance	1	Φ_1^{oo}	0.60 \pm 0.10	0.59 \pm 0.21	-0.10 ± 0.13	0.05 ± 0.10
	2	Φ_2^{oo}	0.00 ± 0.08	-0.26 ± 0.10	-0.01 ± 0.13	-0.13 ± 0.07
Origination \rightarrow extinction covariance	1	Φ_1^{oe}	0.17 ± 0.13	0.08 ± 0.05	0.13 ± 0.13	0.07 ± 0.04
	2	Φ_2^{oe}	-0.07 ± 0.11	-0.02 ± 0.02	0.22 ± 0.12	-0.00 ± 0.03
Extinction \rightarrow origination covariance	1	Φ_1^{eo}	0.08 ± 0.07	0.36 ± 0.38	0.62 ± 0.15	1.37 ± 0.42
	2	Φ_2^{eo}	0.23 ± 0.05	1.16 ± 0.30	0.13 ± 0.13	1.49 ± 0.54
Extinction autocovariance	1	Φ_1^{ee}	0.37 ± 0.17	0.46 ± 0.16	0.06 ± 0.14	0.12 ± 0.12
	2	Φ_2^{ee}	0.05 ± 0.11	0.05 ± 0.10	-0.19 ± 0.15	0.01 ± 0.11

Numbers in boldface type are 5% significant values ($P < 0.05$). Φ components are as in Eq. 1. Uncorrected binned data are from Sepkoski’s compilation (3). Boundary-crosser data, from both uncorrected (pulsed) and corrected (continuous and pulsed) analyses, are as in figures 2, 3, and 4 in ref. 13.




Fig. 2. Correcting for the incompleteness of the fossil record causes autocovariance within origination, and within extinction, to disappear. Scatter plots of origination O_{i-1} in a given interval vs. origination in the following interval O_i are presented for Sepkoski's uncorrected (a) and corrected (c) data. Similar scatterplots of extinction are presented for uncorrected (b), and corrected (d) data. Boundary-crossers with pulsed turnover are used in all panels. Color coding corresponds to geologic time period, as in Fig. 1. Mean averages are indicated by black lines. The autocovariance present in the uncorrected data (a and b), manifest as the clumping of data points around the gray line of slope 1, disappears in the corrected data (c and d).

presented in Table 1, with coefficients reported as the point estimate \pm one standard error, with significant coefficients ($P < 0.05$) in boldface type.

Because several factors might have influenced our results, we also ran several additional regressions: we included t_i as another variable to account for varying interval duration, and we included a linear drift (detrended the data), to account for long-term secular decline. These results, presented in Table 2,

which is published as supporting information on the PNAS web site, agree within error with those in Table 1.

For the uncorrected data, in both binned and boundary-crosser tabulations, we find that per-genus origination and extinction rates have strong autocovariance, even while accounting for extinction-origination covariance (Fig. 2 a and b). This finding agrees with earlier results (4–7), which have formed the basis for the suggestion of inertia in the rates of origination and extinction; i.e., once origination (6, 7) or extinction (5–7) rates increase (decrease), they remain high (low). The only significant covariance is between rates of origination and the rates of extinction two intervals previous, or about 5–10 million years ($\Phi_2^{\text{co}} = 1.16 \pm 0.30$ for boundary-crossers). That is, the present origination rate depends on the extinction rate two intervals previous, and *not* on the rate in the immediately preceding interval (i.e., Φ_1^{co} is insignificant). Although calculated by a completely separate analysis, this result agrees with the peak at 10 million years in the cross-correlation analysis by Kirchner and Weil (6) that has been used to infer delayed recovery after extinction.

The picture drastically changes, however, when Foote's corrections for the incompleteness of the fossil record (13) are applied to Sepkoski's raw data. The autocovariances vanish entirely (Fig. 2 c and d). Note that the statistical result is *not* a negative autocovariance: it is not the case that a high rate of extinction presages a low rate in the next interval, for example. The temporal gap between extinction and subsequent origination also disappears entirely; no evidence for delayed recovery remains. In Foote's continuous model, the two-interval lag Φ_2^{co} becomes insignificant, replaced by a significant one-interval lag Φ_1^{co} . In Foote's pulsed model, both one-interval and two-interval lags contribute significantly.

Discussion

Our analysis suggests that the temporal gap between extinction and subsequent origination in the uncorrected data are an artifact of the smearing back of extinctions [Signor–Lipps effect (16)] and the smearing forward of originations [the Sppil–Rongis (6, 17) or Jaanusson (18, 19) effect] due to incomplete preservation. If a taxon's true last occurrence is missed because of incomplete preservation, its extinction will be mistakenly reported too early; conversely, originations will be misreported too late. However, if the fossil record were uniformly incomplete, then, as Kirchner and Weil point out (6), the Signor–Lipps and




Fig. 3. Episodic incompleteness of the fossil record can cause a rapid recovery from extinction to appear delayed. (a) Complete preservation: an extinction pulse at t_{i-1} followed immediately by complete recovery (origination pulse) at t_i . (b) Uniform, incomplete preservation: assuming an exponential distribution of waiting times between a fossil's apparent and true extinction (6), the pulse of extinction at t_{i-1} (dark purple) is smeared over t_{i-2} and t_{i-3} (lighter purple); the analogous argument applies to origination (blue). Nonetheless, the origination distribution's peak at t_i still immediately follows the extinction distribution's peak at t_{i-1} , leaving temporal correlations qualitatively unchanged relative to the case of complete preservation. (c) Episodic, incomplete preservation: preservability in the time interval t_i immediately after an extinction at t_{i-1} is often comparatively low (13), so that taxa that actually originated during t_i will be reported to have originated in t_{i+1} and t_{i+2} . This drop in preservability at t_i further decreases the reported origination rate at t_i (blue arrow) while concomitantly increasing the apparent rates in t_{i+1} and t_{i+2} (white arrows), creating the artifactual temporal lag/delay in time of recovery from extinction that appears in Sepkoski's data (13).

