

Positional cloning of the wheat vernalization gene *VRN1*

L. Yan*, A. Loukoianov, G. Tranquilli†, M. Helguera‡, T. Fahima§, and J. Dubcovsky*¶

Department of Agronomy and Range Science, University of California, Davis, CA 95616

Edited by Steven D. Tanksley, Cornell University, Ithaca, NY, and approved March 6, 2003 (received for review December 5, 2002)

Winter wheats require several weeks at low temperature to flower. This process, vernalization, is controlled mainly by the *VRN1* gene. Using 6,190 gametes, we found *VRN1* to be completely linked to MADS-box genes *AP1* and *AGLG1* in a 0.03-centimorgan interval flanked by genes *Cysteine* and *Cytochrome B5*. No additional genes were found between the last two genes in the 324-kb *Triticum monococcum* sequence or in the colinear regions in rice and sorghum. Wheat *AP1* and *AGLG1* genes were similar to *Arabidopsis* meristem identity genes *AP1* and *AGL2*, respectively. *AP1* transcription was regulated by vernalization in both apices and leaves, and the progressive increase of *AP1* transcription was consistent with the progressive effect of vernalization on flowering time. Vernalization was required for *AP1* transcription in apices and leaves in winter wheat but not in spring wheat. *AGLG1* transcripts were detected during spike differentiation but not in vernalized apices or leaves, suggesting that *AP1* acts upstream of *AGLG1*. No differences were detected between genotypes with different *VRN1* alleles in the *AP1* and *AGLG1* coding regions, but three independent deletions were found in the promoter region of *AP1*. These results suggest that *AP1* is a better candidate for *VRN1* than *AGLG1*. The epistatic interactions between vernalization genes *VRN1* and *VRN2* suggested a model in which *VRN2* would repress directly or indirectly the expression of *AP1*. A mutation in the promoter region of *AP1* would result in the lack of recognition of the repressor and in a dominant spring growth habit.

Winter wheats differ from spring wheats in their requirement for a long period at low temperatures to become competent to flower. This process, vernalization, prevents the damage of the cold-sensitive flowering meristem during the winter. *VRN1* and *VRN2* (unrelated to the genes with similar names in *Arabidopsis*) are the main genes involved in the vernalization response in diploid wheat *Triticum monococcum* (1, 2). However, most of the variation in vernalization requirement in the economically important polyploid species of wheat is controlled by the *VRN1* locus (2, 3). This gene is critical in polyploid wheats for their adaptation to autumn sowing and divides wheat varieties into the winter and spring market classes.

The *VRN1* gene has been mapped in colinear regions of the long arm of chromosomes 5A (1, 3, 4), 5B (5, 6), and 5D (3). This region of wheat chromosome 5 is colinear with a region from rice chromosome 3 that includes the *HD-6* quantitative trait locus for heading date (7). However, it was recently demonstrated that *VRN1* and *HD-6* are different genes (8).

Despite the progress made in the elucidation of the vernalization pathway in *Arabidopsis*, little progress has been made in the characterization of wheat vernalization genes. The two main genes involved in the vernalization pathway in *Arabidopsis*, *FRI* and *FLC* (9–11), have no clear homologues in the complete draft sequences of the rice genome (12). This finding may not be surprising, considering that rice is a subtropical species that has no vernalization requirement. Because no clear orthologues of the *Arabidopsis* vernalization genes were found in rice or among the wheat or barley ESTs, a map-based cloning project for the wheat *VRN1* gene was initiated in our laboratory.

Chromosome walking in wheat is not a trivial exercise because of the large size of its genomes (5,600 Mb per *T. monococcum*

haploid genome) and the abundance of repetitive elements (13, 14). To minimize the probability that these repetitive elements would stop the chromosome walking, simultaneous efforts were initiated in the orthologous regions in rice, sorghum, and wheat. The initial sequencing of rice, sorghum, and wheat bacterial artificial chromosomes (BACs) selected with restriction fragment length polymorphism marker WG644 [0.1 centimorgan (cM) from *VRN1*] showed good microcolinearity among these genera (14–16). The low gene density observed in the wheat region and the large ratio of physical-to-genetic distances (14) suggested that large mapping populations and comparative physical maps would be necessary for a successful positional cloning of *VRN1*.

In this article we report the construction of detailed genetic and physical maps of *VRN1* in diploid wheat and the comparison of the sequences from selected BACs from these contigs with the sequences of colinear regions in rice and sorghum. We also present the expression patterns of two MADS-box genes completely linked to *VRN1* and discuss the evidence supporting one of these genes as a strong candidate gene for *VRN1*.

Materials and Methods

Mapping Population. The high-density map was based on 3,095 F_2 plants from the cross between *T. monococcum* ssp. *aegilopoides* accessions G2528 (spring, *Vrn1*) with G1777 (winter, *vrn1*). These two lines have the same dominant allele at the *VRN2* locus and therefore, plants from this cross segregate only for *VRN1* in a clear 3:1 ratio (1, 2).

Plants were grown in a greenhouse at 20–25°C without vernalization and under long photoperiod (16-h light). Under these conditions, winter plants flowered 1–2 months later than spring plants. All F_2 plants were analyzed for molecular markers flanking *VRN1*, and progeny tests were performed for plants showing recombination between these markers. The 20–25 individual F_3 plants from each progeny test were characterized with molecular markers flanking the crossover to confirm that the observed segregation in growth habit was determined by variation at the *VRN1* locus. G2528 and G1777 were included as controls in each progeny test.

Procedures for genomic DNA extraction, Southern blots, and hybridizations have been described (17). The first 500 F_2 plants were screened with flanking restriction fragment length polymorphism markers CDO708 and WG644 (1), which were later replaced by closer PCR markers to screen the complete mapping

This paper was submitted directly (Track II) to the PNAS office.

Abbreviations: BAC, bacterial artificial chromosome; cM, centimorgan.

Data deposition: The sequences reported in this paper have been deposited in the GenBank database (accession nos. AY188331, AY188332, AY188333, and AY188330).

*L.Y. and J.D. contributed equally to this work.

†Present address: Instituto de Recursos Biológicos, Instituto Nacional de Tecnología Agropecuaria, Villa Udaondo, 1712 Castelar, Buenos Aires, Argentina.

‡Present address: Estación Experimental Agropecuaria, Instituto Nacional de Tecnología Agropecuaria, 2580 Marcos Juárez, Córdoba, Argentina.

§Present address: Institute of Evolution, University of Haifa, Mount Carmel, Haifa 31905, Israel.

¶To whom correspondence should be addressed. E-mail: jdubcovsky@ucdavis.edu.




Fig. 1. (A) Genetic map of the *VRN1* region on chromosome 5A^m of *T. monococcum*. Genetic distances are in cM (6,190 gametes). (B–D) Physical maps of the colinear *VRN1* regions in rice, sorghum, and wheat. Regions indicated in red have been sequenced. Double dot lines indicate gaps in the current physical maps. (B) Sequence of the colinear region in rice chromosome 3. (C) *S. bicolor* BACs 170F8 (AF503433) and 17E12 (AY188330). (D) *T. monococcum* physical map. BAC clones order from left to right is: 49116, **115G1**, 136F13, 133P9, **116F2**, 89E14, 160C18, 491M20, 328O3, **609E6**, 393O11, **719C13**, 454P4, 54K21, 579P2, 601A24, **231A16**, 638J12, 52F19, 242A12, 668L22, 539M19, and 309P20 (bold indicates sequenced BACs). Black dots indicate validation of BAC connections by hybridization. (E) Gene structure of two MADS-box genes completely linked to the *VRN1* gene (AY188331, AY188333). Bars represent exons. (F) Sequence comparison of the *AP1* promoter regions from genotypes carrying the *Vrn1* and *vrn1* alleles and from two *T. monococcum* accessions with additional deletions. The last two accessions are spring but their genotype has not been determined yet. Numbers indicate distances from the start codon. A putative MADS-box protein-binding site (CARG-box) is indicated in green.

population. Additional markers were developed for the eight genes present between the PCR markers (*Appendix 1*, which is published as supporting information on the PNAS web site, www.pnas.org).

Contig Construction and BAC Sequencing. High-density filters for the BAC libraries from *T. monococcum* accession DV92 (18), *Oryza sativa* var. Nipponbare (19), and *Sorghum bicolor* (20) were screened with segments from the different genes indicated in Fig. 1. Contigs were assembled by using *Hind*III fingerprinting and confirmed by hybridization of BAC ends obtained by plasmid rescue, inverse PCR (20), or BAC sequencing. Restriction maps using single and double digestions with eight-cutter restriction enzymes, pulsed-field electrophoresis, and hybridization of the Southern blots with different genes were used to order genes within the BACs, select the fragment sequenced from the sorghum BAC, and confirm the assembly results from the BAC sequencing. Shotgun libraries for BAC sequencing were constructed as described (15). Complete *T. monococcum* BACs 609E6, 719C13, and 231A16 and a 24-kb fragment from sorghum BAC 17E12 were sequenced. Genes were identified by a combination of comparative genomic analysis, BLAST searches, and gene-finding programs (15).

Phylogenetic Analysis. A phylogenetic study was performed by using the two wheat MADS-box genes found in this study and 24 additional MADS-box genes (*Appendix 2*, which is published as supporting information on the PNAS web site). Phylogenetic trees were generated from the CLUSTALW sequence alignments of the complete proteins by using multiple distance- and parsimony-based methods available in the MEGA 2.1 software package

(21). Distances between each pair of proteins were calculated and a tree was constructed by using the neighbor-joining algorithm. The consensus tree and the confidence values for the nodes were calculated by using 1,000 bootstraps (MEGA 2.1).

RT-PCR and Quantitative PCR. RNA from leaves, undifferentiated apices, and young spikes was extracted by using the Trizol method (Invitrogen). RT-PCR procedures were performed as described (22). Quantitative PCR experiments were performed in an Applied Biosystems ABI 7700 by using three TaqMan systems for *T. monococcum AP1* and for actin and ubiquitin as endogenous controls (*Appendix 3*, which is published as supporting information on the PNAS web site). The $2^{-\Delta\Delta C_T}$ method (23) was used to normalize and calibrate the C_T values of wheat *AP1* relative to the endogenous controls. For the vernalization time course, RNA was extracted from the youngest fully expanded leaf of five winter *T. monococcum* plants (1 month old) immediately before moving the plants into the cold room, and then after 2, 4, and 6 weeks of vernalization (4°C). The last sample was collected 2 weeks after moving plants to the greenhouse (20°C). Plants kept in the greenhouse were sampled as controls at each time point simultaneously with the plants from the cold room (five plants per time point).

Results

Marker Development. In the initial genetic map (1) the *VRN1* gene was flanked on the distal side by WG644 and on the proximal side by CDO708. These markers were used as anchor points to the rice genome sequence to find additional markers.

Distal region. WG644 was previously used to identify rice BAC 3615 that included *GENE1* at its proximal end (15). BLASTN

searches of the different rice genome projects using *GENE1* and the end sequence of BAC 3615 (AY013245) identified the connected contig CL013482.168 (12). Two additional genes, *Phytochelatase synthetase* (*PCS*, *Zea mays* AAF24189.1) and *Cytochrome B5* (*CYB5*, NP_173958.1), were discovered and annotated in this contig. These genes were mapped in wheat by restriction fragment length polymorphism (*Appendix 1*).

Proximal region. Restriction fragment length polymorphism marker CDO708 was mapped 0.9 cM proximal to *VRN1* in the *T. monococcum* map. The sequence of this clone (AY245605) showed a high homology to a putative RNA-binding protein (AAL58954.1) located in rice BAC AC091811. The end of this rice BAC also included gene *MTK4* (putative protein kinase tousel, AAL58952.1), which was converted into a PCR marker and was mapped in wheat (Fig. 1). Rice BAC sequence AC091811 was then connected through contigs CL039395.93, CL039395.83, and CL018222.111.1 (12) to rice BAC sequences AC092556 and AF377947. BAC sequence AC092556 included a *transcriptional adaptor* gene (*ADA2*, AJ430205) that was mapped in *T. monococcum* 0.5 cM from the *VRN1* gene (Fig. 1). The last rice BAC sequence, AF377947, included genes *Phytochrome C* (*PHY-C*, AAM34402.1), *cysteine proteinase* (*CYS*, AAM34401.1) and MADS-box genes AAM34398.1 and AAM34397.1, designated hereafter *API* and *AGL1* (*agamous*-like gene from grasses). The rice proximal region included 318 kb of contiguous sequence.

High-Density Genetic Maps of the *VRN1* Region. The PCR markers developed for *GENE1* and *MTK4* were used to screen 6,190 chromosomes for recombination. Fifty-one recombinant events were detected, and those plants were further characterized by using molecular markers for all of the genes present between these two markers in rice (Fig. 1; *Appendix 1*). Progeny tests were performed for 32 of the 51 F₂ plants to determine the *VRN1* genotype of the parental F₂ plants. Based on the mapping information, the *VRN1* locus was completely linked to *API* and *AGL1*.

On the proximal side, genes *PHY-C* and *CYS* flanked the *VRN1* locus. The last two genes were completely linked to each other and separated from *VRN1* by a single crossover (Fig. 1; *Appendix 1*). On the distal side, the *CYB5* gene was also separated from *VRN1* by a single crossover. Comparison of genotypic and phenotypic data from all of the F₃ plants used in the 30 progeny tests confirmed that the observed segregation in growth habit was determined by variation at the *VRN1* locus. Unvernalized plants homozygous for the G1777 *API* allele flowered 1–2 months later than G2528, whereas the other plants flowered only 1 week before or after the G2528 control. These results confirmed the simple Mendelian segregation for vernalization requirement in this cross (1).

Physical Maps. Distal contig. Genes *CYB5* and *GENE1* were used to screen the BAC libraries from *T. monococcum*, rice, and sorghum. *T. monococcum* BAC clone 609E6 selected with the *CYB5* gene was connected to previously sequenced 116F2 (AF459639) by four BACs (Fig. 1). The *PCS* gene hybridized with two fragments from BAC 609E06 (*PCS1* and *PCS2*, Fig. 1) whereas only one *PCS* copy was found in the colinear region in rice. No single-copy probes were found in BACs 609E6 or in the unique *Hind*III fragments from the most proximal BAC 393O11 to continue the chromosome walking toward the proximal region. **Proximal contig.** Screening of the *T. monococcum* BAC library with *PHY-C*, *CYS*, *API*, and *AGL1* yielded 12 BACs organized in two contigs. The largest contig included eight BACs that hybridized with genes *PHY-C*, *CYS*, and *API*. The four additional BACs hybridized only with the *AGL1* gene (Fig. 1). The location of the *AGL1* contig within the physical map was determined by the complete linkage between the single-copy

genes *AGL1* and *API* and the proximal location of *AGL1* relative to the single-copy gene *CYB5*. No additional single-copy probes were found to close the gaps flanking the *AGL1* contig.

The proximal gap between *AGL1* and *API* was covered by the current rice sequence. However, the distal gap between *CYB5* and *AGL1* was also present in the different rice genome sequencing projects. The screening of the Nipponbare BAC libraries with probe *CYB5* failed to extend the rice region because of the presence of a gap in the current rice physical maps. Fortunately, sorghum BAC 17E12 included *GENE1*, *PCS1*, *PCS2*, and *CYB5* genes from the distal contig, and *AGL1*, *API*, and *CYS* genes from the proximal contig, bridging the gap present in the rice and wheat contigs (Fig. 1). A restriction map of sorghum BAC 17E12 (*Appendix 4*, which is published as supporting information on the PNAS web site) indicated that the sorghum genes were in the same order as that previously found in rice and wheat and that a 24-kb *SwaI*–*SwaI* restriction fragment spanned the region of the rice and wheat gap between *CYB5* and *AGL1*.

Sequence Analysis. Annotated sequences from the three *T. monococcum* BACs (AY188331, AY188332, and AY188333) and the partial sequence of the sorghum BAC 17E12 (AY188330) were deposited in GenBank. Including BACs 115G01 and 116F02 (AF459639), a total of 550 kb was sequenced. Multiple retrotransposons organized in up to four layers of nested elements were the most abundant features, similar to wheat regions analyzed before (13, 14). Annotated retrotransposons and other repetitive elements accounted for 78.4% of the sequence, whereas genes represented only 8.5% of the total. The genes detected in this sequence were in the same order as the ones present in the corresponding regions in rice and sorghum, indicating an almost perfect microcolinearity. The only exception was the duplication of the *PCS* gene in sorghum and wheat relative to the presence of a single *PCS* gene in the colinear rice region (Fig. 1).

No additional genes were found in the rice sequence between the two MADS-box genes corresponding to one of the two gaps in the wheat physical map. These two genes were also adjacent in sorghum (Fig. 1; *Appendix 4*). Similarly, no new genes were found between *CYB5* and *AGL1* in the sequence of the 24-kb *SwaI*–*SwaI* restriction fragment from sorghum BAC 17E12 (AY188330) that covered the other gap in the wheat physical map. The four genes present in the sorghum sequence were in the same order and orientation as that previously found in rice and wheat (*Appendix 4*).

The absence of new genes in the colinear regions of rice and sorghum, together with the excellent microcolinearity detected in this region, suggested that it would be unlikely to find additional genes in the current gaps of the wheat physical map. This assumption was also supported by the absence of any new gene in the 324 kb of wheat sequence flanking these gaps. The presence of almost uninterrupted series of nested retrotransposons flanking the gaps also explained the failure to find single-copy probes to close the two gaps.

Classification of the Two MADS-Box Genes. The *API* and *AGL1* proteins have MADS-box and K domains characteristic of homeotic genes involved in the flowering process and similar exon structure (Fig. 1; *Appendix 5*, which is published as supporting information on the PNAS web site) (24). The consensus tree for 26 plant MADS-box proteins (*Appendix 2*) showed that the closest proteins to wheat *API* and *AGL1* belonged to the SQUAMOSA (bootstrap 97) and AGL2 groups (bootstrap 95), respectively.

The closest *Arabidopsis* MADS-box proteins to wheat *API* were the proteins coded by the three related meristem identity genes *API*, *CAL*, and *FUL* (*Appendix 2*). Two separate clusters




Fig. 2. (A) RT-PCR experiment using *T. monococcum* G3116 (winter growth habit) and *API*-, *AGLG1*-, and actin-specific primers. The PCRs for the three genes were performed by using the same cDNA samples. Lanes 1–5 indicate leaves. Lane 1, before vernalization; lanes 2–4, 2, 4, and 6 weeks of vernalization, respectively; lane 5, 2 weeks after vernalized plants were returned to the greenhouse; lane 6, unvernallized apices; lane 7, 6-week vernalized apices; and lane 8, young spikes. (B) *API* transcription levels in leaves relative to actin measured by quantitative PCR. Lanes 1–5, leaves from plants at the same vernalization stage as samples 1–5 in A. Units are values linearized by using the $2^{-(C_T - \Delta\Delta C_T)}$ method, where C_T is the threshold cycle.

were observed in the SQUAMOSA group dividing the *Arabidopsis* and grass proteins. A similar separation between the monocot and dicot proteins was found in more detailed studies of this group (25). The *API* protein from *T. monococcum* was 98.4% similar to previously described *Triticum aestivum* *WAPI*, formerly *TaMADS11* (26, 27) and 96.0% similar to barley *BM5* (28). These two putative orthologous genes were described in papers characterizing the MADS-box family in wheat and barley but were not mapped or associated with the *VRN1* gene.

The wheat *AGLG1* protein was clustered with members of the *AGL2* subgroup and was closely related with the rice *AGLG1* orthologue and with rice *OsMADS5*, *OsMADS1*, and barley *BM7* proteins (bootstrap 87, *Appendix 2*).

Expression Profiles. No *API* transcripts were detected in apices from unvernallized plants of *T. monococcum* with strong winter growth habit (G3116) even after 10 months in the greenhouse under long-day conditions. However, *API* transcription was detected in the apices of plants from the same genotype after 6 weeks of vernalization (Fig. 2A, lanes 6 and 7). The same result was obtained in three independent experiments. These apices were morphologically at vegetative stage zero according to the developmental scale of Gardner *et al.* (29). In *T. monococcum* accessions with spring growth habit, *API* transcripts were observed in the apices without previous vernalization.

Transcripts of *API* were also detected in the leaves, as reported for *WAPI* (26) and *BM5* (28). A quantitative PCR experiment using the endogenous controls actin (Fig. 2B) and ubiquitin (*Appendix 3*) demonstrated that transcription of *API* in the leaves of the winter genotypes was also regulated by vernalization. The abundance of *API* transcripts started to increase after the first 2 weeks of vernalization and continued to increase during the 4 additional weeks of the vernalization process (Fig. 2B). *API* transcripts were also present in the leaves from

vernallized plants 2 weeks after their transfer to the greenhouse. At this stage, *API* transcripts were detected in young and old green leaves (*Appendix 3*). Control plants kept in the greenhouse showed very low levels of *API* transcription during the 8 weeks of the experiment (Fig. 2B). In the genotypes with a spring growth habit, *API* transcripts were observed in the leaves of unvernallized plants that were initiating the transition to flowering.

AGLG1 transcripts were detected only in young spikes (Fig. 2A, lane 8) but were not observed in the same cDNA samples from apices after 6 weeks of vernalization where the *API* transcripts were already present (Fig. 2A). This finding indicates that *AGLG1* transcription is initiated later than *API*. Transcripts from *AGLG1* were not detected in the leaves (Fig. 2A).

The expression results, together with the known role of the *API* homologues in *Arabidopsis* as meristem identity genes, suggested that *API* was a better candidate gene for *VRN1* than *AGLG1*.

Allelic Variation. Four *API* genes were sequenced from *T. monococcum* accessions G1777, G3116, and DV92 carrying the *vrn1* allele and G2528 carrying the *Vrn1* allele (1, 2). The predicted proteins from DV92 and G2528 were identical and differed from the predicted proteins from G3116 and G1777 by a single amino acid (*Appendix 5*).

Analysis of the 1,024-bp region upstream from the *API* start codon and up to the insertion point of a large repetitive element (AY188331) showed the presence of five polymorphic sites. Two of them differentiated G2528 from the three accessions carrying the *vrn1* allele for winter growth habit. One was a 1-bp insertion located 728 bp upstream from the start codon, and the other one was a 20-bp deletion located 176 bp upstream from the start codon (Fig. 1; *Appendix 5*). No difference were detected in the first 600 bp of the *API* 3' region between the *vrn1* and *Vrn1* alleles.

A PCR screening of a collection of cultivated *T. monococcum* accessions with primers flanking the 20-bp deletion region revealed the presence of deletions of different sizes in agarose gels (*Appendix 5*). Sequencing of these lines showed the presence of two new deletions that overlapped with the 20-bp deletion from G2528. These new deletions included a putative MADS-box protein-binding site adjacent to the 20-bp deletion (Fig. 1; *Appendix 5*).

No DNA differences were detected between accessions DV92 (*vrn1*) and G2528 (*Vrn1*) in the coding region, or the 5' (365 bp) and 3' (583 bp) UTRs of the *AGLG1* gene.

Discussion

Genetic and Physical Maps of the *VRN1* Region. Only eight genes were found in the 556-kb sequence from the *T. monococcum* *VRN1* region, resulting in an estimated gene density of one gene per 70 kb. The low gene density observed in this region was paralleled by a high ratio between physical and genetic distances. Excluding the two gaps in the physical map, a minimum ratio of 6,250 kb·cM⁻¹ was estimated for the region between WG644 and *PHY-C*. This value is two times larger than the average genome-wide estimate of 3,000 kb·cM⁻¹ (30) and four times larger than the 1,400 kb·cM⁻¹ reported for the telomeric region of chromosome 1A (31). Previous cytogenetic studies demonstrated that recombination in the wheat chromosomes decreases exponentially with distance from the telomere (32, 33), predicting an increase of the ratio between physical and genetic distance in the same direction. The region studied here is located between the breakpoints in deletion lines 5AL-6 (FL 0.68) and 5AL-17 (FL 0.78), in a more proximal location than regions used before to estimate ratios between physical and genetic distances in wheat. This result suggests that positional cloning projects in the proximal regions of wheat will be difficult and would greatly

benefit from the use of the rice genomic sequence to jump over large blocks of repetitive elements.

Despite the low recombination rate found in this region, the large number of evaluated gametes was sufficient to find crossovers between most of the genes or at least between pairs of adjacent genes. This detailed genetic study showed that the variation in growth habit determined by the *VRN1* gene was completely linked to only two genes. Although the possibility that additional genes would be found in the two current gaps and unsequenced regions of our *T. monococcum* physical maps cannot be ruled out, this seems unlikely based on the comparative studies with rice and sorghum and the absence of any additional genes in the 324-kb wheat sequence between *CYB5* and *CYS*.

The genetic data reduced the problem of the identification of *VRN1* to the question of which of the two MADS-box genes was the correct candidate. However, because no recombination was found between *AGLG1* and *API* it was not possible to answer this question based on the available genetic results. Therefore, the relationship between *AGLG1* and *API* with MADS-box genes from other species was established as a first step to predict their function from the known function of the related genes.

Phylogenetic Relationships of the *VRN1* Candidate Genes. The similarity between the wheat *API* gene and the *Arabidopsis* meristem identity genes *API*, *CAL*, and *FUL* provided an indication that the wheat *API* gene was a good candidate for *VRN1*. These *Arabidopsis* genes are expressed in the apices and are required for the transition between the vegetative and reproductive phases (34). The triple *Arabidopsis* mutant *ap1-cal-ful* never flowers under standard growing conditions (34). In wheat, the *VRN1* gene is also responsible, directly or indirectly, for the transition between vegetative and reproductive apices. This transition is greatly accelerated by vernalization in the wheat plants carrying the *vrn1* allele for winter growth habit. Therefore, it is reasonable to speculate that the sequence similarity between the wheat *API* gene and the *Arabidopsis* meristem identity genes may indicate similar functions. An evolutionary change in the promoter region of *API* may be sufficient to explain the regulation of *API* by vernalization in wheat (see model below).

The close relationship of wheat *AGLG1* to members of the AGL2 subgroup suggested that *AGLG1* was a less likely candidate for *VRN1* than *API* because transcripts from genes included in this group are usually not observed in the apices in the vegetative phase (25). Expression of *Arabidopsis* *AGL2*, *AGL4*, and *AGL9* begins after the onset of expression of floral meristem identity genes but before the activation of floral organ identity genes, suggesting that members of the AGL2 clade may act as intermediaries between the meristem identity genes and the organ identity genes (35–37). This seems to be valid also for *OsMADS1*, which is more closely related to *AGLG1* than the *Arabidopsis* members of the AGL2 clade. *In situ* hybridization experiments of young rice inflorescences with *OsMADS1* showed strong hybridization signals in flower primordia but not in other tissues (38).

If the functions of wheat *API* and *AGLG1* were similar to the function of the related genes from *Arabidopsis*, the initiation of transcription of *API* should precede the initiation of transcription of *AGLG1* in wheat.

Transcription Profiles of the *VRN1* Candidate Genes. RT-PCR experiments using RNA samples from vernalized apices showed transcription of *API* but not of *AGLG1* (Fig. 2A), indicating that transcription of *AGLG1* occurs after the initiation of transcription of *API*. The similar timing and order of transcription suggests that the wheat genes might perform similar functions to the related *Arabidopsis* genes.




Fig. 3. A model of the regulation of flowering initiation by vernalization in wheat.

It could be argued that any gene in the flowering regulatory pathway would be up-regulated by the initiation of flowering caused by the vernalization process. However, the up-regulation of *API* transcription in the leaves by vernalization (Fig. 2B) indicated a more direct role of the vernalization pathway in the regulation of wheat *API* gene. Four additional characteristics of the transcription profile of *API* paralleled the predicted expression of a vernalization gene. First, vernalization was required to initiate *API* transcription in the plants with winter growth habit but not in the plants with spring growth habit. Second, *API* transcription was initiated after only 2 weeks in the cold room, and a minimum of 2 weeks of vernalization is required by many winter wheat varieties to produce any significant acceleration of flowering (39). Third, the progressive increase of *API* transcripts after the second week of vernalization (Fig. 2B) is consistent with the progressive effect of the length of the vernalization period in the acceleration of flowering time (39). Finally, a high level of *API* transcripts was observed after the plants were moved from the cold to room temperature, indicating that *API* is not just a cold stress-induced gene.

Allelic Variation. No differences were found in the *AGLG1* coding region or in its 5' and 3' regions between *T. monococcum* accessions G2528 (*Vrn1*) and DV92 (*vrn1*), confirming that *AGLG1* was not a good candidate to explain the observed differences in growth habit.

Although no differences were detected in the *API* coding sequences and 3' region, the spring and winter accessions differed in their promoter sequence. The first 600 bp upstream from the start codon were identical among the four genotypes analyzed in this study except for a 20-bp deletion located close to the start of transcription and adjacent to a putative MADS-box protein binding site (CARG-box) in G2528 (ref. 40; Appendix 5). Two additional overlapping deletions were discovered in the same region of the promoter in spring accessions of cultivated *T. monococcum* (Fig. 1). The presence of a putative CARG-box in this region suggests the possibility that a transacting factor may bind to this site and repress *API* transcription until vernalization occurs. This is similar to the case for *FLC* in *Arabidopsis*, which was recently shown to bind to MADS-box gene *SOC1* and repress its transcription before vernalization (41).

A Model for the Regulation of Flowering by Vernalization in Wheat. The results presented in this study can be included in an integrated model (Fig. 3) based on the known epistatic interactions between *VRN1* and *VRN2* (2) and the available information

about the evolution of the vernalization requirement in the Triticeae. The significant epistatic interactions observed between *VRN1* and *VRN2* indicate that these two genes act in the same pathway (2). According to the model presented here (Fig. 3), *VRN2* codes for a dominant repressor of flowering that acts directly or indirectly to repress *VRN1*. As the vernalization process reduces the abundance of the *VRN2* gene product, *VRN1* transcription gradually increases, leading to the competence to flower (Fig. 3 *Middle*).

The growth habit of plants homozygous for the recessive *vrn2* allele for spring growth habit (Fig. 3 *Top*) is independent of variation at the *VRN1* locus (2). According to this model, the *vrn2* allele represents a null or defective repressor that cannot interact with the *VRN1* promoter. Therefore, variation in the promoter of the *VRN1* gene would have no effect on flowering time in homozygous *vrn2* plants. This can be illustrated by the expression pattern of *API* in *T. monococcum* DV92 (*vrn1 vrn2*). In this genotype, the initiation of *API* transcription in leaves and apices did not require vernalization despite the presence of a recessive *vrn1* allele. This result indicated that the *VRN1* gene acts downstream of *VRN2* (Fig. 3).

Conversely, plants homozygous for the *Vrn1* allele for spring growth habit showed no significant effects of the *VRN2* gene on flowering time (2). According to the model in Fig. 3 *Bottom*, the *VRN2* repressor will have no effect on flowering in genotypes carrying the *Vrn1* allele because of the lack of the recognition site in the *VRN1* promoter region. This part of the model can be used to explain the *API* expression profile of G2528 (*Vrn1 Vrn2*). In this genotype, transcription of *API* in leaves and apices is initiated without a requirement for vernalization despite the presence of an active *VRN2* repressor. This finding suggested that the active repressor could not interact with the G2528 *API* promoter region, possibly because of the presence of the 20-bp deletion.

This model also provides an explanation for the parallel

evolution of *VRN1* spring alleles in three different Triticeae lineages. A vernalization gene with a dominant spring growth habit has been mapped in the same map location in diploid wheat (1), barley (42), and rye (43). Most of the wild Triticeae have a winter growth habit, suggesting that the recessive *vrn1* allele is the ancestral character (44–46). This notion is also supported by the fact that it is unlikely that a vernalization requirement would be developed independently at the same locus in the three different lineages from an ancestral spring genotype. According to the model presented here, independent mutations in the promoter regions of winter wheat, barley, and rye genotypes have resulted in the loss of the recognition site of the *VRN2* repressor (or an intermediate gene) and therefore, in a dominant spring growth habit (*Vrn1* allele). Because this is a loss rather than a gain of a new function it is easier to explain its recurrent occurrence in the different Triticeae lineages.

In summary, this article presents the delimitation of the candidate genes for *Vrn1* to *API* and *AGL1* by a high-density genetic map, and the identification of *API* as the most likely candidate based on its similar sequence to meristem identity genes, its transcription profile, and its natural allelic variation. A model is presented to integrate the results from this study with the previous knowledge about the epistatic interactions between vernalization genes and the evolution of vernalization in the Triticeae. Confirmation of the hypotheses generated by the model presented here will contribute to unraveling the complex net of relationships responsible for the regulation of heading date in the temperate cereals.

We thank Drs. P. Gepts and M. Soria for their valuable suggestions; R. Shao, A. Sanchez, S. Olmos, and C. Busso for excellent technical assistance; and Dr. W. Ramakrishna and J. Bennetzen for screening the sorghum BAC library. This work was supported by U.S. Department of Agriculture–National Research Initiative Grant 2000-1678 and National Science Foundation Grant PGRP-99-75793.

- Dubcovsky, J., Lijavetzky, D., Appendino, L. & Tranquilli, G. (1998) *Theor. Appl. Genet.* **97**, 968–975.
- Tranquilli, G. E. & Dubcovsky, J. (1999) *J. Hered.* **91**, 304–306.
- Law, C. N., Worland, A. J. & Giorgi, B. (1975) *Heredity* **36**, 49–58.
- Galiba, G., Quarrie, S. A., Sutka, J. & Morgounov, A. (1995) *Theor. Appl. Genet.* **90**, 1174–1179.
- Iwaki, K., Nishida, J., Yanagisawa, T., Yoshida, H. & Kato, K. (2002) *Theor. Appl. Genet.* **104**, 571–576.
- Barrett, B., Bayram, M. & Kidwell, K. (2002) *Plant Breed.* **121**, 400–406.
- Kato, K., Miura, H. & Sawada, S. (1999) *Genome* **42**, 204–209.
- Kato, K., Kidou, S., Miura, H. & Sawada, S. (2002) *Theor. Appl. Genet.* **104**, 1071–1077.
- Michaels, S. D. & Amasino, R. M. (1999) *Plant Cell* **11**, 949–956.
- Sheldon, C. C., Rouse, D. T., Finnegan, E. J., Peacock, W. J. & Dennis, E. S. (2000) *Proc. Natl. Acad. Sci. USA* **97**, 3753–3758.
- Johanson, U., West, J., Lister, C., Michaels, S., Amasino, R. & Dean, C. (2000) *Science* **290**, 344–347.
- Goff, S. A., Ricke, D., Lan, T. H., Presting, G., Wang, R. L., Dunn, M., Glazebrook, J., Sessions, A., Oeller, P., Varma, H., et al. (2002) *Science* **296**, 92–100.
- Wicker, T., Stein, N., Albar, L., Feuillet, C., Schlagenhauf, E. & Keller, B. (2001) *Plant J.* **26**, 307–316.
- SanMiguel, P., Ramakrishna, W., Bennetzen, J. L., Busso, C. S. & Dubcovsky, J. (2002) *Funct. Integr. Genomics* **2**, 70–80.
- Dubcovsky, J., Ramakrishna, W., SanMiguel, P., Busso, C., Yan, L., Shiloff, B. & Bennetzen, J. (2001) *Plant Physiol.* **125**, 1342–1353.
- Ramakrishna, W., Dubcovsky, J., Park, Y. J., Busso, C. S., Emberton, J., SanMiguel, P. & Bennetzen, J. L. (2002) *Genetics* **162**, 1389–1400.
- Dubcovsky, J., Galvez, A. F. & Dvorak, J. (1994) *Theor. Appl. Genet.* **87**, 957–964.
- Lijavetzky, D., Muzzi, G., Wicker, T., Keller, B., Wing, R. & Dubcovsky, J. (1999) *Genome* **42**, 1176–1182.
- Zhang, H.-B., Choi, S., Woo, S. S., Li, Z. & Wing, R. A. (1996) *Mol. Breeding* **2**, 11–24.
- Woo, S. S., Jiang, J., Gill, B. S., Paterson, A. H. & Wing, R. A. (1994) *Nucleic Acids Res.* **22**, 4922–4931.
- Kumar, S., Tamura, K. & Nei, M. (1994) *Comput. Appl. Biosci.* **10**, 189–191.
- Yan, L., Echenique, V., Busso, C., SanMiguel, P., Ramakrishna, W., Bennetzen, J. L., Harrington, S. & Dubcovsky, J. (2002) *Mol. Genet. Genomics* **268**, 488–499.
- Livak, K. J. & Schmittgen, T. D. (2001) *Methods* **25**, 402–408.
- Ng, M. & Yanofsky, M. F. (2001) *Nat. Rev. Genet.* **2**, 186–195.
- Johansen, B., Pedersen, L. B., Skipper, M. & Frederiksen, S. (2002) *Mol. Phylogenet. Evol.* **23**, 458–480.
- Murai, K., Murai, R. & Ogihara, Y. (1997) *Genes Genet. Syst.* **72**, 317–321.
- Murai, K., Takumi, S., Koga, H. & Ogihara, Y. (2002) *Plant J.* **29**, 169–181.
- Schmitz, J., Franzen, R., Ngyuen, T. H., Garcia-Maroto, F., Pozzi, C., Salamini, F. & Rohde, W. (2000) *Plant Mol. Biol.* **42**, 899–913.
- Gardner, J. S., Hess, W. M. & Trione, E. J. (1985) *Am. J. Bot.* **72**, 548–559.
- Bennett, M. D. & Smith, J. B. (1991) *Philos. Trans. R. Soc. London B* **334**, 309–345.
- Stein, N., Feuillet, C., Wicker, T., Schlagenhauf, E. & Keller, B. (2000) *Proc. Natl. Acad. Sci. USA* **97**, 13436–13441.
- Dvorak, J. & Chen, K.-C. (1984) *Genetics* **106**, 325–333.
- Lukaszewski, A. J. & Curtis, C. A. (1993) *Theor. Appl. Genet.* **84**, 121–127.
- Ferrandiz, C., Gu, Q., Martienssen, R. & Yanofsky, M. F. (2000) *Development (Cambridge, U.K.)* **127**, 725–734.
- Flanagan, C. A. & Ma, H. (1994) *Plant Mol. Biol.* **26**, 581–595.
- Savidge, B., Rounsley, S. D. & Yanofsky, M. F. (1995) *Plant Cell* **7**, 721–733.
- Mandel, M. A. & Yanofsky, M. F. (1998) *Sex Plant Reprod.* **11**, 22–28.
- Chung, Y. Y., Kim, S. R., Finkel, D., Yanofsky, M. F. & An, G. H. (1994) *Plant Mol. Biol.* **26**, 657–665.
- Limin, A. E. & Fowler, D. B. (2002) *Ann. Bot. (London)* **89**, 579–585.
- Tilly, J. J., Allen, D. W. & Jack, T. (1998) *Development (Cambridge, U.K.)* **125**, 1647–1657.
- Hepworth, S. R., Valverde, F., Ravenscroft, D., Mouradov, A. & Coupland, G. (2002) *EMBO J.* **21**, 4327–4337.
- Laurie, D. A., Pratchett, N., Bezant, J. H. & Snape, J. W. (1995) *Genome* **38**, 575–585.
- Plaschke, J., Börner, A., Xie, D. X., Koeber, R. M. D., Schlegel, R. & Gale, M. D. (1993) *Theor. Appl. Genet.* **85**, 1049–1054.
- Kihara, H. & Tanaka, M. (1958) *Preslia* **30**, 241–251.
- Halloran, G. M. (1967) *Genetics* **57**, 401–407.
- Goncharov, N. P. (1998) *Euphytica* **100**, 371–376.