

Human IL-23-producing type 1 macrophages promote but IL-10-producing type 2 macrophages subvert immunity to (myco)bacteria

Frank A. W. Verreck^{*†}, Tjitske de Boer^{*}, Dennis M. L. Langenberg^{*}, Marieke A. Hoeve^{*}, Matthijs Kramer[‡], Elena Vaisberg[‡], Robert Kastelein[‡], Arend Kolk[§], René de Waal-Malefyt[‡], and Tom H. M. Ottenhoff^{*1}

^{*}Department of Immunohematology and Blood Transfusion, Leiden University Medical Centre, Albinusdreef 2, NL-2333-ZA, Leiden, The Netherlands;

[‡]DNAX Research Institute, 901 California Avenue, Palo Alto, CA 94304-1104; and [§]Royal Tropical Institute, Meibergdreef 39, NL-1105-AZ, Amsterdam, The Netherlands

Communicated by Johannes van Rood, Eurodonor Foundation, Leiden, The Netherlands, February 11, 2004 (received for review July 17, 2003)

Macrophages (M ϕ) play a central role as effector cells in immunity to intracellular pathogens such as *Mycobacterium*. Paradoxically, they also provide a habitat for intracellular bacterial survival. This paradoxical role of M ϕ remains poorly understood. Here we report that this dual role may emanate from the functional plasticity of M ϕ : Whereas M ϕ -1 polarized in the presence of granulocyte-M ϕ colony-stimulating factor promoted type 1 immunity, M ϕ -2 polarized with M ϕ colony-stimulating factor subverted type 1 immunity and thus may promote immune escape and chronic infection. Importantly, M ϕ -1 secreted high levels of IL-23 (p40/p19) but no IL-12 (p40/p35) after (myco)bacterial activation. In contrast, activated M ϕ -2 produced neither IL-23 nor IL-12 but predominantly secreted IL-10. M ϕ -1 required IFN- γ as a secondary signal to induce IL-12p35 gene transcription and IL-12 secretion. Activated dendritic cells produced both IL-12 and IL-23, but unlike M ϕ -1 they slightly reduced their IL-23 secretion after addition of IFN- γ . Binding, uptake, and outgrowth of a mycobacterial reporter strain was supported by both M ϕ subsets, but more efficiently by M ϕ -2 than M ϕ -1. Whereas M ϕ -1 efficiently stimulated type 1 helper cells, M ϕ -2 only poorly supported type 1 helper function. Accordingly, activated M ϕ -2 but not M ϕ -1 down-modulated their antigen-presenting and costimulatory molecules (HLA-DR, CD86, and CD40). These findings indicate that (i) M ϕ -1 and M ϕ -2 play opposing roles in cellular immunity and (ii) IL-23 rather than IL-12 is the primary type 1 cytokine produced by activated proinflammatory M ϕ -1. M ϕ heterogeneity thus may be an important determinant of immunity and disease outcome in intracellular bacterial infection.

Mycobacteria can infect human macrophages (M ϕ) and cause serious chronic infectious diseases such as tuberculosis and leprosy. M ϕ play a crucial role in human host defense by secreting cytokines and chemokines, presenting antigen to T lymphocytes and clearing infectious agents. Type 1 cell-mediated immunity is required for granuloma formation and effective host defense against intracellular pathogens (1), but mycobacteria are able to escape immunity and persist in a nonreplicating state inside M ϕ for many years (2). The molecular and cellular mechanisms that underlie the development of effective immunity versus latent infection (or immune escape) and the induction of immunopathology after *Mycobacterium tuberculosis* infection, however, remain poorly understood.

Mononuclear phagocytes including M ϕ are activated through ligation of pattern-recognition receptors such as Toll-like receptors (TLRs) by microbial ligands (3, 4), which is generally considered to potentiate the production of the type 1 cytokine IL-12. IL-12 is a heterodimer of p40 and p35 that drives polarization of naive T cells toward type 1 helper (Th1) cells and induces the release of IFN- γ from T and natural killer cells (5). IFN- γ , in turn, activates M ϕ and enhances cytokine secretion, antigen presentation and, supposedly, the bactericidal activity of M ϕ (6). The IL-12/IFN- γ axis is critical indeed for the establishment of effective host defense against intracellular pathogens: We and others have reported that human

genetic deficiencies in this type 1 cytokine signaling cascade (affecting IL-12p40, IL-12R, IFN- γ R, or STAT-1) lead to increased susceptibility to otherwise weakly pathogenic mycobacteria and salmonellae (reviewed in ref. 1).

Dendritic cells (DCs) are highly potent phagocytes that prime naive T cells and control the development of Th1 cells (7, 8). The role of M ϕ in Th1-mediated immunity, however, is less clear. It has been reported that M ϕ fail to release IL-12 (p40/p35) heterodimer after mycobacterial stimulation (9, 10). Recently, IL-23 was identified as an IL-12-like heterodimer, consisting of IL-12p40 and a novel p19 chain. IL-23, similar to IL-12, induces IFN- γ secretion from T cells (11) and may be involved in type 1 immune defense against mycobacteria (1, 12).

Beside the classical route of M ϕ activation, it has become evident in recent years that alternative activation modes of M ϕ can be distinguished (for recent reviews see refs. 13 and 14). Whereas classical activation of M ϕ by microbial compounds yields a phenotype that is hallmarked by the production of proinflammatory cytokines, alternative activation can lead to an antiinflammatory phenotype, hallmarked by IL-10 as the signature cytokine. Non-classical antiinflammatory M ϕ may also evolve by natural neuroendocrine control mechanisms and play a role in homeostatic processes such as dampening inflammation, scavenging debris, angiogenesis, and wound healing (but also tumor outgrowth) (14–16). The characterization of functional human M ϕ profiles thus far is incomplete, and the role of M ϕ in immunity to mycobacteria remains elusive.

Here we show that highly pure subsets of M ϕ with polarized pro- and antiinflammatory phenotypes can be obtained by differentiating human blood monocytes in the presence of the lineage-determining cytokines granulocyte-M ϕ colony-stimulating factor (GM-CSF) and M ϕ CSF (M-CSF), respectively. Pro- and antiinflammatory M ϕ , designated M ϕ -1 and M ϕ -2, respectively, were characterized for their potential to produce IL-12 and IL-23 to support intracellular mycobacterial growth and present antigens to Th1 lymphocytes. Both M ϕ subsets supported the outgrowth of a mycobacterial reporter strain in the absence of any other immune components, but our findings also indicate that M ϕ -1 promote whereas M ϕ -2 subvert type 1 immunity in the face of (myco)bacterial infection. These results have important implications for our

Abbreviations: M ϕ , macrophage(s); TLR, Toll-like receptor; Th1, type 1 helper; DC, dendritic cell; M-CSF, M ϕ colony-stimulating factor; GM-CSF, granulocyte-M ϕ CSF; mo.DC, monocyte-derived DC; LPS, lipopolysaccharide.

[†]To whom correspondence should be sent at the present address: Biomedical Primate Research Centre, Department of Parasitology, Lange Kleiweg 157; NL-2288-GJ, Rijswijk, The Netherlands. E-mail: verreck@bprc.nl.

¹To whom correspondence should be addressed at: Department of Immunohematology and Blood Transfusion, Leiden University Medical Centre, Albinusdreef 2, NL-2333-ZA, Leiden, The Netherlands. E-mail: t.h.m.ottenhoff@lumc.nl.

© 2004 by The National Academy of Sciences of the USA

understanding of M ϕ biology in cell-mediated immunity to intracellular infections.

Materials and Methods

Cells and Microbial Reagents. Monocytes were isolated to high purity by magnetic cell sorting using anti-CD14-coated beads (per manufacturer recommendations, Miltenyi Biotec, Auburn, CA) and subsequently cultured for 6 days in medium (RPMI medium 1640, GIBCO/Invitrogen) with 10% FCS (HyClone) and either 50 units/ml recombinant human GM-CSF (Novartis Pharma, Arnhem, The Netherlands) to generate M ϕ -1 or 50 ng/ml recombinant human M-CSF (R & D Systems) to generate M ϕ -2. As a control, DCs [monocyte-derived DCs (mo.DCs)] were generated with 1,000 units/ml GM-CSF and 500 units/ml IL-4 as described (17).

Lipopolysaccharide (LPS) from *Escherichia coli* (serotype 055:B5, Sigma-Aldrich) was used to stimulate M ϕ and DCs at 10 ng/ml (unless indicated otherwise). Mycobacterial lysate was obtained by ultrasonication of heat-inactivated *M. tuberculosis* H37Rv, lyophilized, and resuspended in PBS as described (18). The lysate was quantified on the basis of bacterial dry weight; cells were stimulated with 10 μ g/ml unless indicated otherwise.

Cell Cytometry. To analyze cell surface marker expression, aliquots of 10^5 M ϕ were stained for 30 min at 4°C by using appropriate isotype controls and phycoerythrin-conjugated anti-CD14, anti-CD1a, anti-CD83, anti-HLA-DR, anti-CD80, anti-CD86, and anti-CD40 (BD Biosciences/Pharmingen). Samples were analyzed on a FACSCalibur using CELLQUEST software (BD Biosciences).

Cytokine and Chemokine Measurements. IFN- γ secretion was quantified by ELISA (U-CyTech, Utrecht, The Netherlands) with a sensitivity of 20 pg/ml. Specific ELISAs for IL-12p40 and IL-10 were purchased from BioSource International (Camarillo, CA; sensitivity: 20 pg/ml), and IL-12p40/p35 heterodimer was measured by using the cytometric bead assay (BD Biosciences/Pharmingen; sensitivity: 40 pg/ml). IL-23 was measured by ELISA using anti-IL-12p40 monoclonal antibody BP40 (Diacclone) for coating and rat-anti-hp19 monoclonal antibody 12F12 for detection (with a sensitivity of 60 pg/ml).

Polarized M ϕ were harvested by using trypsin-EDTA in Hanks' balanced salt solution without Ca/Mg (GIBCO/Invitrogen). DCs and M ϕ were washed, counted, and seeded in triplicate at 10^5 cells per 200 μ l in 96-well flat-bottom culture plates (Corning Life Sciences) in the presence or absence of stimulating agents as indicated. Supernatants were collected after 24 h (unless indicated otherwise).

mRNA levels of IL-12p40, IL-12p35, and IL-23p19 were measured by real-time quantitative PCR 8 h after activation. RNA was extracted by using an RNeasy kit (Qiagen, Valencia, CA) according to manufacturer protocol and reverse-transcribed with oligo(dT)14–18 (Life Technologies) and random hexamer primers (Promega) by using standard protocols. cDNA was analyzed by PCR with a Perkin-Elmer ABI Prism 7700 sequence-detection system. Gene expression was quantified by correcting for the relative expression to 18S rRNA levels (19).

Mycobacterial Binding, Uptake, and Survival Assays. Mycobacterial binding and uptake were measured by incubating M ϕ at 0 and 37°C with chemically killed (streptomycin/amikacin), GFP-tagged *Mycobacterium bovis* bacillus Calmette–Guérin for 20 min. M ϕ were washed and analyzed by flow cytometry. Intracellular mycobacterial survival was measured by using luciferase-transfected *M. bovis* bacillus Calmette–Guérin as a reporter strain (20). Transfected *M. bovis* bacillus Calmette–Guérin was a generous gift from D. Young (Imperial College, London, U.K.). Briefly, polarized M ϕ were harvested, counted, and seeded at 10^5 cells per 100 μ l per well in 96-well plates in culture medium. After overnight incubation of the M ϕ , 100 μ l of log-phase *M. bovis* bacillus Calmette–Guérin were

added in tissue-culture medium at the concentration indicated. After 2–4 h of infection, M ϕ were washed twice with warm medium and incubated in a final volume of 200 μ l. After 6 days, cells were washed, and luciferase activity was measured after adding Triton X-100 for 10 min [25 μ l of a 10% (vol/vol) solution] by the conversion of *n*-decyl aldehyde [25 μ l of a 1% (wt/vol) solution in ethanol]. Luminescence was measured in a Victor² multilabel plate reader (Perkin-Elmer) equipped with an automatic injector. Host cell viability was determined with a fluorescent calcein viability stain (Molecular Probes). To this end, M ϕ were washed and incubated for 30 min at room temperature in the dark with 50 μ l of 1 mM calcein in anhydrous DMSO (Sigma-Aldrich). Subsequently, the cells were washed twice with PBS, resuspended in 100 μ l of 1% Triton X-100, and measured in the multilabel plate reader.

T Cell Activation Assays. Antigen presentation by M ϕ and DCs was determined by measuring proliferation and IFN- γ secretion of the HLA-DR2-restricted CD4⁺ T cell clone R2F10 that is specific for the 60-kDa heat-shock protein (amino acids 418–427) of *Mycobacterium leprae* (21) or the HLA-DR1-restricted CD4⁺ T cell clone HA1.7 that is specific for the hemagglutinin (amino acids 306–318) of influenza virus (22). Briefly, antigen-presenting cells were harvested and seeded in triplicates at 2.5×10^3 cells per well in 96-well flat-bottom culture plates in 100 μ l in the presence or absence of LPS or mycobacterial sonicate for 24 h. Subsequently, 10^4 R2F10 cells and recombinant protein or synthetic peptide were added to a final concentration of 10 μ g/ml and 100 ng/ml, respectively, in a final volume of 200 μ l and incubated for another 72 h. Aliquots of 50 μ l of supernatant were harvested and pooled per triplicate to measure IFN- γ secretion (see above), and 0.5 μ Ci of [³H]thymidine (1 Ci = 37 GBq) was added for another 18 h to measure T cell proliferation.

Results

Polarization of Monocytes into Type 1 and Type 2 M ϕ Subsets. Highly pure proinflammatory (IL-12p40⁺) M ϕ -1 were obtained after culturing CD14⁺ human blood monocytes for 6 days in the presence of recombinant human GM-CSF, and antiinflammatory (IL-10⁺) M ϕ -2 were obtained by using M-CSF. Both procedures yielded CD14⁺CD1a⁻ cells (Fig. 1A) that were predominantly adherent and had an apparent M ϕ morphology as judged by microscopy (data not shown). For control purposes, from the same donors CD14⁻CD1a⁺CD83⁺ DCs (mo.DCs) were generated (Fig. 1A). Fig. 1B illustrates that M ϕ -1, similar to mo.DCs, secreted high levels of IL-12p40 after activation with either the TLR4-agonist LPS or a sonicate of heat-killed *M. tuberculosis* that stimulates via TLR2 (23). Although M ϕ -1 and M ϕ -2 showed similar levels of (IFN- γ -induced) TLR2 and TLR4 expression (data not shown), activated M ϕ -2 did not secrete IL-12p40 (Fig. 1B) but produced high levels of IL-10 in comparison with M ϕ -1 or mo.DCs (Fig. 1C). These signature cytokine profiles were found consistently in >10 independent experiments using cells from different donors. Moreover, M ϕ -1 but not M ϕ -2 produced (high levels of) IL-18, tumor necrosis factor- α , IL-6, and IL-1 β (F.A.W.V., unpublished observations).

Regulation of IL-12 and IL-23 Production by M ϕ and DCs. Because IL-12p40 can pair with either p35 or p19 to form IL-12 (p40/p35) and IL-23 (p40/p19), respectively, we studied the capacity of M ϕ and DCs to produce these cytokines. Fig. 2 shows that M ϕ -1 and DCs required exogenous IFN- γ to secrete high levels of IL-12 in response to mycobacteria or LPS. Accordingly, IFN- γ was required to induce strong IL-12p35 gene transcription (Fig. 3B). In the absence of IFN- γ , only LPS-activated mo.DCs but not M ϕ -1 produced low levels of IL-12, concordant with the low level of LPS-induced p35 mRNA in these cells (Fig. 3A).

In contrast to IL-12, however, IL-23 was secreted by M ϕ -1 as well as mo.DCs in response to both LPS and mycobacteria without requiring exogenous IFN- γ (Fig. 2). These findings were supported




Fig. 1. Differentiation of human blood monocytes into functionally distinct M ϕ subsets, M ϕ -1 and M ϕ -2. (A) In contrast to mo.DCs, M ϕ -1 and M ϕ -2 highly expressed CD14 but showed no or only weak expression of CD1a or CD83 (the latter after activation by LPS) as determined by flow cytometry. Cytokine secretion was measured up to 72 h after stimulation with *M. tuberculosis* sonicate (myc) (■), LPS (●), or medium control (◆). (B) Whereas activated M ϕ -1, similar to DCs, secreted high levels of IL-12p40, M ϕ -2 failed to secrete IL-12p40. (C) IL-10, in contrast, was most predominant in M ϕ -2. Similar cytokine profiles were obtained with cells from at least 10 independent donors.

by the induction of p40 and p19 mRNA in activated M ϕ -1 and mo.DCs (Fig. 3). The highest levels of IL-23 were reproducibly secreted from *Mycobacterium*-activated M ϕ -1. Together, mycobacterial activation of M ϕ -1 initially results in the induction of IL-23, whereas IL-12 production from these cells requires IFN- γ as an essential second signal. Remarkably, the addition of IFN- γ mildly but reproducibly suppressed the production of IL-23 by DCs, whereas it strongly enhanced IL-23 secretion by M ϕ -1 (Fig. 2). Although IFN- γ enhanced the transcription of p40 mRNA, activation-induced p19 transcription was slightly decreased after the addition of IFN- γ both in DCs and M ϕ -1 (Fig. 3B). This paradoxical finding of IFN- γ -mediated decrease of p19 gene transcription (at $t = 8$ h) and elevation of IL-23 secretion (at $t = 24$ h) in M ϕ -1 suggests an as-yet-unknown posttranscriptional mechanism that affects IL-23 production by M ϕ -1 differently than in DCs.

In agreement with the lack of IL-12p40 production, activated M ϕ -2 completely failed to secrete IL-12 and IL-23 and failed to induce p35, p40, or p19 mRNA (Figs. 2 and 3A).




Fig. 2. Type 1 cytokine secretion by monocyte-derived M ϕ and mo.DCs. The capacity of mo.DCs, M ϕ -1, and M ϕ -2 to secrete IL-12 (p40/p35) and IL-23 (p40/p19) proteins was determined 24 h after microbial stimulation in the absence (black bars) or presence (gray bars) of 500 units/ml IFN- γ . M ϕ -1 secreted IL-23 but failed to produce IL-12 after activation with mycobacterial sonicate (10 μ g/ml) or LPS (10 ng/ml) unless IFN- γ was added. Also, DCs showed IFN- γ -enhanced IL-12 secretion, but unlike M ϕ -1 they yielded decreased rather than elevated IL-23 levels with IFN- γ . M ϕ -2 failed to produce IL-12 or IL-23. Depicted are average protein levels plus standard deviation ($n = 3$; $n = 5$ for IL-12 in DCs and M ϕ -1).

Binding, Uptake, and Outgrowth of Mycobacteria Infecting M ϕ -1 and M ϕ -2. To compare binding and/or uptake of mycobacteria by M ϕ -1 versus M ϕ -2, cells were infected with *M. bovis* bacillus Calmette–Guérin–GFP and analyzed by flow cytometry. Within 20 min at 0°C, \approx 4-fold more M ϕ -2 acquired mycobacteria as compared with M ϕ -1 (Fig. 4A). Similarly, metabolically active M ϕ -2 at 37°C showed at least 2-fold better binding and/or uptake of fluorescent *M. bovis* bacillus Calmette–Guérin (Fig. 4A).

To study the capacity of mycobacteria to survive within M ϕ -1 versus M ϕ -2, we used a luciferase-transfected *M. bovis* bacillus Calmette–Guérin reporter strain (bacillus Calmette–Guérin–lux). As reported previously (20, 24), the luciferase activity of bacillus Calmette–Guérin–lux correlated well with bacterial viability as determined by classical colony-forming unit counting (data not shown). In accordance with the higher capacity to bind and/or phagocytose mycobacteria, we found reproducibly enhanced outgrowth of mycobacteria at day 6 after infection in M ϕ -2 over M ϕ -1 over a range of 20–2.5 infecting mycobacteria per host cell (Fig. 4B). The condition of the infected M ϕ -1 and M ϕ -2 as judged by light microscopy revealed no significant differences during the incubation period. This was corroborated further by calcein viability staining of the host cells [M ϕ -1: 31,977 (\pm 3,449) cps; M ϕ -2: 28,936 (\pm 3,656) cps ($n = 3$)]. Notably, the predominant cytokines in the supernatants of M ϕ -1 and M ϕ -2 at the end of the infection period remained IL-12p40 and IL-10, respectively, suggesting that M ϕ -1 and M ϕ -2 represent stable subsets in the M ϕ spectrum.

M ϕ -1 but Not M ϕ -2 Support Th1 Function After Mycobacterial Stimulation. To address the antigen-presenting capacity of M ϕ -1 and M ϕ -2, we used an HLA-DR2-restricted Th1 reporter clone (R2F10) that recognizes a *M. leprae*-specific HSP60 epitope that is lacking from *M. tuberculosis*. Fig. 5A shows that M ϕ -1 and mo.DCs both were effective in inducing proliferation of R2F10 when presenting *M. leprae* HSP60 protein or peptide antigen. Proliferation of R2F10 Th1 cells induced by M ϕ -1 was not significantly affected by preactivation of M ϕ -1 with *M. tuberculosis* sonicate before adding specific antigen (Fig. 5A). In contrast to M ϕ -1, however, M ϕ -2 stimulated R2F10 relatively poorly, and both pro-




Fig. 3. Type 1 cytokine mRNA levels in monocyte-derived $M\phi$ and mo.DCs. (A) Quantitative reverse transcription–PCR analyses 8 h after LPS stimulation of $M\phi$ -1 revealed that expression of IL-12-specific p35, but not p40 or p19 mRNA, required costimulation with IFN- γ (500 units/ml). DCs showed low but significant transcription of p35 in response to LPS and profound levels of p40 and p19 mRNA. $M\phi$ -2 failed to induce any of these cytokine transcripts. (B) Whereas stimulation with 10 μ g/ml mycobacterial sonicate strongly induced the production of p40 and p19 mRNA, high-level transcription of p35 required costimulation with IFN- γ . (Note that the y axes are logarithmic in B!) Stimulation indexes were calculated from the average level of mRNA from two (DCs) to four ($M\phi$ -1) experiments. Activation-induced p19 transcription was inhibited by IFN- γ in both mo.DCs and $M\phi$ -1.

tein- and peptide-mediated T cell activation were reduced further when $M\phi$ -2 had been preactivated by mycobacterial stimulation (Fig. 5A). Similar poor antigen presentation by $M\phi$ -2 was observed when LPS was used to preactivate the cells (Fig. 5B), indicating that the subversion of Th1 cell activation by $M\phi$ -2 was not unique to *M. tuberculosis* activation. Similar results were obtained with the HLA-DR1-restricted, influenza hemagglutinin-specific T cell clone HA1.7, as illustrated by the dose-dependent reduction in the proliferation of these cells induced by $M\phi$ -2 that had been preactivated with increasing doses of *M. tuberculosis* sonicate (Fig. 5C). Fig. 5D shows that (preactivated) $M\phi$ -1, albeit less efficiently than mo.DCs, supported antigen-specific IFN- γ secretion by R2F10 cells. In contrast, IFN- γ secretion was poorly supported by $M\phi$ -2, and preactivation further reduced this ability (Fig. 5D). In accordance with their poor antigen-presenting capacity, activated $M\phi$ -2 but not $M\phi$ -1 or mo.DCs down-regulated the expression of HLA-DR and CD86 (but not CD80; Fig. 6) as well as CD40 (data not shown).

Thus, although $M\phi$ -1 (and mo.DCs) promote activation of Th1 cells, $M\phi$ -2 fail to promote Th1-mediated immunity efficiently after activation by microbial components.




Fig. 4. Binding, uptake, and outgrowth of mycobacteria by $M\phi$ -1 and $M\phi$ -2. (A) Binding (at 0°C) and/or uptake (at 37°C) of an *M. bovis* bacillus Calmette–Guérin–GFP reporter strain was determined by flow cytometry. After 20 min, more $M\phi$ -2 than $M\phi$ -1 had acquired fluorescent mycobacteria (representative of three independent experiments). (B) Accordingly, over a wide range of multiplicities of infection (MOI) from 20 to 2.5 mycobacteria per host cell, $M\phi$ -2 displayed enhanced outgrowth of *M. bovis* bacillus Calmette–Guérin–lux at day 6 compared with $M\phi$ -1. Depicted are averaged luminescence signals as cps [± standard deviation ($n = 4$)]. Similar results were found with cells from different donors in four independent experiments.

Discussion

$M\phi$ are the major population of tissue-resident mononuclear phagocytes and the predominant targets for infection by intracellular pathogens including mycobacteria. $M\phi$ play a dual role in antimycobacterial host defense that currently is poorly understood: They contribute to cell-mediated immunity and bacterial elimination but also provide an essential niche for intracellular bacterial survival and escape from host defense mechanisms. Here we identify two distinct human $M\phi$ subsets, $M\phi$ -1 and $M\phi$ -2, that display largely opposite functions. Although both $M\phi$ populations can be infected and support the outgrowth of mycobacteria (in the absence of any other immune components), $M\phi$ -1 promote type 1 cellular immunity, whereas $M\phi$ -2 are poor antigen-presenting cells for supporting type 1 immunity. *Mycobacterium*-activated $M\phi$ -1 secrete IL-23 (p40/p19) but no IL-12 (p40/p35), whereas activated $M\phi$ -2 fail to produce IL-23 or IL-12 and predominantly secrete IL-10.

It is well established that type 1 cell-mediated immunity is essential for optimal host defense against intracellular pathogens (1, 6, 25), but it is unresolved how (different functional) $M\phi$ (subsets) contribute to type 1 immunity in antimycobacterial host defense. In the present study, we generated proinflammatory $M\phi$ -1 using the lineage-determining cytokine GM-CSF, which is associated with inflammation. These $M\phi$ -1 efficiently supported the antigen-specific function of Th1 cells. Importantly, our findings indicate that these $M\phi$ -1 initially secrete IL-23 (p40/p19) but no IL-12 (p40/p35) after mycobacterial stimulation. Similar to that for monocytes and mo.DCs (26–28), enhancement of IL-12 production after microbial stimulation of $M\phi$ -1 required IFN- γ , which activates transcription of the IL-12p35 gene. This regulation of IL-12 production may reflect an important function of IFN- γ in enhancing type 1 cellular immunity against intracellular pathogens. Although activation of both DCs and $M\phi$ -1 in the presence of IFN- γ reduced the induction of IL-23-specific p19 mRNA, only DCs reduced IL-23 protein levels under these conditions. $M\phi$ -1, in contrast, enhanced IL-23 secretion after (myco)bacterial stimulation in the presence of IFN- γ . Additional studies are required to unravel the (posttranscriptional) mechanism that is responsible for this differential regulation of IL-23 secretion between mo.DCs and $M\phi$ -1. Although we confirm previous studies that were unable to detect IL-12 heterodimer (9, 10), our study now demonstrates that mycobacterial stimulation of human proinflammatory $M\phi$ triggers IL-23 secretion.




Fig. 5. Antigen-presenting capacity of $M\phi$ -1 versus $M\phi$ -2. To determine the antigen-presenting capacity of $M\phi$ -1 and $M\phi$ -2, cells were pretreated (black bars) or not (gray bars) for 24 h with a (myco)bacterial stimulus before incubation with antigen and antigen-specific Th1 cells. Mo.DCs were included for control purposes. (A) In contrast to $M\phi$ -1 and mo.DCs, $M\phi$ -2 relatively poorly supported proliferation of the *M. leprae*-specific Th1 clone R2F10 toward protein or peptide antigen, which was reduced further after activation of $M\phi$ -2 by *M. tuberculosis* sonicate (myc). (B) Similar results were obtained when antigen-presenting cells were pretreated further after activation with LPS. (C) The antigen-presenting capacity of $M\phi$ -2 toward the influenza hemagglutinin-specific T cell clone HA1.7 was reduced also by mycobacterial stimulation in a dose-dependent fashion. (D) Protein or peptide antigen-specific IFN- γ secretion of R2F10 Th1 cells was supported by $M\phi$ -1, albeit less efficiently than by mo.DCs. IFN- γ secretion by R2F10 cells responding to $M\phi$ -2 was substantially lower and reduced further by mycobacterial activation of $M\phi$ -2. IFN- γ production is depicted as secreted protein in the pooled supernatant of triplicate cultures. Proliferation is depicted as the average incorporation of [3 H]thymidine in triplicate cultures (cpm; + standard deviation). Experiments were repeated at least twice using independent donors to generate $M\phi$ and DCs.

The finding that IL-23 rather than IL-12 is the initial type 1 cytokine released by activated proinflammatory $M\phi$ may point toward an important role for IL-23 in type 1 immunity and anti(myco)bacterial host defense. Because IL-23 has been shown to induce IFN- γ in memory T cells (11), $M\phi$ -derived IL-23 may play a significant role in immunological memory and/or the effector phase of T cell-dependent immunity toward mycobacteria. Murine knock-out models in which either IL-12 or both IL-12 and IL-23 signaling were abrogated indeed have suggested a significant role for IL-23 in host resistance to intracellular pathogens including mycobacteria (12, 29). Interestingly, IL-23 rather than IL-12 was reported to be critical for inflammation in mice (30–32). In support of its unique function, IL-23 rather than IL-12 stimulates mouse T cells to produce GM-CSF (33). Thus, by enhancing GM-CSF




Fig. 6. Expression of HLA and costimulatory molecules on $M\phi$ -1 and $M\phi$ -2. $M\phi$ were pretreated (gray bars) or not (black bars) with mycobacterial sonicate for 48 h before flow-cytometric analysis of HLA-DR, CD80, and CD86 expression. $M\phi$ -2 but not $M\phi$ -1 or mo.DCs down-regulated their cell surface levels of HLA-DR and the costimulatory molecule CD86 (but not CD80) after activation. Depicted are the mean fluorescent intensities (mfi; + standard deviation) of cells from three independent donors.

production, IL-23 may drive differentiation of newly recruited monocytes to a proinflammatory $M\phi$ phenotype.

Over the last few years it has become clear that $M\phi$ are highly heterogeneous, and nonclassical antiinflammatory subsets have been identified (13, 14). Our results show that human $M\phi$ polarized by M-CSF stably display such a nonclassical phenotype, secreting IL-10 but no IL-12(p40) in response to (myco)bacteria. The lack of type 1 cytokine secretion, the high production of IL-10, and the profound down-regulation of HLA, CD86, and CD40 by activated $M\phi$ -2 are likely to contribute to their poor Th1-activating capacity (34–37). $M\phi$ -2 expressed similar levels of TLR4 and TLR2 as $M\phi$ -1, suggesting that the functional differentiation between these two subsets emanates from differential signaling or gene-expression profiles rather than from divergent patterns of innate immune recognition. The enhanced binding and uptake of mycobacteria by $M\phi$ -2 compared with $M\phi$ -1 (already at 0°C) suggests differential expression of a cell surface receptor and also fits with the notion that nonclassical $M\phi$ display enhanced endocytosis (15). This may account for the elevated outgrowth of *M. bovis* bacillus Calmette-Guérin in $M\phi$ -2.

Gordon and coworkers (13, 38) have described alternatively activated $M\phi$ after treatment with IL-4 or IL-13, which in contrast to $M\phi$ -2 produce IL-10 without microbial stimulation. Also unlike alternatively activated $M\phi$, activated $M\phi$ -2 failed to release $M\phi$ -derived chemokine (MDC/CCL22) or thymus- and activation-regulated chemokine (TARC/CCL17) (unpublished observations). However, activated $M\phi$ -2 readily produced other chemokines (e.g., MCP-1/CCL2, IP-10/CXCL10, and MIP-1 β /CCL4; unpublished data), suggesting that $M\phi$ -2 can attract and regulate other immune cells such as monocytes and lymphocytes. The term “type 2 $M\phi$ ” has been used to describe IL-12 $^{-}$ IL-10 $^{+}$ $M\phi$ in the mouse, obtained by stimulation through TLR, CD40, or CD44 ligation in the presence of Fc γ R-ligating immune complexes (39). Additional analyses should reveal how these type 2 $M\phi$ relate to the human $M\phi$ -2 in this study.

A murine *Leishmania* infection model has indicated that IL-10 plays a central role in the maintenance of latent infection by intracellular pathogens (40). Moreover, IL-10-deficient mice display increased antimycobacterial immunity with concordant higher levels of tumor necrosis factor- α and lower bacterial burden (41). Therefore, IL-10 produced by activated $M\phi$ -2 may inhibit optimal host defense and promote latent infection and immune escape by mycobacteria. M-CSF used to generate $M\phi$ -2 is a ubiquitous serum protein, which may indicate that under normal homeostatic conditions $M\phi$ are switched to an antiinflammatory mode. M-CSF-treated $M\phi$ have been described to induce T cell hyporesponsiveness in an indoleamine 2,3-dehydrogenase-dependent fashion and have been implicated in the maintenance of peripheral tolerance (42–44). Furthermore, MCP-1, which is strongly secreted by $M\phi$ -2

(see above) and up-regulated by IL-10, stimulates type 2 helper polarization, and overexpression of MCP-1 in mice increased their susceptibility to infection by *M. tuberculosis* (45–47). Altogether, whereas M ϕ -1 promote cell-mediated immunity, M ϕ -2 seem to down-regulate type 1 cell-mediated immunity by various mechanisms and may promote chronic mycobacterial infection.

Intracellular pathogens generally interact with and modulate their host cells, and *M. tuberculosis* has an intrinsic capacity to interfere with the classical IFN- γ - and TLR-mediated activation pathways of M ϕ (48, 49). Our findings imply that not only active interference of pathogens with M ϕ signaling pathways but also the plasticity in the human M ϕ compartment may critically affect host defense against intracellular infections. Beside providing a

rationale for the paradoxical role of M ϕ in combating intracellular pathogens while also providing a niche that is sequestered from immunosurveillance, our study also provides a model for additional investigation of M ϕ plasticity and M ϕ -derived IL-23 during inflammation, protective immunity, and immunopathology in mycobacterial infections and immunological diseases in general.

We thank Drs. F. Koning and D. Roelen for critically reading the manuscript. Our work was supported by the Netherlands Leprosy Foundation, the Netherlands Organization for Scientific Research, the Commission of the European Community, and the Royal Netherlands Academy of Arts and Sciences.

- Ottenhoff, T. H., Verreck, F. A., Lichtenauer-Kaligis, E. G., Hoeve, M. A., Sanal, O. & Van Dissel, J. T. (2002) *Nat. Genet.* **32**, 97–105.
- Kaufmann, S. H. (2001) *Nat. Rev. Immunol.* **1**, 20–30.
- Janeway, C. A., Jr., & Medzhitov, R. (2002) *Annu. Rev. Immunol.* **20**, 197–216.
- Sieling, P. A. & Modlin, R. L. (2002) *Curr. Opin. Microbiol.* **5**, 70–75.
- Trinchieri, G. (1995) *Annu. Rev. Immunol.* **13**, 251–276.
- Raupach, B. & Kaufmann, S. H. (2001) *Curr. Opin. Immunol.* **13**, 417–428.
- Kalinski, P., Hilkens, C. M., Wierenga, E. A. & Kapsenberg, M. L. (1999) *Immunol. Today* **20**, 561–567.
- Reid, S. D., Penna, G. & Adorini, L. (2000) *Curr. Opin. Immunol.* **12**, 114–121.
- Giacomini, E., Iona, E., Ferroni, L., Miettinen, M., Fattorini, L., Orefici, G., Julkunen, I. & Coccia, E. M. (2001) *J. Immunol.* **166**, 7033–7041.
- Hickman, S. P., Chan, J. & Salgame, P. (2002) *J. Immunol.* **168**, 4636–4642.
- Oppmann, B., Lesley, R., Blom, B., Timans, J. C., Xu, Y., Hunte, B., Vega, F., Yu, N., Wang, J., Singh, K., et al. (2000) *Immunity* **13**, 715–725.
- Cooper, A. M., Kipnis, A., Turner, J., Magram, J., Ferrante, J. & Orme, I. M. (2002) *J. Immunol.* **168**, 1322–1327.
- Gordon, S. (2003) *Nat. Rev. Immunol.* **3**, 23–35.
- Mosser, D. M. (2003) *J. Leukocyte Biol.* **73**, 209–212.
- Goerdts, S. & Orfanos, C. E. (1999) *Immunity* **10**, 137–142.
- Mantovani, A., Sozzani, S., Locati, M., Allavena, P. & Sica, A. (2002) *Trends Immunol.* **23**, 549–555.
- Sallusto, F. & Lanzavecchia, A. (1994) *J. Exp. Med.* **179**, 1109–1118.
- Verbon, A., Kuijper, S., Jansen, H. M., Speelman, P. & Kolk, A. H. (1990) *J. Gen. Microbiol.* **136**, 955–964.
- Scheerens, H., Hessel, E., Waal-Malefyt, R., Leach, M. W. & Rennick, D. (2001) *Eur. J. Immunol.* **31**, 1465–1474.
- Snewin, V. A., Gares, M. P., Gaora, P. O., Hasan, Z., Brown, I. N. & Young, D. B. (1999) *Infect. Immun.* **67**, 4586–4593.
- Ottenhoff, T. H., Klatser, P. R., Ivanyi, J., Elferink, D. G., de Wit, M. Y. & de Vries, R. R. (1986) *Nature* **319**, 66–68.
- Lamb, J. R., Eckels, D. D., Lake, P., Woody, J. N. & Green, N. (1982) *Nature* **300**, 66–69.
- Means, T. K., Wang, S., Lien, E., Yoshimura, A., Golenbock, D. T. & Fenton, M. J. (1999) *J. Immunol.* **163**, 3920–3927.
- Boechat, N., Bouchonnet, F., Bonay, M., Grodet, A., Pelicic, V., Gicquel, B. & Hance, A. J. (2001) *J. Immunol.* **166**, 6203–6211.
- Flynn, J. L. & Chan, J. (2001) *Annu. Rev. Immunol.* **19**, 93–129.
- Snijders, A., Hilkens, C. M., van der Pouw Kraan, T. C., Engel, M., Aarden, L. A. & Kapsenberg, M. L. (1996) *J. Immunol.* **156**, 1207–1212.
- Hayes, M. P., Wang, J. & Norcross, M. A. (1995) *Blood* **86**, 646–650.
- Trinchieri, G. (2003) *Nat. Rev. Immunol.* **3**, 133–146.
- Decken, K., Kohler, G., Palmer-Lehmann, K., Wunderlin, A., Mattner, F., Magram, J., Gately, M. K. & Alber, G. (1998) *Infect. Immun.* **66**, 4994–5000.
- Wiekowski, M. T., Leach, M. W., Evans, E. W., Sullivan, L., Chen, S. C., Vassileva, G., Bazan, J. F., Gorman, D. M., Kastelein, R. A., Narula, S., et al. (2001) *J. Immunol.* **166**, 7563–7570.
- Zhang, G. X., Gran, B., Yu, S., Li, J., Siglienti, I., Chen, X., Kamoun, M. & Rostami, A. (2003) *J. Immunol.* **170**, 2153–2160.
- Cua, D. J., Sherlock, J., Chen, Y., Murphy, C. A., Joyce, B., Seymour, B., Lucian, L., To, W., Kwan, S., Churakova, T., et al. (2003) *Nature* **421**, 744–748.
- Aggarwal, S., Ghilardi, N., Xie, M. H., De Sauvage, F. J. & Gurney, A. L. (2003) *J. Biol. Chem.* **278**, 1910–1914.
- de Waal Malefyt, R., Haanen, J., Spits, H., Roncarolo, M. G., te Velde, A., Figdor, C., Johnson, K., Kastelein, R., Yssel, H. & de Vries, J. E. (1991) *J. Exp. Med.* **174**, 915–924.
- de Waal Malefyt, R., Abrams, J., Bennett, B., Figdor, C. G. & de Vries, J. E. (1991) *J. Exp. Med.* **174**, 1209–1220.
- Koppelman, B., Neeffjes, J. J., de Vries, J. E. & de Waal Malefyt, R. (1997) *Immunity* **7**, 861–871.
- Lim, W., Ma, W., Gee, K., Aucoin, S., Nandan, D., Diaz-Mitoma, F., Kozlowski, M. & Kumar, A. (2002) *J. Immunol.* **168**, 1759–1769.
- Stein, M., Keshav, S., Harris, N. & Gordon, S. (1992) *J. Exp. Med.* **176**, 287–292.
- Anderson, C. F. & Mosser, D. M. (2002) *J. Leukocyte Biol.* **72**, 101–106.
- Belkaid, Y., Hoffmann, K. F., Mendez, S., Kamhawi, S., Udey, M. C., Wynn, T. A. & Sacks, D. L. (2001) *J. Exp. Med.* **194**, 1497–1506.
- Murray, P. J. & Young, R. A. (1999) *Infect. Immun.* **67**, 3087–3095.
- Munn, D. H., Shafizadeh, E., Attwood, J. T., Bondarev, I., Pashine, A. & Mellor, A. L. (1999) *J. Exp. Med.* **189**, 1363–1372.
- Fruento, G., Rotondo, R., Tonetti, M., Damonte, G., Benatti, U. & Ferrara, G. B. (2002) *J. Exp. Med.* **196**, 459–468.
- Terness, P., Bauer, T. M., Rose, L., Dufter, C., Watzlik, A., Simon, H. & Opelz, G. (2002) *J. Exp. Med.* **196**, 447–457.
- Gu, L., Tseng, S., Horner, R. M., Tam, C., Loda, M. & Rollins, B. J. (2000) *Nature* **404**, 407–411.
- Gu, L., Rutledge, B., Fiorillo, J., Ernst, C., Grewal, I., Flavell, R., Gladue, R. & Rollins, B. (1997) *J. Leukocyte Biol.* **62**, 577–580.
- Rutledge, B. J., Rayburn, H., Rosenberg, R., North, R. J., Gladue, R. P., Corless, C. L. & Rollins, B. J. (1995) *J. Immunol.* **155**, 4838–4843.
- Ting, L. M., Kim, A. C., Cattamanchi, A. & Ernst, J. D. (1999) *J. Immunol.* **163**, 3898–3906.
- Nau, G. J., Richmond, J. F., Schlesinger, A., Jennings, E. G., Lander, E. S. & Young, R. A. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 1503–1508.