

Corrections

BIOCHEMISTRY. For the article “Scanning the human proteome for calmodulin-binding proteins,” by Xinchun Shen, C. Alexander Valencia, Jack Szostak, Biao Dong, and Rihe Liu, which appeared in issue 17, April 26, 2005, of *Proc. Natl. Acad. Sci. USA* (**102**, 5969–5974; first published April 19, 2005; 10.1073/pnas.0407928102), the author name Jack Szostak should have appeared as Jack W. Szostak. The online version has been corrected. The corrected author line appears below.

Xinchun Shen, C. Alexander Valencia, Jack W. Szostak, Biao Dong, and Rihe Liu

The authors note several additions to the author contribution footnote. The corrected author contribution footnote, which appears online only, is shown below.

Author contributions: X.S., C.A.V., J.W.S., B.D., and R.L. designed research; X.S., C.A.V., B.D., and R.L. performed research; R.L. contributed new reagents/analytic tools; X.S., C.A.V., B.D., and R.L. analyzed data; and X.S., C.A.V., J.W.S., and R.L. wrote the paper.

www.pnas.org/cgi/doi/10.1073/pnas.0504552102

ECOLOGY. For the article “Diverse taxa of cyanobacteria produce β -N-methylamino-L-alanine, a neurotoxic amino acid,” by Paul Alan Cox, Sandra Anne Banack, Susan J. Murch, Ulla Rasmussen, Georgia Tien, Robert Richard Bidigare, James S. Metcalf, Louise F. Morrison, Geoffrey A. Codd, and Birgitta Bergman, which appeared in issue 14, April 5, 2005, of *Proc. Natl. Acad. Sci. USA* (**102**, 5074–5078; first published April 4, 2005; 10.1073/pnas.0501526102), the authors note that on page 5074, the first sentence of the second full paragraph, right column, should read “BMAA was recently discovered in the brain tissues of eight of nine Alzheimer’s patients,” and not “BMAA was recently discovered in the brain tissues of nine Canadian Alzheimer’s patients.” The authors also note on page 5075, first line in the right column, “Lyophilized samples (5–20 μ g dry weight)” should read “Lyophilized samples (5–20 mg dry weight).” In addition, on page 5076, the unit quantification “mol” in both columns (line 8 in left column; line 1 in right column) should read “ μ mol.” These errors do not change the conclusion of the article.

www.pnas.org/cgi/doi/10.1073/pnas.0504625102

IMMUNOLOGY. For the article “Recombinant NY-ESO-1 protein with ISCOMATRIX adjuvant induces broad integrated antibody and CD4+ and CD8+ T cell responses in humans,” by Ian D. Davis, Weisan Chen, Heather Jackson, Phillip Parente, Mark Shackleton, Wendie Hopkins, Qiyuan Chen, Nektaria Dimopoulos, Tina Luke, Roger Murphy, Andrew M. Scott, Eugene Maraskovsky, Grant McArthur, Duncan MacGregor, Sue Sturrock, Tsin Yee Tai, Simon Green, Andrew Cuthbertson, Darryl Maher, Lena Miloradovic, Susan V. Mitchell, Gerd Ritter, Achim A. Jungbluth, Yao-Tseng Chen, Sacha Gnjjatic, Eric W. Hoffman, Lloyd J. Old, and Jonathan S. Cebon, which appeared in issue 29, July 20, 2004, of *Proc. Natl. Acad. Sci. USA* (**101**, 10697–10702; first published July 13, 2004; 10.1073/pnas.0403572101), the authors note that on page 10701, first column, second full paragraph, “With a median followup of 748 days, 16 have relapsed: five of seven placebo pts, nine of 16 who received protein alone and two of 19 who received NY-ESO-1 with IMX” should read: “With a median followup of 748 days, 16 have relapsed: five of seven placebo pts, six of 16 who received protein alone and two of 19 who received NY-ESO-1 with IMX.” The conclusions of the article remain unchanged.

www.pnas.org/cgi/doi/10.1073/pnas.0504548102

PLANT BIOLOGY. For the article “The *Arabidopsis* SUMO E3 ligase SIZ1 controls phosphate deficiency responses,” by Kenji Miura, Ana Rus, Altanbadralt Sharkhuu, Shuji Yokoi, Athikkattuvalasu S. Karthikeyan, Kashchandra G. Raghohama, Dongwon Baek, Yoon Duck Koo, Jing Bo Jin, Ray A. Bressan, Dae-Jin Yun, and Paul M. Hasegawa, which appeared in issue 21, May 24, 2005, of *Proc. Natl. Acad. Sci. USA* (**102**, 7760–7765; first published May 13, 2005; 10.1073/pnas.0500778102), the authors note that in the Acknowledgments, the “Ministry of Science and Technology” was inadvertently identified as the “Ministry of Sports and Technology,” due to a printer’s error.

www.pnas.org/cgi/doi/10.1073/pnas.0504375102

Diverse taxa of cyanobacteria produce β -N-methylamino-L-alanine, a neurotoxic amino acid

Paul Alan Cox^{*†}, Sandra Anne Banack[‡], Susan J. Murch^{*}, Ulla Rasmussen[§], Georgia Tien[¶], Robert Richard Bidigare[¶], James S. Metcalf^{||}, Louise F. Morrison^{||}, Geoffrey A. Codd^{||}, and Birgitta Bergman[§]

^{*}Institute for Ethnomedicine, National Tropical Botanical Garden, Kalaheo, HI 96741; [†]Institute for Ethnomedicine and Department of Biological Science, California State University, Fullerton, CA 92834; [§]Department of Botany, Stockholm University, S-106 91 Stockholm, Sweden; [¶]Center for Marine Microbial Ecology and Diversity, University of Hawaii at Manoa, Honolulu, HI 96822; and ^{||}Division of Environmental and Applied Biology, School of Life Sciences, University of Dundee, Dundee DD1 4HN, United Kingdom

Communicated by William S. Bowers, University of Arizona, Tucson, AZ, February 24, 2005 (received for review December 3, 2004)

Cyanobacteria can generate molecules hazardous to human health, but production of the known cyanotoxins is taxonomically sporadic. For example, members of a few genera produce hepatotoxic microcystins, whereas production of hepatotoxic nodularins appears to be limited to a single genus. Production of known neurotoxins has also been considered phylogenetically unpredictable. We report here that a single neurotoxin, β -N-methylamino-L-alanine, may be produced by all known groups of cyanobacteria, including cyanobacterial symbionts and free-living cyanobacteria. The ubiquity of cyanobacteria in terrestrial, as well as freshwater, brackish, and marine environments, suggests a potential for widespread human exposure.

biomagnification | neurotoxin | symbiosis | amyotrophic lateral sclerosis/parkinsonism–dementia complex

Cyanotoxin production by cyanobacteria appears to be of wide, but irregular, occurrence throughout the five principal classification sections of cyanobacteria based on axenic strains (1). For example, the hepatotoxic cyclic heptapeptide microcystins are produced by members of the genera *Microcystis*, *Anabaena*, *Planktothrix*, *Hapalosiphon*, *Anabaenopsis*, and *Nostoc*. In contrast, production of the cyclic pentapeptide nodularins appears to be limited to *Nodularia* among the free-living cyanobacterial genera, with indications of nodularin production by cyanobacterial symbionts in a sponge (2). Phylogenetic investigations have indicated an early and widespread occurrence of genes involved in nonribosomal peptide synthesis, including microcystin synthesis. Sporadic distribution of microcystin synthesis in extant cyanobacteria appears to have occurred because of repeated losses of genes for microcystin biosynthesis in the later-derived lineages of these organisms (3). This pattern appears to have arisen because of an early development of microcystin production followed by repeated gene loss in cyanobacterial evolution. The consequences of cyanobacterial toxins on human health, water-based industries, recreation, and wildlife are of increasing concern as eutrophication and rising global temperatures trigger increases in the geographical extent, population densities, and duration of cyanobacterial blooms in fresh, brackish, and marine waters. Human poisonings from cyanobacterial blooms can be serious; 150 persons who drank cyanobacteria-contaminated water in Australia were hospitalized, and >50 kidney dialysis patients at a Brazilian clinic who were exposed to microcystins died (4–7).

β -N-methylamino-L-alanine (BMAA), a nonprotein amino acid, was found to be produced (8) by cyanobacterial root symbionts of the genus *Nostoc* (9). Originally discovered in cycad seeds (10), BMAA was suggested as a possible cause of the amyotrophic lateral sclerosis/parkinsonism–dementia complex (ALS/PDC) that has an extremely high rate of incidence among the Chamorro people of Guam compared with incidence rates of ALS elsewhere (11, 12). Although BMAA as a putative cause of ALS/PDC was initially disputed (13), this hypothesis has re-

cently regained attention when it was discovered that BMAA is biomagnified within the Guam ecosystem (8, 14–16) and is found in the brain tissues of Chamorros who died of ALS/PDC, but not in patients who died of causes unrelated to neurodegenerative disease (8, 16, 17).

BMAA accumulates in ascending trophic levels within the Guam ecosystem. Axenic cultures of *Nostoc* (isolated from the coralloid roots of the cycad tree *Cycas micronesica*) produce BMAA at 0.3 μ g/g. BMAA occurs at 37 μ g/g within mildly infected *C. micronesica* coralloid roots, and at 3,556 μ g/g in the flying foxes that forage on the sarcotesta of *C. micronesica* seeds, a roughly 100-fold increase for each trophic level (8, 15, 18). Because flying foxes are a traditional delicacy (19) of the indigenous Chamorro people, this 10,000-fold increase in BMAA concentration from cyanobacteria to flying foxes suggests that the Chamorros may unwittingly ingest high levels of BMAA in their traditional diet. Flying foxes are not the only BMAA source in the Chamorro diet: cycad seed flour, which shows little free BMAA after washing, releases up to 169 μ g/g BMAA on acid hydrolysis (16). Other protein-associated fractions from each organism in the Guam food chain typically exhibit a 100-fold BMAA increase over the amount of free BMAA (16). Once ingested, BMAA can be bound by proteins within the body, resulting in a slow release of free BMAA over years as contaminated proteins are metabolized. Demonstration, through enzymatic cleavage, that BMAA is incorporated within the actual amino acid sequence of the protein would add weight to this hypothesis. Such a mechanism may explain the observed long latency period of ALS/PDC among the Chamorros (16). We suggest that the Chamorro people of Guam may not be the only human population exposed to this cyanobacteria-produced “slow toxin” (20).

BMAA was recently discovered in the brain tissues of nine Canadian Alzheimer’s patients, but it was not detected in the brain tissues of 14 other Canadians who died of causes unrelated to neurodegeneration (8, 16, 17). Because cycads are not part of the Canadian flora, we suggested that cyanobacteria might be the ultimate source of the BMAA in the Canadian Alzheimer’s patients (8). We also found BMAA in other cyanobacterial–plant symbioses (*Azolla filiculoides*, 2 μ g/g; *Gunnera kauaiensis*, 4 μ g/g) (8). These new findings raised additional questions. Is BMAA produced by other taxa of cyanobacteria? Is the biomagnification of cyanobacteria-produced BMAA unique to the Guam ecosystem or can it occur elsewhere?

Because molecular phylogenies unite the cyanobacteria into a single monophyletic group (21), we decided to examine BMAA production in representative free-living and morphological gen-

Freely available online through the PNAS open access option.

Abbreviations: BMAA, β -N-methylamino-L-alanine; ALS, amyotrophic lateral sclerosis; PDC, parkinsonism–dementia complex.

[†]To whom correspondence should be addressed. E-mail: paul@ethnomedicine.org.

© 2005 by The National Academy of Sciences of the USA




Fig. 1. Cyanobacterial strains that produce BMAA representing different morphological sections. (a) The bloom-forming, filamentous, and colony-forming *Trichodesmium thiebautii* (section III). (Scale bar: 100 μm .) (b) The unicellular *Synechococcus* PCC 6301 (section I). (Scale bar: 1 μm .) (c) The filamentous *Symploca* PCC 8002 (section III). (Scale bar: 10 μm .) (d) The filamentous, nonbranching, and heterocystous (H) *Nostoc* PCC 7107 (section IV). (Scale bar: 10 μm .) (e) The filamentous, heterocystous (H) and branching (arrow) *Fischerella* PCC 7521 (section V). (Scale bar: 15 μm .)

era from all five of the principal sections (Fig. 1) of cyanobacteria (1) as well as in cyanobacterial symbionts isolated from lichen and a diversity of plant species. We analyzed cyanobacterial cultures maintained at the University of Dundee in Scotland, Stockholm University in Sweden, and the University of Hawaii in the United States, as well as natural bloom samples, by using a fluorescent derivatization of amino acids coupled with HPLC to quantify free and protein-associated BMAA for each sample (Fig. 2). We then confirmed selected BMAA peaks through liquid chromatography-MS.

Materials and Methods

Samples of cyanobacterial cultures maintained at the Universities of Stockholm, Hawaii, and Dundee were analyzed for BMAA production. Some cultures had been isolated from symbioses with species of the flowering plant *Gunnera* or from lichens, hornworts, and liverworts; others came from marine, brackish, and freshwater

environments collected throughout the world. Lyophilized samples (5–20 μg dry weight) of cyanobacterial cultures were extracted with 0.1 M trichloroacetic acid and centrifuged at $15,800 \times g$ for 3 min to precipitate proteins. Bound BMAA was released from the protein pellets by hydrolysis (6 M HCl at 110°C for 24 h). Extracts underwent ultrafiltration before derivatization with 6-aminoquinolyl-*N*-hydroxysuccinimidyl carbamate. BMAA was quantified by using a validated HPLC separation. Free amino acids were separated by reverse-phase separation on a gradient HPLC system (Waters 717 Automated Injector and 1525 Binary Solvent Delivery System) and Waters Nova-Pak C18 column at 37°C . Individual compounds were eluted from the column with a gradient elution of 140 mM sodium acetate, 5.6 mM triethylamine (pH 5.2), and 60% or 52% acetonitrile (15). The identity of the BMAA peak was confirmed by comparison with an authenticated standard (Sigma) and was reverified by a modified gradient elution. The concentration of BMAA in samples was determined by fluorescence detec-




Fig. 2. HPLC chromatograph of BMAA peak in *Chroococcidiopsis indica* GT-3–26 (section II) (solid line) and BMAA authenticated standard (dashed line) obtained by using fluorescence detection.

Table 1. BMAA in cyanobacteria isolated from symbioses

Cyanobacteria	Host	Symbiont	Free BMAA, $\mu\text{g/g}$	Protein BMAA, $\mu\text{g/g}$
<i>Nostoc</i> Pc	Lichen	<i>Peltigera</i>	ND	1,671
<i>Nostoc</i> PCC 9305	Hornwort	<i>Anthoceros</i>	156	1,400
<i>Nostoc</i> LBG1	Liverwort	<i>Blasia</i>	ND	ND
<i>Nostoc</i> PCC 73102	Cycad	<i>Macrozamia</i>	ND	ND
<i>Nostoc</i> PCC 7422	Cycad	<i>Cycas</i>	ND	962
<i>Nostoc</i> enc	Cycad	<i>Encephalartos</i>	137	ND
<i>Nostoc</i> PCC 9229	Flowering plant	<i>Gunnera monoica</i>	110	ND
<i>Nostoc</i> 8001	Flowering plant	<i>Gunnera monoica</i>	203	664
<i>Nostoc</i> 9401	Flowering plant	<i>Gunnera perpensa</i>	ND	ND
<i>Nostoc</i> 8963	Flowering plant	<i>Gunnera prorepens</i>	ND	58
<i>Nostoc</i> 8964	Flowering plant	<i>Gunnera macrophylla</i>	9	ND

ND, not detected.

tion (Waters 2487 Dual-Fluorescence Detector) with excitation at 250 nm and emission at 395 nm. Detection of the 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate-derivatized BMAA depended on concentration and comparison of equal amounts of BMAA and a norleucine internal standard, resulting in a mean response of 51.2%. The lower limits of detection and lower limits of quantification were determined by a concentration gradient of an authenticated standard. The lower limits of detection was 0.000065 mol per injection for all analyses. The lower limits of quantification was

0.0065 mol per injection for all analyses. For data interpretation, all sample analyses were quantified within the range of the lower limits of quantification or were reported as not detected (Tables 1 and 2).

The presence of BMAA in the samples as well as the identity and purity of the BMAA peak in the HPLC separations was confirmed by liquid chromatography-MS by using the same HPLC system in-line with a Micromass ZQ/EMD1000 MS (Waters) mass spectrometer, single quadrupole MS with an atmospheric pressure ionization source by using the electrospray

Table 2. BMAA in free-living cyanobacteria

Cyanobacterial species/strain	Section*	Habitat	Origin	Free BMAA, $\mu\text{g/g}$	Protein BMAA, $\mu\text{g/g}$
<i>Microcystis</i> PCC 7806	I	Freshwater	The Netherlands	4	6
<i>Microcystis</i> PCC 7820	I	Freshwater	Scotland	6	12
<i>Prochlorococcus marinus</i> CCMP1377	I	Marine	Sargasso Sea	32 [†]	57 [†]
<i>Synechocystis</i> PCC 6308	I	Freshwater	U.S.A.	ND	ND
<i>Synechococcus</i> PCC 6301	I	Freshwater	U.S.A.	25	ND
<i>Chroococciopsis indica</i> GQ2-7	II	Marine coral	Unknown	435	76
<i>Chroococciopsis indica</i> GT-3-26	II	Marine rock	Unknown	1,306	5,415
<i>Myxosarcina burmensis</i> GB-9-4	II	Marine coral	Marshall Islands	79	1,943
<i>Myxosarcina concinna</i> GT-7-6	II	Marine coral	Unknown	1,501	960
<i>Lyngbya majuscula</i>	III	Marine	Zanzibar	32	4
<i>Planktothrix agardhii</i> NIES 595	III	Freshwater	Northern Ireland	318	30
<i>Plectonema</i> PCC 73110	III	Unknown	Unknown	155	150
<i>Phormidium</i>	III	Unknown	Unknown	11	270
<i>Symploca</i> PCC 8002	III	Marine, intertidal	U.K.	3	262
<i>Trichodesmium thiebautii</i>	III	Marine	Caribbean	145	8
<i>Trichodesmium</i> CCMP1985	III	Marine, coastal	North Carolina	13 [†]	17 [†]
<i>Anabaena</i> PCC 7120	IV	Unknown	U.S.A.	32	ND
<i>Anabaena variabilis</i> ATCC 29413	IV	Freshwater	U.S.A.	35	ND
<i>Aphanizomenon flos-aquae</i>	IV	Marine	Baltic Sea	ND	866
<i>Cylindrospermopsis raciborskii</i> CR3	IV	Freshwater	Australia	6,478	14
<i>Nodularia spumigena</i>	IV	Brackish water	Baltic Sea	16 [†]	50 [†]
<i>Nodularia harveyana</i> CCAP 1452/1	IV	Marine	Unknown	20	11
<i>Nostoc</i> 268	IV	Brackish Water	Baltic Sea	34	274
<i>Nostoc</i> PCC 6310	IV	Freshwater	Israel	42	ND
<i>Nostoc</i> PCC 7107	IV	Freshwater	U.S.A.	27	1,772
<i>Nostoc</i> sp. CMMED 01	IV	Marine	Hawaiian Islands	1,243	1,070
<i>Calothrix</i> PCC 7103	IV	Unknown	Unknown	13 [†]	92 [†]
<i>Chlorogloeopsis</i> PCC 6912	V	Soil	India	758	ND
<i>Fischerella</i> PCC 7521	V	Yellowstone, hot spring	U.S.A.	44	175
<i>Scytonema</i> PCC 7110	V	Limestone cave	Bermuda	ND	1,733

ND, not detected.

*Morphological groupings are as defined in ref.1. Section I, unicellular cyanobacteria that reproduce by binary fusion or budding; section II, unicellular cyanobacteria that reproduce by multiple fission or by both multiple fission and binary fission; section III, filamentous, nonheterocystous cyanobacteria that divide in one plane; section IV, filamentous, heterocystous cyanobacteria that divide in only one plane; section V, heterocystous, filamentous cyanobacteria that divide in more than one plane.

[†]Estimate of concentration based on <1 mg dry weight of sample.

[‡]BMAA was not detected in all samples of this isolate.




Fig. 3. Satellite photograph of a *Trichodesmium* bloom by using SeaWiFS imagery for spectral imaging at 443, 490, and 550 nm off the eastern coast of Florida on October 30, 1998.

ionization interface all controlled by the same computer. Compounds were separated on a reverse-phase column (Nova-Pak C18, 4 Fm, 3.9×300 mm, Waters) heated at 30°C with a linear gradient elution of acetonitrile (10–60% over 25 min) in water. Nitrogen gas was purified and supplied to the electrospray ionization interface with a nebulizing pressure of 100 psi. BMAA was identified through selective ion monitoring in both positive and negative ion modes with detection of the derivatized parent molecule and two daughter ions. All BMAA ions were detected with a dwell time of 1 sec and a cone voltage of 35 V.

Results

Analysis of *Nostoc* strains isolated from symbiotic relationships with lichen and host plants of broad taxonomic diversity indicated that 73% (8/11) of these strains produced BMAA (Table 1).

For free-living cyanobacteria, we found that BMAA is produced by members of all five cyanobacterial sections, as well as in 95% (20/21) of the genera tested, and 97% (29/30) of the strains we tested, representing a wide phylogenetic and ecological diversity (Table 2). Because we found BMAA in an isolate of bloom-forming *Nodularia* from the Baltic Sea (Table 2), we have now begun testing samples from freshwater and marine cyanobacterial blooms. Our preliminary results are intriguing. For example, *Trichodesmium* sp. concentrated from two separate frozen 500-ml samples of seawater from a Hawaiian marine bloom, collected August 18, 2004, had $0.0079 \mu\text{g/g}$ and $0.0071 \mu\text{g/g}$ BMAA wet weight, respectively. The supernatant from the pelleted *Trichodesmium* in seawater had detectable BMAA; however, it was below our lower limits of quantification.

Discussion

Because cyanobacteria are arguably among the most widespread, abundant, and ancient organisms on planet Earth, our finding

that all five sections and 95% of all genera of cyanobacteria tested produce the neurotoxin BMAA is of both ecological and evolutionary significance. For example, *Trichodesmium* blooms are known to cover thousands of square kilometers of the Earth's oceans (Fig. 3). They are observable by satellite from earth orbit and reflect light frequencies that allow taxonomic determination of the cyanobacterium (22). Our detection of BMAA in samples of Baltic Sea and oceanic blooms suggest that significant quantities of BMAA may be released into the world's oceans. Previous analysis of marine blooms of *Trichodesmium thiebautii* and *Trichodesmium erythraeum* have revealed high neurotoxicity to mice, but the low molecular weight neurotoxic entity remained a mystery because peptide toxins and anatoxin-a could not be detected in the bloom extracts (23); if this work is repeated, we suggest that an analysis of BMAA be conducted. Similarly, the occurrence of BMAA in *Prochlorococcus* (Table 2), the most abundant oxyphototroph found in tropical and subtropical waters, could indicate a significant input of neurotoxin at the lowest trophic level of such marine ecosystems (24).

Although we found 95% of cyanobacterial genera and 97% of cyanobacterial strains in our samples to produce BMAA, it is possible that given the right conditions, all cyanobacteria are capable of producing BMAA. Our data (Table 2) show significant variance in the range of BMAA content between genera and species, and also within the ratio of free to protein-bound BMAA. This variance suggests that BMAA production and storage is a function of growth conditions and/or life cycle stages. For example, samples of a single *Calothrix* culture taken 3 months apart showed no BMAA in one sample but quantifiable amounts in the second, and a similar situation was found in *Nodularia spumigena*. This finding is unlike the situation for other known cyanotoxins (neuro- and hepatotoxins), which seem to be consistently produced by

individual toxigenic strains, but with a relatively uneven occurrence among genera (2). Additionally, as our analysis shows that free-living (terrestrial, as well as freshwater, brackish, and marine) cyanobacteria produce BMAA, symbiosis is not a prerequisite for cyanobacterial BMAA production; instead, BMAA production appears to be a common trait within the entire cyanobacterial radiation.

The widespread occurrence of BMAA in our survey suggests that production of this unusual nonprotein amino acid may be a synapomorphy, a common conserved evolutionary feature that unites the cyanobacteria as a group. Because cyanobacteria were among the earliest biological entities to develop on Earth, with Precambrian fossil records conservatively stretching to 1.5 billion years B.P. (25), BMAA is likely to have been produced long before the evolution of organisms with neuronal systems. If so, any selective value of BMAA for the cyanobacteria must have been initially unrelated to its neurotoxicity. Because BMAA likely alters the tertiary conformation of proteins, and because incorporation of BMAA into cyanobacterial peptides may occur as the result of nonribosomal peptide synthesis (26), it is possible that the function of certain aspects of cyanobacterial biochemistry may depend on the presence of this nonprotein amino acid.

Our analysis greatly expands the potential distribution of BMAA in nature as well as the total quantity of BMAA released to the environment. Before 2003, BMAA was known to occur only in cycads, limiting potential human exposure to tropical and subtropical terrestrial environments where cycads grow and then only to indigenous peoples who consume cycad products or animals that feed on cycads. However, these new data demonstrate that BMAA is produced by cyanobacteria from geographical regions and diverse environments throughout the world (Table 2). Because cyanobacteria function as primary producers in many food chains, it is likely that human populations far from Guam may be exposed to this environmental neurotoxin. This suggestion is partially corroborated by the discovery of BMAA in brain tissues of Canadian Alzheimer's patients (16, 17).

In Guam, human exposure to high quantities of BMAA results from unique components of the traditional Chamorro diet including cycad tortillas, flying foxes, and possibly other feral

animals (8). Here, we show that cyanobacterial symbionts of other plants also contain BMAA. Symbioses between cyanobacteria and plants are taxonomically uncommon but can be of ecological importance. For example, ungulates that graze on cyanolichen genera such as *Peltigera* (9, 27, 28) may ingest BMAA at certain seasons of the year (Table 1). The ubiquity of cyanobacteria in diverse terrestrial and aquatic environments suggests that ingestion of BMAA may occur through even less esoteric routes, including direct consumption of cyanobacteria or cyanobacterial hosts, bioaccumulation in additional food chains, or exposure to cyanobacteria-contaminated water supplies.

The recent hypothesis that BMAA accumulates in proteins, which collectively function as an endogenous neurotoxic reservoir within the human body, and then is slowly released through time as these proteins are metabolized (16) suggests that possible health consequences of chronic exposure to low doses of BMAA deserve further investigation. It may now be prudent to monitor BMAA concentrations in drinking waters contaminated by cyanobacterial blooms. BMAA concentrations should also be monitored within invertebrates, fish, or grazing animals used for human consumption that either directly consume cyanobacteria or forage on plants or prey that may have accumulated cyanobacteria-produced BMAA. Given the global importance of marine cyanobacterial blooms, such as those generated by iron-laden dust in the Atlantic and Pacific Oceans (29–31), a broader analysis of the production and fate of BMAA in marine ecosystems is also needed.

We thank M. K. Asay and H. Johnson at the Institute for Ethnomedicine for technical support and R. Honegger of the University of Zurich for useful discussions of cyanolichens. We thank the Acacia Foundation (Larkspur, CA), B. and J. Lane, P. and H. Henry, and C. Childs for laboratory equipment, and N. Kuring of the National Aeronautics and Space Administration (Washington, DC) and A. Subramaniam of the LaMont Doherty Observatory (Columbia University, NY) for satellite imagery. This work was funded by grants from the Harold K. L. Castle Foundation (Kailua, HI) (to P.A.C.); the Swedish Research Council (Stockholm) (to B.B.); the European Commission (Brussels) (to G.A.C.); the Swedish International Development Cooperation Agency (Stockholm) (to U.R.); National Institute for Environmental Health Sciences (Research Triangle Park, NC) (to R.R.B.); and the National Science Foundation (Arlington, VA) (to R.R.B.).

- Rippka, R., Deruelles, J. B., Waterbury, J. B., Herdman, M. & Stanier, R. Y. (1979) *J. Gen. Microbiol.* **111**, 1–61.
- Sivonen, K. & Jones, G. (1999) in *Toxic Cyanobacteria in Water*, eds. Chorus, I. & Bartram, J. (E & F. N. Spon, London), pp. 41–111.
- Rantala, A., Fewer, D. P., Hisbergues, M., Rouhiainen, L., Vaitomaa, J., Börner, T. & Sivonen, K. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 568–573.
- Kuiper-Goodman, T., Falconer, I. & Fitzgerald, J. (1999) in *Toxic Cyanobacteria in Water*, eds. Chorus, I. & Bartram, J. (E & F. N. Spon, London), pp. 113–153.
- Pouria, S., de Andrade, A., Barbosa, J., Cavalcanti, R. L., Barreto, V. T. S., Ward, C. J., Preiser, W., Poon, G. K., Neild, G. H. & Codd, G. A. (1998) *Lancet* **352**, 21–26.
- Carmichael, W. W. (1994) *Sci. Am.* **270**, 78–86.
- Codd, G. A., Bell, S. G., Kaya, K., Ward, C. J., Beattie, K. A. & Metcalf, J. S. (1999) *Eur. J. Phycol.* **34**, 405–415.
- Cox, P. A., Banack, S. A. & Murch, S. J. (2003) *Proc. Natl. Acad. Sci. USA* **100**, 13380–13383.
- Vessey, J. K., Pawlowski, K. & Bergman, B. (2005) *Plant Soil* **266**, 205–230.
- Vega, A. & Bell, E. A. (1967) *Phytochemistry* **6**, 759–762.
- Spencer, P. S., Nunn, P. B., Hugan, J., Ludolph, A. & Roy, D. N. (1986) *Lancet* **i**, 965.
- Spencer, P. S., Nunn, P. B., Hugon, J., Ludolph, A. C., Ross, S. M., Roy, D. N. & Robertson, R. C. (1987) *Science* **237**, 517–522.
- Duncan, M. W., Steele, J. C., Kopin, I. J. & Markey, S. P. (1990) *Neurology* **40**, 767–772.
- Cox, P. A. & Sacks, O. W. (2002) *Neurology* **58**, 956–959.
- Banack, S. A. & Cox, P. A. (2003) *Neurology* **61**, 387–389.
- Murch, S. J., Cox, P. A. & Banack, S. A. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 12228–12231.
- Murch, S. J., Cox, P. A., Banack, S. A., Steele, J. C. & Sacks, O. W. (2004) *Acta Neurol. Scand.* **110**, 267–269.
- Banack, S. A. & Cox, P. A. (2003) *Bot. J. Linn. Soc.* **143**, 165–168.
- Monson, C. S., Banack, S. A. & Cox, P. A. (2003) *Conserv. Biol.* **17**, 678–686.
- Spencer, P. S., Kisby, G. E. & Ludolph, A. C. (1991) *Neurology* **41**, Suppl. 2, 41–68.
- Turner, S. (1988) *Plant Syst. Evol.* **11**, 13–52.
- Subramaniam, A., Brown, C. W., Hood, R. R., Carpenter, E. J. & Capone, D. G. (2002) *Deep-Sea Res.* **49**, 107–121.
- Hawser, S. P. & Codd, G. A. (1992) in *Marine Pelagic Cyanobacteria: Trichodesmium and Other Diazotrophs*, eds. Carpenter, E. J., Capone, D. G. & Rueter, J. G. (Kluwer, Dordrecht, The Netherlands), pp. 319–329.
- Rocap, G., Alrimer, F. W., Lamerdin, J., Malfatti, S., Chain, P., Ahlgren, N. A., Arellano, A., Coleman, M., Hauser, L., Hess, W. R., et al. (2003) *Nature* **424**, 1042–1047.
- Schopf, J. W. (1996) *Nova Hedwigia* **112**, 12–32.
- Neilan, B. A., Dittmann, E., Rouhiainen, L., Bass, R. A., Schaub, V., Sivonen, K. & Börner, T. (1999) *J. Bacteriol.* **181**, 4089–4097.
- Rai, A. N., Bergman, B. & Rasmussen, U. (2002) *Cyanobacteria in Symbiosis* (Kluwer, Dordrecht, The Netherlands).
- Bergman, B., Matveyev, A. & Rasmussen, U. (1996) *Trends Plant Sci.* **1**, 191–197.
- Capone, D. G., Zehr, J. P., Paerl, H. W., Bergman, B. & Carpenter, E. J. (1997) *Science* **276**, 1221–1229.
- Karl, D., Michaels, A., Bergman, B., Capone, D., Carpenter, E., Letelier, R., Lipschultz, F., Paerl, H., Sigma, D. & Stal, L. (2002) *Biogeochemistry* **57/58**, 47–98.
- Lenes, J. M., Darrow, B. P., Cattrell, C., Heil, C. A., Callahan, M., Vargo, G. A., Byrne, R. H., Prospero, J. M., Bates, D., Farning, K. A., et al. (2001) *Limnol. Oceanogr.* **46**, 1261–1277.