

The Müller-Lyer illusion explained by the statistics of image–source relationships

Catherine Q. Howe and Dale Purves*

Center for Cognitive Neuroscience, Duke University, Box 90999, Durham, NC 27708

Contributed by Dale Purves, December 16, 2004

The Müller-Lyer effect, the apparent difference in the length of a line as the result of its adornment with arrowheads or arrow tails, is the best known and most controversial of the classical geometrical illusions. By sampling a range-image database of natural scenes, we show that the perceptual effects elicited by the Müller-Lyer stimulus and its major variants are correctly predicted by the probability distributions of the possible physical sources underlying the relevant retinal images. These results support the conclusion that the Müller-Lyer illusion is a manifestation of the probabilistic strategy of visual processing that has evolved to contend with the uncertain provenance of retinal stimuli.

geometrical illusions | natural scene statistics | vision

The standard Müller-Lyer stimulus (Fig. 1A) has been the subject of hundreds of studies since its introduction in the late 19th century (1). The perceptual effect is that two identical straight lines appear different in length when they are terminated, respectively, with “arrowheads” that extend inward or “arrow tails” that extend outward with respect to the “shaft.” Although there is considerable variation in the reported magnitude of the effect (presumably due to the different experimental conditions in various studies), the line terminated by the arrowheads always appears shorter than same line terminated by arrow tails (2–8).

Rationalizing this illusion has been made especially difficult by persistence of the effect when the identical lines are terminated with a variety of other adornments (9), a fact that undermines intuitive explanations based on what arrowheads and tails might signify. In Fig. 1B, for instance, the same perceptual discrepancy is generated when identical lines are terminated by outward and inward squares. A further obstacle for any simple explanation of the Müller-Lyer effect is that neither the shaft (Fig. 1C) nor continuous lines (Fig. 1D) is needed to elicit a misperception of the relevant spatial interval. Although the effects produced by these several variants have not been quantitatively compared, there is a general agreement that the shaft or the corresponding interval in the “outward” figure always appears longer than its counterpart in the “inward” figure. As a result, there has been much controversy about the genesis of the Müller-Lyer effect (6, 10–20), which still has no generally accepted explanation (21, 22).

Here we test the hypothesis that the standard Müller-Lyer effect and its variants are a result of the fundamentally probabilistic strategy of visual processing that contends with inverse optics problem. Any geometrical stimulus (or indeed any visual stimulus) can have been generated by many different real-world sources (23–27), presenting a quandary to observers whose survival depends on appropriate visually guided behavior. A plausible solution would be to generate visual percepts predicated on the probability distributions of the physical sources of retinal images. In these terms, the identical shafts or intervals in Müller-Lyer stimuli appear different in length because the probability distributions of the real-world sources of the lines or intervals, given the contexts provided by the arrowheads or arrow tails, are in fact different. To test this idea, we determined the physical sources of the standard Müller-Lyer stimulus and its

variants in a range-image database that specified the distance and direction of every point in these natural scenes.

Methods

The range-image database of natural scenes is described in refs. 26 and 27. In keeping with the general approach used to identify the physical sources of lines and angles in these studies, we sampled the range images for sets of pixels whose positions matched the geometrical configurations of the Müller-Lyer stimuli tested.

Fig. 2A shows examples of the geometrical templates used to identify the physical sources of different components of Müller-Lyer stimuli. As a first step, a template was applied to the images to identify areas of the scenes containing the physical sources of one of the pair of adornments in a Müller-Lyer figure (i.e., an arrowhead, an arrow tail, or the equivalent in the Müller-Lyer variants). As indicated in Fig. 2B, the set of pixels underlying the template was then screened to determine whether the physical points corresponding to each straight line in the template formed a geometrically defined straight line in 3D space. If this criterion was met, the points were accepted as a valid sample of the physical source of what we subsequently refer to as the “conditional adornment.”

After identifying a valid physical source of the conditional adornment, the same region of the scene was examined for the occurrence of the other components of the Müller-Lyer figure. For this purpose a series of templates complementary to the template for the conditional adornment was sequentially overlaid on the image (see Fig. 2A). For a standard Müller-Lyer figure, the complementary templates comprised a shaft of increasing length and an arrow adornment configured as the mirror reflection of the conditional adornment. For the Müller-Lyer variant with squares, the complementary templates comprised a square with a shaft of increasing length attached to either the left or the right edge of the square. In the case of the variants without a shaft, or comprising only dots, the complementary template was simply a mirror reflection of the conditional template. This second step thus identifies the “complementary adornment” and the shaft or interval between the two adornments.

The length of the shaft (or the corresponding interval between the two adornments) was varied incrementally from –128 to 128 pixels (negative values indicating that the complementary template was to the left of the conditional adornment and positive values to the right). Thus as the complementary template shifted from the left to the right of the conditional adornment, the overall configuration of the stimulus formed by these two components was reversed (see Fig. 2). As above, the physical points corresponding to each straight line in the complementary template were also evaluated to see whether they formed a straight line in 3D space. If this further criterion was met, the

Freely available online through the PNAS open access option.

Abbreviation: *L*, length of the shaft or corresponding interval in a Müller-Lyer stimulus.

*To whom correspondence should be addressed. E-mail: purves@neuro.duke.edu.

© 2005 by The National Academy of Sciences of the USA




Fig. 3. Statistical analysis of the fully natural scenes in the range-image database for the standard Müller-Lyer stimulus. (A) Probability distributions of the physical sources of Müller-Lyer figures with various shaft lengths (L , in pixels), given the presence of a conditional adornment with its apex pointing either to the right (black) or to the left (gray). In the diagram above, the conditional adornments are indicated by solid lines and the complementary components by dotted lines. (B) The cumulative probability distributions derived from the probability distributions in A. The dotted parts of the curves were computed by extrapolation. (C) Superimposition of the two functions in B. (D) Examples of two shafts 50 pixels in length, one adorned with arrow tails and the other with arrowheads (Upper). The left adornments are arbitrarily designated the conditional adornments and are indicated by solid lines at position 0 (Lower). Given each of these conditional adornments, the probability distributions shown in A–C indicate that the complementary adornment and shaft (dotted lines) can occur at different positions with varying probabilities. The summed probability of occurrence of all possible complementary adornments to the left of position 50 is greater when the fins of the conditional adornment extend to the left of position 0 (black) than when they extend to the right (gray), and conversely. This statistical fact means that a complementary adornment at position 50, given a conditional adornment extending to the left of position 0, is further to the right in the empirical range of possible positions for complementary adornments than is a complementary adornment at position 50, given a conditional adornment extending to the right of position 0.

Perceptual Implications. To understand the perceptual implications of the differences between the two probability distributions in Fig. 3 A–C, consider, as an example, two identical shafts 50 pixels in length, one adorned with arrow tails and the other with arrowheads (Fig. 3D). Imagine taking one of the adornments on each shaft, the one on the left for instance, as the conditional adornment, defining the position of its apex as 0 (the same argument of course applies if the right adornment is selected). The complementary adornments are thus at position 50. Given the conditional adornment in the arrow-tails configuration in Fig. 3D (black), the summed probability of occurrence of the physical sources of complementary adornments whose positions are to the left of position 50 is greater than the comparable summed probability given the conditional adornment in the arrowheads configuration (gray). Conversely, the summed probability of occurrence of the physical sources of complementary adornments located to the right of position 50, given the conditional adornment in the arrow-tails configuration, is less than the comparable summed probability given the conditional adornment in the arrowheads configuration.

These differences in the cumulative probabilities of the stimulus sources mean that the complementary adornment that actually occurred at position 50 in the arrow-tails configuration lies further to the *right* in the empirical range of possible positions of complementary adornments than does the complementary adornment at position 50 in the arrowheads configuration. If the perceptions of the Müller-Lyer stimulus in Fig. 3D are determined by these probabilities, the complementary adornment in the arrow-tails configuration should appear fur-

ther separated from the conditional adornment than the interval between the complementary adornment and the conditional adornment in the arrowheads configuration. Thus the shaft connecting the two adornments in the arrow-tails configuration should be seen as longer than the same line in the arrowheads configuration. The same reasoning can be generalized to Müller-Lyer stimuli with any shaft length, meaning that a shaft adorned with arrow tails should always look longer than the same shaft adorned with arrowheads. These predictions are, of course, consistent with the percepts elicited by the standard Müller-Lyer stimulus.

Statistics Derived from Different Types of Scenes. The results presented so far were derived from the set of fully natural scenes in the database, which is presumably the most important visual environment in the evolution of human perception. We also carried out the same analyses on the scenes in the database that include human constructions because the more rectilinear structure of man-made environments has sometimes been considered a factor contributing to the Müller-Lyer effect (see *Discussion*). The probability distribution of the physical sources of the standard Müller-Lyer stimulus derived from this type of environments is shown in Fig. 4. There is no obvious difference among the results obtained from these different types of scenes.

Variants of the Müller-Lyer Stimulus. Finally, we examined several variants of the standard Müller-Lyer stimulus that elicit the same perceptual effect, including the Müller-Lyer stimulus oriented vertically, identical shafts adorned with squares rather than

15. Gregory, R. L. (1966) *Eye and Brain: The Psychology of Seeing* (McGraw Hill, New York).
16. Griggs, R. (1974) *Bull. Psychonom. Soc.* **4**, 168–170.
17. Morrison, L. C. (1977) *Br. J. Psychol.* **68**, 23–27.
18. Pressey, A. W. (1967) *Percept. Motor Skills* **25**, 569–572.
19. Skottun, B. C. (2000) *Perception* **29**, 201–209.
20. Prinzmetal, W. & Beck, D. M. (2001) *J. Exp. Psychol. Hum. Percept. Perform.* **27**, 206–217.
21. Robinson, J. O. (1998) *The Psychology of Visual Illusion* (Dover, New York).
22. Rock, I. (1995) *Perception* (Scientific Am. Library, New York).
23. Helmholtz, H. L. F. v. (1924) *Helmholtz's Treatise on Physiological Optics* (Opt. Soc. Am., Rochester, NY).
24. Hershenson, M. (1999) *Visual Space Perception* (MIT Press, Cambridge, MA).
25. Mamassian, P., Landy, M. & Maloney, L. T. (2002) in *Probabilistic Models of the Brain: Perception and Neural Function*, eds. Rao, R. P. N., Olshausen, B. A. & Lewicki, M. S. (MIT Press, Cambridge, MA), pp. 13–36.
26. Howe, C. Q. & Purves, D. (2005) *Proc. Natl. Acad. Sci. USA* **102**, 1228–1233.
27. Howe, C. Q. & Purves, D. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 13184–13188.
28. Purves, D. & Lotto, B. (2003) *Why We See What We Do: An Empirical Theory of Vision* (Sinauer, Sunderland, MA).
29. Dragoi, V. & Lockhead, G. (1999) *Vision Res.* **39**, 1657–1670.
30. Carr, H. A. (1935) *An Introduction to Space Perception* (Longmans, New York).
31. Pressey, A. W. (1970) *Percept. Psychophys.* **8**, 411–412.
32. Evans, C. R. & Marsden, R. P. (1966) *Br. J. Physiologic. Opt.* **23**, 242–248.
33. Howe, C. Q. & Purves, D. (2004) *J. Cognit. Neurosci.* **16**, 90–102.