

Increased substrate oxidation and mitochondrial uncoupling in skeletal muscle of endurance-trained individuals

Douglas E. Befroy^{*†}, Kitt Falk Petersen^{*}, Sylvie Dufour[‡], Graeme F. Mason[†], Douglas L. Rothman[†], and Gerald I. Shulman^{*‡§¶}

Departments of ^{*}Internal Medicine, [†]Diagnostic Radiology, [‡]Cellular and Molecular Physiology, and [§]Howard Hughes Medical Institute, Yale University School of Medicine, New Haven, CT 06536

Contributed by Gerald I. Shulman, September 9, 2008 (sent for review June 26, 2008)

Endurance exercise training is accompanied by physiological changes that improve muscle function and performance. Several studies have demonstrated that markers of mitochondrial capacity are elevated, however, these studies tend to be performed *ex vivo* under conditions that yield maximal enzyme activities or *in vivo* but monitoring the response to exercise. Therefore, it is unclear whether basal mitochondrial metabolism is affected by exercise training. To explore whether resting muscle metabolism was altered in trained individuals *in vivo*, two independent parameters of metabolic function—tricarboxylic acid (TCA) cycle flux (V_{TCA}), and ATP synthesis (V_{ATP})—were assessed noninvasively by using magnetic resonance spectroscopy in a cohort of young endurance trained subjects ($n = 7$) and a group of matched sedentary subjects ($n = 8$). V_{TCA} was 54% higher in the muscle of endurance trained compared with sedentary subjects (91.7 ± 7.6 vs. 59.6 ± 4.9 nmol/g/min, $P < 0.01$); however, V_{ATP} was not different between the trained and sedentary subjects (5.98 ± 0.43 vs. 6.35 ± 0.70 μ mol/g/min, $P = 0.67$). The ratio V_{ATP}/V_{TCA} (an estimate of mitochondrial coupling) was also significantly reduced in trained subjects ($P < 0.04$). These data demonstrate that basal mitochondrial substrate oxidation is increased in the muscle of endurance trained individuals yet energy production is unaltered, leading to an uncoupling of oxidative phosphorylation at rest. Increased mitochondrial uncoupling may represent another mechanism by which exercise training enhances muscle insulin sensitivity via increased fatty acid oxidation in the resting state.

ATP synthesis | exercise | magnetic resonance spectroscopy | mitochondria | TCA cycle

Endurance exercise training is accompanied by a number of physiological adaptations that improve muscle function and exercise performance (1). In particular, trained muscle exhibits remodeling toward a more oxidative phenotype (2, 3) that involves changes from the ultrastructural to the subcellular levels. These modifications include an increase in muscle capillary density (3), expanded intramyocellular storage of glycogen (4, 5), and lipid (6) and enhanced insulin responsiveness (7). However, one of the most profound effects of endurance training is the stimulation of mitochondrial biogenesis with an increase in mitochondrial number that is apparent after relatively few weeks of training (8, 9) and is likely mediated by the chronic activation of AMPK (10, 11).

The metabolic response of the muscle to endurance training has been estimated by measuring markers of oxidative capacity with increases in the activities of enzymes involved in mitochondrial metabolism and fat utilization catalogued by numerous studies (5, 12, 14), suggesting that the maximal capacity of the mitochondria, in general, and fat oxidation, in particular, may be increased. These data have been substantiated by studies of mitochondria isolated from trained muscle that demonstrate that elevated mitochondrial enzyme activities are accompanied by enhanced respiratory capacity (9, 15, 16), fatty-acid oxidation

(13, 17), and ATP production (18). However, one limitation of these studies is that they were performed *in vitro* or *ex vivo* under which conditions enable full activation of enzymes and thus represent measurements of maximal capacity rather than function.

Other studies have used indirect calorimetry to estimate whole-body oxidation rates either at rest or during exercise. The data from studies assessing resting metabolic rate are equivocal with no clear effect of endurance training on energy expenditure (19, 20), whereas endurance training consistently appears to improve fatty-acid oxidation in exercising muscle (21, 22). ³¹P magnetic resonance spectroscopy (MRS) has also been used to monitor the recovery of phosphocreatine (PCr) after muscle contraction with increased rates of recovery observed in trained individuals (23, 24), suggesting elevated oxidative capacity. However, calorimetry techniques are constrained by the extrapolation of whole-body oxygen consumption data to muscle metabolism and studies performed during exercise give no insight into basal mitochondrial function.

Therefore, despite a considerable body of work investigating the response of muscle metabolism to endurance training, it is unclear whether resting metabolism, including mitochondrial function, is altered *in vivo*. This is particularly relevant because a reduction in basal muscle metabolism has been associated with the development of insulin resistance in both the elderly (25) and offspring of type 2 diabetic patients (26, 27). Endurance exercise training may represent an effective treatment not only to improve insulin responsiveness in these individuals (28, 29) but also to elevate resting muscle metabolism.

The aim of this study was to investigate the effect of endurance training on basal muscle metabolism *in vivo* using two independent MRS techniques (25–27). The rate of flux through the tri-carboxylic acid (TCA) cycle, a direct measure of mitochondrial oxidative function, was determined by using ¹³C-MRS to monitor the oxidation of intravenously infused [2-¹³C]acetate. The unidirectional rate of muscle ATP synthesis, an assessment of overall energy production, was estimated during the same session by using ³¹P saturation-transfer MRS. These data provide insight into the effects that endurance training may have on resting mitochondrial function *in vivo*.

Author contributions: D.E.B., K.F.P., S.D., and G.I.S. designed research; D.E.B., K.F.P., S.D., D.L.R., and G.I.S. performed research; D.E.B., K.F.P., S.D., G.F.M., and G.I.S. contributed new reagents/analytic tools; D.E.B., K.F.P., S.D., G.F.M., D.L.R., and G.I.S. analyzed data; D.E.B., K.F.P., S.D., G.F.M., D.L.R., and G.I.S. wrote the paper.

The authors declare no conflict of interest.

Freely available online through the PNAS open access option.

[¶]To whom correspondence should be addressed. Yale University School of Medicine, The Anlyan Center, 5269, P.O. Box 9812, New Haven, CT 06536-8012. E-mail: gerald.shulman@yale.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0808889105/DCSupplemental.

© 2008 by The National Academy of Sciences of the USA

Table 1. Characteristics of endurance trained and sedentary subject groups

	Sedentary	Trained
n	8	7
Age, yr	26.8 ± 3.0	26.1 ± 1.6
Height, m	1.80 ± 0.02	1.81 ± 0.03
Weight, kg	74.8 ± 2.8	78.4 ± 3.9
BMI, kg/m ²	23.0 ± 1.03	24.0 ± 1.3
Activity Index	2.24 ± 0.16	5.37 ± 0.26*

BMI, body-mass index.

**P* < 0.0001.

Results

Subject Characteristics. Endurance trained and sedentary subjects were matched for age, height, weight, and body-mass index (BMI) (Table 1). The endurance trained group undertook significantly more exercise than the sedentary group as assessed by the exercise index of the activity questionnaire (5.37 ± 0.26 vs. 2.24 ± 0.16 , *P* < 0.0001). Consistent with these data, the peak oxygen consumption (VO_2 peak) in the trained group (60.9 ± 4.6 ml/kg/min) was increased well above the normal sedentary range (30–40 ml/kg/min).

Muscle Substrate Oxidation – TCA Cycle Flux. Rates of substrate oxidation via the TCA cycle (V_{TCA}) were 54% higher in the muscle of endurance trained individuals compared with sedentary control subjects (91.7 ± 7.6 vs. 59.6 ± 4.9 nmol/g/min, *P* < 0.005, Fig. 1). There were no differences in the steady-state concentration or enrichment of plasma acetate between the groups, or in the maximal enrichment of the muscle C₂ glutamate pool (Table 2). The proportion of acetate oxidized via the TCA cycle ($V_{\text{AC}}/V_{\text{TCA}}$) was also equivalent between groups, indicating that entry of acetate into the myocyte was not limiting (Table 2). There was a trend for a decreased layer of subcutaneous fat between the MR probe and the gastrocnemius muscles of the calf in the trained subjects (3.0 ± 0.5 mm vs. 4.25 ± 0.43 mm, *P* = 0.09), and the total muscle cross-sectional area (CSA) detectable during the ¹³C-MRS experiment was 18% larger in the endurance trained group (19.6 ± 0.5 vs. 16.7 ± 0.6 cm², *P* < 0.005) because of a significant increase in the MR-detectable CSA of the gastrocnemius (12.9 ± 0.9 vs. 7.7 ± 1.1 cm², *P* < 0.005, Table 3). However, V_{TCA} was not correlated with any parameter of muscle CSA within either subject group or for all subjects combined. Whole-body energy expenditure, respiratory quo-




Fig. 1. Muscle TCA cycle flux [nmol/g(muscle)/min] for sedentary (white bar, *n* = 8) and endurance trained (black bar, *n* = 7) subjects.

Table 2. Muscle and plasma metabolite enrichment data and metabolic fluxes obtained during the TCA cycle flux experiment

	Sedentary	Trained
Plasma [acetate], mM	0.63 ± 0.07	0.65 ± 0.03
Plasma [2- ¹³ C]acetate APE, %	85.4 ± 1.1	86.4 ± 1.0
Muscle C ₂ -glutamate APE, %	1.31 ± 0.32	1.39 ± 0.18
Muscle C ₄ -glutamate APE [†] , %	2.62 ± 0.64	2.80 ± 0.36
V_{TCA} , nmol/g (muscle) per min	59.6 ± 4.9	91.7 ± 7.6*
$V_{\text{AC}}/V_{\text{TCA}}$, %	3.6 ± 0.9	3.4 ± 0.4

APE, atom percent excess.

**P* < 0.05.

[†]Calculated from C₂-glutamate assuming 5% anaplerosis.

tient, and fat and glucose oxidation, estimated from indirect calorimetry performed during the MR session, were not different between the trained and sedentary groups (Table 3). There was no correlation between V_{TCA} and any parameter of whole-body metabolism.

Muscle ATP Synthesis. There was no difference in the resting rate of muscle ATP synthesis (V_{ATP}) between the endurance trained and sedentary groups (Fig. 2), with similar values for both the pseudo first order rate constant for P_i exchange (*k'*) and the intracellular inorganic phosphate (P_i) concentration (Table 4). The ratio of $V_{\text{ATP}}/V_{\text{TCA}}$ was significantly decreased in the endurance trained subjects (0.067 ± 0.007 vs. 0.113 ± 0.017 , *P* < 0.05, Fig. 3). Muscle ³¹P metabolite concentrations and ratios were unchanged in the endurance trained group compared with sedentary subjects, and there was no difference in the concentration of free ADP ([ADP]_{free}) estimated from muscle ³¹P metabolite concentrations and the intracellular pH (Table 4).

Discussion

In this study, we found that resting substrate oxidation, calculated by monitoring the rate of incorporation of [2-¹³C]acetate into [4-¹³C]glutamate using ¹³C-MRS, was significantly increased in the muscle of endurance trained male individuals compared with healthy age, height, weight, and sex-matched sedentary subjects. However, the basal rate of muscle ATP synthesis, measured by ³¹P-saturation-transfer MRS, was not different between the endurance trained and sedentary subjects, suggesting that there is an uncoupling of oxidation from energy production in endurance-trained muscle at rest. These MRS-based techniques for measuring metabolic fluxes are independent of the mass of tissue that contributes to the MR signal; therefore, the observed increase in TCA cycle flux rate and

Table 3. Parameters of whole-body energy expenditure, assessed by indirect calorimetry and the cross-sectional areas of muscles contributing to the MR signal in endurance trained (*n* = 7) and sedentary subjects (*n* = 8)

	Sedentary	Trained
Energy expenditure, kcal/kg/24 hr	23.7 ± 0.8	24.2 ± 1.2
Respiratory quotient [†]	0.78 ± 0.02	0.77 ± 0.02
Fat oxidation [†] , mg/kg per min	4.88 ± 0.79	4.89 ± 0.72
Glucose Oxidation [†] , mg/kg per min	0.46 ± 0.20	0.47 ± 0.12
Total muscle CSA, cm ²	16.7 ± 0.6	19.6 ± 0.5*
CSA soleus, cm ²	9.0 ± 1.0	6.7 ± 1.0
CSA gastrocnemius, cm ²	7.7 ± 1.1	12.9 ± 0.9*

CSA, cross sectional area.

**P* < 0.005.

[†]*n* = 7.




Fig. 2. Muscle ATP synthesis rates [$\mu\text{mol/g}(\text{muscle})/\text{min}$] for sedentary (white bar, $n = 8$) and endurance trained (black bar, $n = 7$) subjects.




Fig. 3. The ratio of muscle ATP synthesis/TCA cycle flux (V_{ATP}/V_{TCA}), an estimate of the efficiency of muscle energy production, for sedentary (white bar, $n = 8$) and endurance trained (black bar, $n = 7$) subjects.

decrease in V_{ATP}/V_{TCA} are unaffected by the larger muscle mass of subjects in the trained group.

Our observation of increased basal TCA cycle flux in the muscle of endurance trained individuals is consistent with prior studies that have reported increases in mitochondrial number and oxidative enzyme content in trained muscle, accompanied by an enhanced capacity for respiration, fatty-acid oxidation, and ATP production under maximal conditions, and faster recovery of PCr after a muscle contraction (5, 8, 9, 12–18, 21–24). However, using *in vivo* MRS techniques to investigate the muscle under basal conditions, we have shown that resting muscle oxidation is augmented as well as maximal capacity.

Similar relationships have also been observed in isolated muscle preparations. Resting oxygen consumption (VO_2) was severalfold higher in the highly oxidative soleus muscle compared with the predominantly glycolytic gracilis muscle group in cat, and this was correlated with the mitochondrial density of the muscles (30). Furthermore, chronic low frequency electrical stimulation of the gracilis muscle, designed to mimic exercise training, caused an increase in mitochondrial density accompanied by enhanced VO_2 . Six weeks of endurance training in humans caused an increase in VO_2 in permeabilized vastus-lateralis fibers (31), whereas four weeks of treadmill running in rats increased maximal oxygen consumption in skinned soleus fibers but not gastrocnemius (15).

These data also provide an explanation for the inconsistent results of studies which used indirect calorimetry to investigate the effects of endurance training on resting metabolic rate (19, 20). Although substrate oxidation was elevated in the muscle of trained subjects in this study, we observed no significant effect on energy expenditure measured simultaneously by indirect calorimetry, demonstrating that measurements of whole-body energy expenditure cannot accurately predict significant changes in muscle TCA cycle activity.

Table 4. Muscle ^{31}P metabolite concentrations and ATP synthesis rate (V_{ATP}) in endurance trained and sedentary subjects

	Sedentary	Trained
V_{ATP} , $\mu\text{mol/g per min}$	6.35 ± 0.70	5.98 ± 0.43
k' , min^{-1}	0.043 ± 0.006	0.040 ± 0.003
$[P_i]$, $\mu\text{mol/g}$	2.67 ± 0.24	2.49 ± 0.15
$[PCr]$, $\mu\text{mol/g}$	32.2 ± 1.5	29.9 ± 1.4
pH	7.08 ± 0.01	7.06 ± 0.01
$[ADP]_{\text{free}}$, nmol/g	10.5 ± 0.2	10.1 ± 0.3

Despite increased basal rates of oxidation via the TCA cycle, resting rates of ATP synthesis were unchanged in the muscle of endurance trained individuals, resulting in a significant decrease in the ratio of V_{ATP}/V_{TCA} . Although this ratio is not directly equivalent to the traditional parameter of mitochondrial coupling, the P:O ratio (the amount of ATP produced per oxygen atom consumed), it does reflect an estimate of the efficiency between energy production and oxidative metabolism, and suggests that this relationship is less efficient at rest in trained muscle. Endurance training alters the relative protein composition of the mitochondria (32) which may affect the efficiency of multiple steps in the oxidative phosphorylation pathway. Loss of mitochondrial efficiency—“uncoupling”—may occur via proton (H^+) leak across the inner mitochondrial membrane (IMM), redox “slip” of the electron transport chain (ETC), and modulated stoichiometry of the ATP synthase (33). Substrate selection and mitochondrial respiratory control may also play a role in dictating the efficiency of energy transduction.

Uncoupling protein 3 (UCP3) is a putative uncoupler of mitochondrial oxidative phosphorylation, mediating H^+ leak across the IMM (34). Acute bouts of exercise have been demonstrated to transiently induce the expression of UCP3 in both rodents and humans (34, 35). However, UCP3 expression is lower in oxidative type I muscle fibers (36, 37) than the more glycolytic type II fibers, and expression and protein content appear to be down-regulated in endurance trained muscle (16, 37), making it an unlikely candidate for the decreased efficiency observed in these studies. However, the adenine nucleotide translocase (ANT) has been shown to be increased in endurance trained muscle to a greater extent than other mitochondrial proteins (16). This protein is fundamental to viable mitochondrial function and its increased expression in trained muscle is likely necessary to cope with the high ATP production during exercise. ATP/ADP exchange via the ANT incurs an energetic cost because nucleotide transport is driven by the mitochondrial membrane potential and the ANT may facilitate fatty-acid induced uncoupling (38) or basal mitochondrial H^+ leak (33) proportional to its content, but not activity (39).

The efficiency of oxidative phosphorylation is also dependent on substrate selection with different proportions of reducing equivalents (NADH and $FADH_2$) generated by the full oxidation of glucose (glycolysis plus TCA cycle) compared with fatty-acids (β -oxidation plus TCA cycle) such that the P:O ratio of fatty acid oxidation is lower and thus less efficient than of glucose. The ^{31}P -saturation-transfer technique measures ATP synthesized from all metabolic sources, and the methodology used in the present study does not permit the relative contribu-

repetition time (T_R) of 1.4 seconds (26). Signals from overlapping lipid resonances were suppressed by T_1 -selective nulling after an adiabatic inversion pulse. Temporal resolution was 10 min, corresponding to 424 averages. Spectra were acquired for 20 min before and during a 120-min infusion of 99% enriched $[2-^{13}\text{C}]$ acetate (350 mmol/liter sodium salt) at a rate of 3.0 mg/kg/min. Plasma samples were obtained at 10-min intervals throughout the study for the measurement of plasma acetate concentration and fractional enrichment by gas chromatography/mass spectrometry (26).

To determine the time course of ^{13}C incorporation into C_4 -glutamate during the $[2-^{13}\text{C}]$ acetate infusion, difference spectra for each time-point were obtained by subtracting averaged reference spectra and the increment in the C_4 -glutamate peak was determined by integration (± 0.5 ppm). Absolute enrichment of the C_2 -glutamate peak at the end of infusion was determined by integration (± 0.5 ppm) relative to the natural abundance enrichment (1.1%) of the reference spectra; maximal enrichment at C_4 -glutamate was calculated from that of C_2 -glutamate (assuming 5% dilution of the C_2 pool because of anaplerosis). V_{TCA} for each individual was determined by computer modeling by using CWave software (see below).

Muscle ATP Synthesis – ^{31}P Saturation-Transfer MRS. After a short break, the subject was repositioned in the magnet with the right calf muscles positioned over an MR probe consisting of a 9-cm diameter ^{31}P surface coil and a concentric 13-cm diameter ^1H surface coil. After imaging and shimming, V_{ATP} was measured in each subject by using the ^{31}P saturation-transfer experiment to determine the rate of phosphate exchange between P_i and ATP. Irradiating the γATP peak causes a decrease in the P_i signal that is equivalent to the pseudo-first-order rate constant of the exchange reaction (k') multiplied by the longitudinal relaxation time (T_1) of P_i (Eq. 1). V_{ATP} is equal to the rate constant multiplied by the concentration of P_i (Eq. 2).

$$k' = (1 - M'/M_0)/T_1 \quad [1]$$

$$M' = \text{P}_i \text{ signal during } \gamma\text{ATP saturation}$$

$$M_0 = \text{P}_i \text{ signal with symmetric saturation}$$

$$V_{\text{ATP}} = k'(P_i) \quad [2]$$

Nonlocalized ^{31}P MR spectra were acquired with frequency-selective saturation of the γATP peak or with saturation at a downfield frequency equidistant from P_i . The T_1 relaxation time of P_i during γATP saturation was estimated by using a modified inversion-recovery sequence with γATP saturation during the T_R and the inversion delay. Concentrations of ^{31}P metabolites were determined from control spectra assuming a constant muscle ATP concentration of 5.5 mmol/kg. Intracellular pH was calculated from the chemical shift (δ) of the P_i peak relative to the PCr peak (53).

$$\text{pH} = 6.75 + \log[(\delta - 3.27)/(5.69 - \delta)]$$

$[\text{ADP}]_{\text{free}}$ was determined from the ^{31}P spectra according to the creatine-kinase equilibrium (54), with the equilibrium constant (K_{eq}) corrected for the effects of pH (55) and assuming that 15% of total creatine is unphosphorylated at rest (56).

$$[\text{ADP}]_{\text{free}} = \{[\text{ATP}][\text{Cr}]\}/\{[\text{PCr}][\text{H}^+]\text{K}_{\text{eq}}\}$$

- Fitts RH (2003) Effects of regular exercise training on skeletal muscle contractile function. *Am J Phys Med Rehabil* 82:320–331.
- Bergh U, et al. (1978) Maximal oxygen uptake and muscle fiber types in trained and untrained humans. *Med Sci Sports* 10:151–154.
- Coggan AR, et al. (1992) Skeletal muscle adaptations to endurance training in 60- to 70-yr-old men and women. *J Appl Physiol* 72:1780–1786.
- Holloszy JO, Booth FW (1976) Biochemical adaptations to endurance exercise in muscle. *Annu Rev Physiol* 38:273–291.
- Gollnick PD, et al. (1973) Effect of training on enzyme activity and fiber composition of human skeletal muscle. *J Appl Physiol* 34:107–111.
- Hoppeler H, et al. (1985) Endurance training in humans: Aerobic capacity and structure of skeletal muscle. *J Appl Physiol* 59:320–327.
- Ebeling P, et al. (1993) Mechanism of enhanced insulin sensitivity in athletes. Increased blood flow, muscle glucose transport protein (GLUT-4) concentration, and glycogen synthase activity. *J Clin Invest* 92:1623–1631.
- Hood DA (2001) Invited Review: Contractile activity-induced mitochondrial biogenesis in skeletal muscle. *J Appl Physiol* 90:1137–1157.
- Holloszy JO, Coyle EF (1984) Adaptations of skeletal muscle to endurance exercise and their metabolic consequences. *J Appl Physiol* 56:831–838.
- Bergeron R, et al. (1999) Effect of AMPK activation on muscle glucose metabolism in conscious rats. *Am J Physiol Endocrinol Metab* 276:E938–E944.

CWave Modeling. V_{TCA} was determined by modeling the incorporation of ^{13}C label from plasma $[2-^{13}\text{C}]$ acetate into the muscle $[4-^{13}\text{C}]$ glutamate pool [supporting information (SI) Fig. S1], using CWave software (26). The CWave model consists of isotopic mass balance equations that describe the metabolic fate of the plasma $[2-^{13}\text{C}]$ acetate (see SI Text). Upon entry into the myocyte, $[2-^{13}\text{C}]$ acetate is converted (V_{AC}) into $[2-^{13}\text{C}]$ acetylCoA which enters the TCA cycle by condensing with oxaloacetate to form $[4-^{13}\text{C}]$ citrate. Entry of unlabeled substrates into the TCA cycle via acetyl CoA or anaplerosis was incorporated into the model as a separate reaction ($V_{\text{PDH+FA}}$). The position of the ^{13}C label is conserved through the initial steps of the TCA cycle, labeling α -ketoglutarate at the C_4 position. Glutamate and α -ketoglutarate are in rapid exchange and equilibration results in the formation of $[4-^{13}\text{C}]$ glutamate. As the TCA cycle progresses, the ^{13}C label becomes scrambled between the C_2 and C_3 positions because of the symmetry of the succinate molecule; a second turn of the TCA cycle yields $[2-^{13}\text{C}]$ and $[3-^{13}\text{C}]$ glutamate. CWave determines V_{TCA} as the rate of total carbon flow from acetyl CoA to α -ketoglutarate by using a nonlinear least-squares algorithm to fit the curve of C_4 -glutamate enrichment. In this model, V_{TCA} is equal to the sum of ($V_{\text{AC}}+V_{\text{PDH+FA}}$) but is independent of the absolute fractional enrichment at C_4 -glutamate ($V_{\text{AC}}/V_{\text{PDH+FA}}$). The intracellular concentration of glutamate was measured by muscle biopsy and was assumed to be 2.41 mmol/liter (25) and unchanged by endurance training (57). The ratio of natural abundance C_2 -glutamate/ C_2 -creatine was the same between the groups and there was no significant difference in the fractional enrichment at C_2 -glutamate at the end of the $[2-^{13}\text{C}]$ acetate infusion, confirming that the intracellular concentration of glutamate was unchanged in endurance trained individuals. The rate of exchange between α -ketoglutarate and glutamate (V_X) was determined in prior studies (25) and was fixed at 150 nmol/g(muscle)/min that is significantly faster than the TCA cycle flux.

Muscle Cross-sectional Area. The relative contributions of the soleus and gastrocnemius muscles to the observed MR signal were estimated from MR images obtained during the study. The MR-detectable CSA of each muscle group was measured in the image slice that corresponded with the center of the MR probe.

Indirect Calorimetry. Whole-body energy expenditure was measured during the MR studies by the ventilated hood technique by using a Deltatrac Metabolic Monitor. Nonprotein respiratory quotients assuming 100% oxidation of fat or carbohydrates were 0.707 and 1.00, respectively.

Statistical Analysis. All data are expressed as mean \pm SE. Statistical analyses were performed by using InStat3 software (Graphpad Software). Statistically significant differences between trained and sedentary subjects were detected by using an unpaired 2-tailed student's t test. A P value of <0.05 was considered statistically significant.

ACKNOWLEDGMENTS. We thank the subjects for their participation in this study, Dr. James Dziura for statistical assistance, and Yanna Kosover, Anthony Romanelli, Mikhail Smolgovsky, and the staff of the Yale Center for Clinical Investigation for technical assistance with the studies. This work was supported by National Institutes of Health Grants R01 NS-037527 (to D.L.R.), K02 AA-13430 (to G.F.M.), R01 AG-23686 (to K.F.P.), M01 RR-00125, R01 DK-49230 (to G.I.S.), and P01 DK-68229 (to G.I.S.), The Keck Foundation, and a Distinguished Clinical Scientist Award from the American Diabetes Association (to G.I.S.).

- Zong H, et al. (2002) AMP kinase is required for mitochondrial biogenesis in skeletal muscle in response to chronic energy deprivation. *Proc Natl Acad Sci USA* 99:15983–15987.
- Holloszy JO (1967) Biochemical adaptations in muscle. Effects of exercise on mitochondrial oxygen uptake and respiratory enzyme activity in skeletal muscle. *J Biol Chem* 242:2278–2282.
- Mole PA, Oscari LB, Holloszy JO (1971) Adaptation of muscle to exercise. Increase in levels of palmityl Coa synthetase, carnitine palmityltransferase, and palmityl Coa dehydrogenase, and in the capacity to oxidize fatty acids. *J Clin Invest* 50:2323–2330.
- Kiens B, et al. (1997) Membrane associated fatty acid binding protein (FABPpm) in human skeletal muscle is increased by endurance training. *Biochem Biophys Res Commun* 231:463–465.
- Burelle Y, Hochachka PW (2002) Endurance training induces muscle-specific changes in mitochondrial function in skinned muscle fibers. *J Appl Physiol* 92:2429–2438.
- Fernstrom M, Tonkonogi M, Sahlin K (2004) Effects of acute and chronic endurance exercise on mitochondrial uncoupling in human skeletal muscle. *J Physiol* 554:755–763.
- Koves TR, et al. (2005) Subsarcolemmal and intermyofibrillar mitochondria play distinct roles in regulating skeletal muscle fatty acid metabolism. *Am J Physiol Cell Physiol* 288:C1074–C1082.
- Wibom R, et al. (1992) Adaptation of mitochondrial ATP production in human skeletal muscle to endurance training and detraining. *J Appl Physiol* 73:2004–2010.

19. Poehlman ET, Melby CL, Badylak SF, Calles J (1989) Aerobic fitness and resting energy expenditure in young adult males. *Metabolism* 38:85–90.
20. Bullough RC, Gillette CA, Harris MA, Melby CL (1995) Interaction of acute changes in exercise energy expenditure and energy intake on resting metabolic rate. *Am J Clin Nutr* 61:473–481.
21. Martin WH, 3rd, et al. (1993) Effect of endurance training on plasma free fatty acid turnover and oxidation during exercise. *Am J Physiol* 265:E708–E714.
22. Coggan AR, et al. (2000) Fat metabolism during high-intensity exercise in endurance-trained and untrained men. *Metabolism* 49:122–128.
23. McCully KK, et al. (1989) Wrist flexor muscles of elite rowers measured with magnetic resonance spectroscopy. *J Appl Physiol* 67:926–932.
24. Takahashi M, et al. (1995) Control of the rate of phosphocreatine resynthesis after exercise in trained and untrained human quadriceps muscles. *Eur J Appl Physiol Occup Physiol* 71:396–404.
25. Petersen KF, et al. (2003) Mitochondrial dysfunction in the elderly: Possible role in insulin resistance. *Science* 300:1140–1142.
26. Befroy DE, et al. (2007) Impaired mitochondrial substrate oxidation in muscle of insulin-resistant offspring of type 2 diabetic patients. *Diabetes* 56:1376–1381.
27. Petersen KF, et al. (2004) Impaired mitochondrial activity in the insulin-resistant offspring of patients with type 2 diabetes. *N Engl J Med* 350:664–671.
28. Perseghin G, et al. (1996) Increased glucose transport-phosphorylation and muscle glycogen synthesis after exercise training in insulin-resistant subjects. *N Engl J Med* 335:1357–1362.
29. Goodpaster BH, Katsiaras A, Kelley DE (2003) Enhanced fat oxidation through physical activity is associated with improvements in insulin sensitivity in obesity. *Diabetes* 52:2191–2197.
30. Hoppeler H, Hudlicka O, Uhlmann E (1987) Relationship between mitochondria and oxygen consumption in isolated cat muscles. *J Physiol* 385:661–675.
31. Walsh B, Tonkonogi M, Sahlin K (2001) Effect of endurance training on oxidative and antioxidative function in human permeabilized muscle fibres. *Pflugers Arch* 442:420–425.
32. Holloszy JO, Oscai LB, Don JI, Mole PA (1970) Mitochondrial citric acid cycle and related enzymes: Adaptive response to exercise. *Biochem Biophys Res Commun* 40:1368–1373.
33. Brand MD (2005) The efficiency and plasticity of mitochondrial energy transduction. *Biochem Soc Trans* 33:897–904.
34. Schrauwen P, Saris WH, Hesselink MK (2001) An alternative function for human uncoupling protein 3: Protection of mitochondria against accumulation of nonesterified fatty acids inside the mitochondrial matrix. *FASEB J* 15:2497–2502.
35. Pilegaard H, Ordway GA, Saltin B, Neufer PD (2000) Transcriptional regulation of gene expression in human skeletal muscle during recovery from exercise. *Am J Physiol Endocrinol Metab* 279:E806–E814.
36. Hesselink MK, et al. (2001) Protein expression of UCP3 differs between human type 1, type 2a, and type 2b fibers. *FASEB J* 15:1071–1073.
37. Russell AP, et al. (2003) UCP3 protein expression is lower in type I, IIa and IIx muscle fiber types of endurance-trained compared to untrained subjects. *Pflugers Arch* 445:563–569.
38. Simonyan RA, Skulachev VP (1998) Thermoregulatory uncoupling in heart muscle mitochondria: Involvement of the ATP/ADP antiporter and uncoupling protein. *FEBS Lett* 436:81–84.
39. Brand MD, et al. (2005) The basal proton conductance of mitochondria depends on adenine nucleotide translocase content. *Biochem J* 392:353–362.
40. Cocco T, Pacelli C, Sgobbo P, Villani G (2008) Control of OXPHOS efficiency by complex I in brain mitochondria. *Neurobiol Aging*, in press.
41. Mogensen M, Sahlin K (2005) Mitochondrial efficiency in rat skeletal muscle: Influence of respiration rate, substrate and muscle type. *Acta Physiol Scand* 185:229–236.
42. Pande SV, Blanchaer MC (1971) Carbohydrate and fat in energy metabolism of red and white muscle. *Am J Physiol* 220:549–553.
43. Krieger DA, Tate CA, McMillin-Wood J, Booth FW (1980) Populations of rat skeletal muscle mitochondria after exercise and immobilization. *J Appl Physiol* 48:23–28.
44. Cogswell AM, Stevens RJ, Hood DA (1993) Properties of skeletal muscle mitochondria isolated from subsarcolemmal and intermyofibrillar regions. *Am J Physiol* 264:C383–C389.
45. Palmer JW, Tandler B, Hoppel CL (1977) Biochemical properties of subsarcolemmal and interfibrillar mitochondria isolated from rat cardiac muscle. *J Biol Chem* 252:8731–8739.
46. Kemp GJ, Taylor DJ, Radda GK (1993) Control of phosphocreatine resynthesis during recovery from exercise in human skeletal muscle. *NMR Biomed* 6:66–72.
47. Gates AC, et al. (2007) Respiratory uncoupling in skeletal muscle delays death and diminishes age-related disease. *Cell Metab* 6:497–505.
48. Choi CS, et al. (2007) Overexpression of uncoupling protein 3 in skeletal muscle protects against fat-induced insulin resistance. *J Clin Invest* 117:1995–2003.
49. Baecke JA, Burema J, Frijters JE (1982) A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr* 36:936–942.
50. Watt MJ, Heigenhauser GJ, Spriet LL (2002) Intramuscular triacylglycerol utilization in human skeletal muscle during exercise: Is there a controversy? *J Appl Physiol* 93:1185–1195.
51. Krssak M, et al. (2000) Intramuscular glycogen and intramyocellular lipid utilization during prolonged exercise and recovery in man: A ¹³C and ¹H nuclear magnetic resonance spectroscopy study. *J Clin Endocrinol Metab* 85:748–754.
52. Shen J, Rycyna RE, Rothman DL (1997) Improvements on an in vivo automatic shimming method [FASTERMAP]. *Magn Reson Med* 38:834–839.
53. Taylor DJ, et al. (1983) Bioenergetics of intact human muscle. A ³¹P nuclear magnetic resonance study. *Mol Biol Med* 1:77–94.
54. Lawson JW, Veech RL (1979) Effects of pH and free Mg²⁺ on the K_{eq} of the creatine kinase reaction and other phosphate hydrolyses and phosphate transfer reactions. *J Biol Chem* 254:6528–6537.
55. Golding EM, Teague WE, Jr., Dobson GP (1995) Adjustment of K⁺ to varying pH and pMg for the creatine kinase, adenylate kinase and ATP hydrolysis equilibria permitting quantitative bioenergetic assessment. *J Exp Biol* 198:1775–1782.
56. Boska M (1994) ATP production rates as a function of force level in the human gastrocnemius/soleus using ³¹P MRS. *Magn Reson Med* 32:1–10.
57. Dawson KD, et al. (2003) Short-term training attenuates muscle TCA cycle expansion during exercise in women. *J Appl Physiol* 95:999–1004.