

Candicidin-producing *Streptomyces* support leaf-cutting ants to protect their fungus garden against the pathogenic fungus *Escovopsis*

Susanne Haeder^a, Rainer Wirth^b, Hubert Herz^{c,d}, and Dieter Spiteller^{a,1}

^aDepartment of Bio-Organic Chemistry, Max Planck Institute for Chemical Ecology, Hans-Knöll-Strasse 8, D-07745 Jena, Germany; ^bDepartment of Plant Ecology and Systematics, Technische Universität Kaiserslautern, Erwin-Schrödinger-Strasse 13, D-67663 Kaiserslautern, Germany; and ^cSmithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancón, Republic of Panamá; and ^dBiology Department, San Francisco State University, 1600 Holloway Avenue, San Francisco, CA 94132

Edited by Jerrold Meinwald, Cornell University, Ithaca, NY, and approved January 28, 2009 (received for review December 1, 2008)

Leaf-cutting ants such as *Acromyrmex octospinosus* live in obligate symbiosis with fungi of the genus *Leucoagaricus*, which they grow with harvested leaf material. The symbiotic fungi, in turn, serve as a major food source for the ants. This mutualistic relation is disturbed by the specialized pathogenic fungus *Escovopsis* sp., which can overcome *Leucoagaricus* sp. and thus destroy the ant colony. Microbial symbionts of leaf-cutting ants have been suggested to protect the fungus garden against *Escovopsis* by producing antifungal compounds [Currie CR, Scott JA, Summerbell RC, Malloch D (1999) Fungus-growing ants use antibiotic-producing bacteria to control garden parasites. *Nature* 398:701–704.]. To date, however, the chemical nature of these compounds has remained elusive. We characterized 19 leaf-cutting ant-associated microorganisms (5 *Pseudonocardia*, 1 *Dermaococcus*, and 13 *Streptomyces*) from 3 *Acromyrmex* species, *A. octospinosus*, *A. echinator*, and *A. volcanus*, using 16S-rDNA analysis. Because the strain *Streptomyces* sp. Ao10 proved highly active against the pathogen *Escovopsis*, we identified the molecular basis of its antifungal activity. Using bioassay-guided fractionation, high-resolution electrospray mass spectrometry (HR-ESI-MS), and UV spectroscopy, and comparing the results with an authentic standard, we were able to identify candicidin macrolides. Candicidin macrolides are highly active against *Escovopsis* but do not significantly affect the growth of the symbiotic fungus. At least one of the microbial isolates from each of the 3 leaf-cutting ant species analyzed produced candicidin macrolides. This suggests that candicidins play an important role in protecting the fungus gardens of leaf-cutting ants against pathogenic fungi.

Acromyrmex | antifungal agent | symbionts | Attini | polyketides

Ants have fascinated people for a long time, in part because many aspects of their social organization appear so similar to those of humans (1). The highly evolved leaf-cutting ants of *Acromyrmex* and *Atta* spp. (Hymenoptera: Formicidae: Attini) have attracted particular attention because these ants cultivate a symbiotic fungus (Agaricales: mostly Lepiotaceae: Leucocoprineae) in specialized chambers of their nests. The ants cut leaves, preprocess them into small pieces, and feed them to the symbiotic fungus, which, in turn, serves as their major food source (2). However, their highly sophisticated behavior aside, leaf-cutting ants are both dominant herbivores of neotropical forests (2) and a serious problem to crop plants in South America (3).

Leaf-cutting ants, like other fungus-growing ants (tribe Attini), are threatened by specialized fungal pathogens; *Escovopsis* (Ascomycota: anamorphic Hypocreales) was shown to have detrimental effects on the health of the fungal gardens and, consequently, on the survival of the ant colony (4, 5). To avoid infection, leaf-cutting ants have evolved an elaborate cleaning behavior, carefully maintaining their fungus garden and placing old or suspicious material into waste chambers (6). In addition to mechanically removing waste, leaf-cutting ants secrete anti-

microbial and antifungal agents from their metapleural glands [e.g., myrmicacin (3-hydroxydecanoic acid)], which help to protect against pathogens (7–10).

In their pioneering work, Currie et al. (11) discovered symbiotic microorganisms on the integument of leaf-cutting ants as additional players in these interactions. Those microorganisms, identified as *Pseudonocardia* spp., were suggested to be part of a 50 million-year-old tripartite coevolutionary process. Microorganisms producing antifungal compounds coevolved to protect the ants' fungus against the pathogens *Escovopsis aspergilloides* and *Escovopsis weberi* (12, 13). However, recent evidence indicates that diverse microbial communities rather than a single symbiont exist in the leaf-cutting ants' gardens. This contradicts the prevailing view, according to which attine ants, their garden fungus, and the pathogens specifically coevolved with *Pseudonocardia*. To give an example, *Streptomyces* and *Burkholderia* are now thought to be involved in this multitrophic interaction (14–16). The studies of Kost et al. (14) and Mueller et al. (16) provide strong evidence that the microbial communities of leaf-cutting ants are flexible, highly diverse, and likely to result when microorganisms are recruited from the environment.

Although our knowledge about the ants' microbial symbionts has evolved dramatically in recent years (11), the chemical basis of their antifungal activities has not yet been elucidated. Using a bioassay-guided approach, we isolated an antifungal compound produced by a *Streptomyces* strain associated with leaf-cutting ants. Structure elucidation revealed candicidin macrolides that are highly active against the fungal pathogen *Escovopsis* but not against the ants' garden fungus *Leucoagaricus gongylophorus*.

Results

Isolation and Characterization of Microorganisms Associated with Leaf-Cutting Ants. Microorganisms from 3 leaf-cutting ant species (*Acromyrmex octospinosus*, *Acromyrmex echinator*, and *Acromyrmex volcanus*) were isolated from individual worker ants or small fungus garden samples. *Streptomyces*-type microorganisms were isolated using selective agar plates (17), and their typical morphology was considered as the selection criterion (18). Microorganisms were found in all 3 ant species. 16S-rDNA analysis of the isolates allowed us to classify the isolates [supporting information (SI) Table S1 and Fig. S1].

Six strains were obtained from *A. octospinosus*: 4 *Pseudono-*

Author contributions: D.S. designed research; S.H., R.W., H.H., and D.S. performed research; S.H. and D.S. analyzed data; and D.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Data deposition: The sequences reported in this paper have been deposited in the BankIt database (accession nos. FJ490527–FJ490550).

¹To whom correspondence should be addressed. E-mail: dspiteller@ice.mpg.de.

This article contains supporting information online at www.pnas.org/cgi/content/full/0812082106/DCSupplemental.




Fig. 2. Antifungal activity assay of candicidin macrolides (90 nmol) against *E. weberi*, *B. bassiana*, *M. anisopliae*, *L. lecanii*, and the symbiotic fungus *L. gongylophorus* in comparison to the solvent control.

of candicidin-producing microorganisms on the leaf-cutting ants' bodies.

Having identified candicidin macrolides as an antifungal principle of *Streptomyces* sp. Ao10, we screened for candicidins and related polyketide polyenes in the other microbial isolates from leaf-cutting ants. Six of 19 microorganisms were able to produce candicidin macrolides. At least 1 associated microorganism in all 3 leaf-cutting ant species (*A. octospinosus*, *A. echinator*, and *A. volcanus*) was found to produce candicidins. All candicidin-producing microorganisms belonged to *Streptomyces* (Table S1).

Candicidin Macrolides as Selective Antifungal Compounds Against *Escovopsis*. After the candicidin macrolides were identified, their antifungal potential was addressed. In the agar diffusion assay against *E. weberi*, 90-nmol candicidin macrolides created a large inhibition zone of \varnothing 1.6 cm after 4 days of incubation (Fig. 2). Nine nanomoles of candicidin macrolides was clearly sufficient to inhibit the growth of *E. weberi* (\varnothing 1.3-cm inhibition zone) and *E. aspergilloides* (\varnothing 0.6-cm inhibition zone) in the agar diffusion assay (Figs. S2–S4). At 4.5 nmol, however, even though growth is still visibly inhibited, the minimum concentration necessary for the inhibition of *E. weberi* to occur is reached. Candicidins seemed to inhibit *Escovopsis* specifically, because the growth of other pathogenic fungi was either not or only weakly inhibited (*Metarhizium anisopliae*, *Beauveria bassiana*, and *Lecanicillium lecanii*) (Figs. S5–S7). Importantly, the ants' symbiotic fungus *L. gongylophorus* is not significantly affected by candicidin macrolides (Fig. 2 and Fig. S8).

Discussion

To isolate an antifungal compound from microorganisms associated with leaf-cutting ants, we focused on *Streptomyces*-type bacteria. These microorganisms are both well known for their valuable bioactive secondary metabolites (25) and have previously been isolated from leaf-cutting ants (5, 14, 16). However, despite our selective isolation procedure, we obtained 19 strains, with 13 of them belonging to *Streptomyces* (Table S1 and Fig. S1). This supports the recent finding that diverse microbial communities are associated with leaf-cutting ants (14, 16) rather than a single coevolved *Pseudonocardia* strain, as previously believed (12, 13). Because conventional isolation techniques yield only a small proportion of the naturally occurring microorganisms, the microbial communities of leaf-cutting ants are likely extremely complex (26).

Because our initial screening for their antifungal potential against *Escovopsis* showed that some *Streptomyces* exhibited pronounced activity, we isolated the antifungal compound from the *Streptomyces*

sp. Ao10. We did not opt for a *Pseudonocardia* because most of our *Pseudonocardia* isolates did not appear to be very active against *Escovopsis*, an observation supported by previous studies (16). *Streptomyces* sp. Ao10 has been isolated directly from the *A. octospinosus* fungus garden. The presence of microorganisms producing antifungal compounds in the fungus garden that is the target of *Escovopsis* may provide the most efficient protection to the symbiotic fungus, whereas the bacterial biofilms on the body of the leaf-cutting ants may mainly protect individuals and help to distribute beneficial microorganisms. This conclusion is also supported by the observation that leaf-cutting workers preparing leaf material as a substrate for the symbiotic fungus inoculate the leaf material with microorganisms (27).

The bioassay-guided fractionation of the highly active *Streptomyces* sp. Ao10 yielded candicidin macrolides, which were highly active against *E. weberi* and *E. aspergilloides*. Candicidin macrolides were isolated in the 1950s (20), but elucidating their structure took until 1979 (21). Their name reflects the antifungal properties of the compounds against the human pathogenic fungus *Candida albicans*. Although candicidins are highly active antifungals, their medical use is restricted to severe cases (28). Like other polyketide polyenes, candicidins interact with sterols in the fungal cell membranes, causing K^+ -leakage and inducing cell death (29). The relatively rare development of resistance to polyketide polyenes among pathogenic fungi (28) might explain why leaf-cutting ants benefit from the presence of candicidin-producing microorganisms and do not face the risk that *Escovopsis* will adapt quickly. Recently, the candicidin biosynthetic gene cluster was characterized from both *Streptomyces* sp. FR-008 (23) and *S. griseus* (22), which allowed us to confirm its presence by PCR with gene-specific primers not only in *Streptomyces* sp. Ao10 but in total microbial DNA from individual *A. echinator* worker ants. Thus, we could directly prove that candicidin-producing microorganisms were on the ants' bodies without having to isolate the microorganisms. Ninety nanomoles of candicidin macrolides caused a large inhibition zone against *E. weberi* and *E. aspergilloides* in the agar diffusion assay (Fig. 2 and Figs. S2–S4); this zone was not overgrown even after 14 days of incubation. In the same assay, only 9 nmol of candicidin macrolides clearly inhibited the growth of *E. weberi* (Fig. S4). This result illustrates that candicidin macrolides effectively help leaf-cutting ants to defend their fungus gardens against *Escovopsis*. In contrast, candicidins did not inhibit the growth of the insect pathogenic fungi *M. anisopliae*, *B. bassiana*, and *L. lecanii* (Figs. S5–S7). In particular, the growth performance of the symbiotic fungus *L. gongylophorus* was not significantly altered by candicidins (Fig. S8).

Notably, of 19 bacterial isolates from leaf-cutting ants, 6 produced candicidin macrolides. In fact, we found candicidin-producing microorganisms associated with all leaf-cutting ant species analyzed (*A. octospinosus*, *A. echinator*, and *A. volcanus*). Moreover, the microbial screening by Mueller et al. (16) yielded an *S. griseus* strain from the fungus garden of *Cyphomyrmex muelleri*. This strain is closely related to our candicidin-producing *Streptomyces* strains, and thus is highly likely to produce candicidin macrolides. Consequently, candicidin-producing *Streptomyces* appear to be widespread in fungus gardens of leaf-cutting ants. In this context, it is interesting that Zhang et al. (19) concluded in their behavioral study that *Acromyrmex* ants are able to select for beneficial microorganisms. Furthermore, Poulsen et al. (30) suggest that the metapleural gland influences the growth of microbial symbionts on the ants' bodies. However, the chemical basis of both observations remains unknown.

Even though candicidin-producing *Streptomyces* sp. seem to be widespread among leaf-cutting ants, other antifungal compounds likely help to protect the fungus garden. *Pseudonocardia autotrophica*, which is closely related to the *Pseudonocardia* strains of leaf-cutting ants, has been found to contain a cryptic gene cluster specific for polyketide polyenes (31). However, we did not observe an antifungal polyene produced by our *Pseudonocardia* isolates. In addition to polyene macrolides, an as yet uncharacterized depsipeptide (32) and an unknown antifungal from *Burkholderia* (15) may contribute to the protection of the symbiotic fungus of different leaf-cutting ant species. These studies suggest that according to the microbial diversity among leaf-cutting ant-associated bacteria (14, 16), there is considerable variety of antifungal compounds.

In summary, with the identification of candicidins, we characterized an antifungal agent produced by the symbiotic bacteria of *Acromyrmex* leaf-cutting ants. Candicidins proved to be efficient agents against the fungal pathogen *Escovopsis*. The identification of candicidin macrolides as active antifungal compounds in the leaf-cutting ants' ecosystem opens a door for future studies of the chemical ecology of the microbial communities associated with leaf-cutting ants.

Materials and Methods

Fungal Cultures. *E. aspergilloides* CBS 423.93 and *E. weberi* CBS 110660 were obtained from the Centraalbureau voor Schimmelcultures. *L. gongylophorus* was an isolate from the fungus garden of *Atta colombica*. *M. anisopliae* DSMZ 1490, *B. bassiana* DSMZ 875, and *L. lecanii* DSMZ 3411 originated from the German Collection of Microorganisms and Cell Cultures. The strains were maintained on SFM agar plates.

Collection of the Ants and Fungus Garden Samples. Leaf-cutting ants and fungus garden samples were taken from *A. volcanus*, *A. octospinosus*, and *A. echinator* colonies (colonies A, B, C, and 1, respectively) collected in Gamboa, Panama. In addition, microorganisms were isolated from the fungus garden of an established laboratory colony of *A. echinator* (colony 2) collected in 2002 in Panama.

Isolation of Microorganisms. Microorganisms from the body of leaf-cutting ants were isolated by touching each ant with a sterile toothpick and drawing this over *Streptomyces* isolation agar (SIA) (17) medium agar plates. To collect microorganisms from the fungus garden, 2 mL of sterile water was added to a few milligrams of fungus garden. After samples were vortexed for 2 min, the aqueous supernatant was either plated directly onto SIA agar plates or diluted further before plating. After growing for 3 to 14 days, *Streptomyces*-type colonies were identified by their morphology, picked, and isolated by transfer onto new plates. Such pure colonies were maintained on SFM agar plates (20 g of soy flour, 20 g of mannitol, 20 g of agar, and 1 L of water) (25).

16S-rDNA Analysis of Isolated Symbionts. Genomic DNA was prepared from 3–7-day-old 25-mL cultures grown in liquid SFM medium following "procedure B" (25). For 16S rDNA analysis, the primers 8F (AGAGTTTGATCAGGCT-CAG) and 1492r (GGTACCTTGTACGACT) (33), fD2 (GAGTTTGATCAGGCT-CAG) (34), and 16Sr (TTGCGGGACTTAACCAACAT) (35) were used. The PCR

products were gel-purified and sequenced using the same primers. A minimum size of ca. 1,400 bp was used for database comparison (<http://greengenes.lbl.gov/cgi-bin/nph-index.cgi>, simrank algorithm). Phylogenetic analyses using the neighbor joining method (bootstrap value, $n = 1,000$) were conducted with MEGA version 4.1 (36).

Bioassay of the Antifungal Potential. *E. aspergilloides*, *E. weberi*, *M. anisopliae*, *B. bassiana*, *L. lecanii*, and *L. gongylophorus* were used as test organisms in the agar diffusion assay against isolated microorganisms, culture extracts, or pure samples. In the initial screening, the growth performance of *E. aspergilloides* was monitored on plates that 2 days earlier had been inoculated with one of the isolated microorganisms on one side of the plate. For the bioassay-guided fractionation, 100 μ L of mycelium or spore suspensions (ca. 5 mg wet weight/mL Luria–Bertani medium) of the test organisms was spread onto SFM plates (5.5-cm diameter). A 6-mm hole was cut in the middle of the plate, or a piece of filter paper was placed onto the agar plate, to apply 5–200 μ L of the test solution or an appropriate solvent control (e.g., MeOH, DMSO). To assay *L. gongylophorus*, a SFM agar piece (0.5 cm \times 0.5 cm) with the fully grown fungus was used to inoculate the test plate. The inhibition zones were monitored after 3 to 14 days at 28 $^{\circ}$ C. All assays were performed at least in triplicate and were compared with equally prepared solvent controls. Various amounts of purified candicidin macrolides dissolved in DMSO (2 mg/mL) were tested (4.5–90 nmol) (S): inhibition zones *E. weberi*: 90 nmol \varnothing 1.6 cm, 9 nmol \varnothing 1.3 cm, and 4.5 nmol \varnothing 0.6 cm; inhibition zones *E. aspergilloides*: 90 nmol \varnothing 1.4 cm, 45 nmol \varnothing 1.3 cm, 18 nmol \varnothing 1 cm, and 9 nmol \varnothing 0.6 cm.

Bioassay-Guided Fractionation and Structure Elucidation. *Streptomyces* sp. Ao10 was grown in 500-mL Erlenmeyer flasks fitted with springs for aeration and containing 200 mL of liquid SFM medium. The flasks were incubated at 28 $^{\circ}$ C on a rotating shaker (Infors Mutitron II MT25) shaker (200 rpm) for 7 days. From 2 to 10 L of culture was used for the bioassay-guided fractionation and purification of the antifungal compound. After harvesting, the cultures were centrifuged at 5,000 g for 30 min. The supernatant was extracted 3 times with an equal volume of 1-butanol. The extract was concentrated using a roti-vap. The residue was then resuspended in 20–50 mL of MeOH and subjected to SiO₂-column chromatography (60 M SiO₂, 50 cm \times 4 cm; Macherey–Nagel): 1 L of ethylacetate, 1 L of MeOH, and 1 L of water. Two hundred-milliliter fractions were collected, concentrated, redissolved in 4 mL of MeOH, and tested in the agar diffusion assay (50 μ L) against *E. aspergilloides*. Bioactive fractions were combined and subjected to RP 18 MPLC (40–63 μ m, LiChroprep RP18 resin, 45 cm \times 4 cm; Merck) using gradient elution at a flow rate of 10 mL: 100% A 0 min, in 30 min 100% B, 100% B 15 min (A: water, B: MeCN). Twenty-milliliter fractions were collected, concentrated, redissolved in 4 mL of MeOH, and used for the bioassay (50 μ L). The bioactive fraction was subjected to LC-MS analysis and HPLC separation using a Phenomenex Synergi Polar RP column (250 mm \times 2 mm, 5 μ m). A HP1100 system with a diode array detector (DAD) (Agilent) hooked to a Thermo Fisher LTQ or a Gilson 207 fraction collector was used. HPLC conditions were 3 min 100% A, in 27 min to 100% B, 10 min 100% B (A: water 0.1% AcOH, B: MeCN 0.1% AcOH), with a flow rate of 0.25 mL/min. Retention time of candicidin D was 25.4 min (UV: 408 nm, 384 nm, 364 nm, and 344 nm). High-resolution ESI-MS was performed with a Thermo Fisher Orbitrap injecting the purified samples directly via a syringe pump. Measured HR-ESI-MS was 1109.57938, and calculated HR-ESI-MS was 1109.57974 (C₅₉H₈₅O₁₈N₂). HR-ESI-MS/MS of 1109 was as follows: [M+H–H₂O]⁺ 1091.56827 (C₅₉H₈₃O₁₇N₂), [M+H–3H₂O]⁺ 1055.54762 (C₅₉H₇₉O₁₅N₂), [M+H–mycosamine–H₂O]⁺ 928.48458 (C₅₃H₇₀O₁₃N), 874.45283 (C₅₃H₆₄O₁₀N). Semipreparative HPLC was performed with a Grom-Sil ODS 5 ST RP18 column (250 mm \times 10 mm, 5 μ m; Alltech) using the fraction collector. HPLC conditions were 3 min 100% A, in 27 min to 100% B, 10 min 100% B (A: water 0.1% AcOH, B: MeCN 0.1% AcOH), at a flow rate of 3 mL/min and with a retention time of 20 min.

Screening Microbial Isolates for Candicidin Production. All microorganisms isolated from the different leaf-cutting ants (Table S1) were screened for their ability to produce candicidins. The strains were cultivated for 7 days in 100 mL of SFM medium at 28 $^{\circ}$ C/200 rpm. One milliliter of MeOH was added to 1 mL of the culture broth; the sample was then vortexed and centrifuged to pellet the cells. Fifty microliters of the supernatant was analyzed by LC-UV-MS.

Screening for Candicidin Biosynthetic Genes. Primers specific for the candicidin biosynthetic genes were used to amplify characteristic fragments from isolated microorganisms as well as directly from individual ants (*A. echinator*). DNA of microorganisms associated with individual ants was obtained by washing an ant in 1 mL of sterile water, transferring the resulting solution into a new Eppendorf tube, and centrifuging it. The pellet was utilized for DNA isolation using the Promega DNA Isolation Kit following its protocol for genomic DNA isolation from Gram-positive microorganisms. About 1,000 kb

regions from fscA, fscM, and fscP were amplified by PCR using fscAfor (ATG-GTGCCCGTCCACGCACACGACTACGTGACCGATCCGC) and fscArev (GGCGGC-CAGCACCTCGGGCAGCGGGTACCACCAG), fscMfor (TCGCTGGGCGAGGTG-GTCCCGAACTGT) and fscMrev (CGGCTTGTCAGGGTACGGTGTGATGCCG), and fscPfor (GGTTCTGGCCAGGCACTGGTGGACGCCGTAGCC) and fscPprev (ATGACGACCAGCCCCGCCCCGACGGTGTGCTGACTTCCAC). PCR products were gel-purified and sequenced using the gene-specific primers. The sequences were compared by Blast search to the corresponding genes from *Streptomyces* sp. FR-008 (23) and *S. griseus* (22).

- Hölldobler B, Wilson EO (1990) *The Ants* (Springer, Berlin).
- Wirth R, Beyschlag W, Ryel R, Herz H, Hölldobler B (2003) *Ecological Studies* (Springer, Berlin).
- Cherrett JM (1986) The biology, pest status and control of leaf cutting ants. *Agric Zool Rev* 1:1–27.
- Seifert KA, Samson RA, Chapela IH (1995) *Escovopsis aspergilloides*, a rediscovered hyphomycete from leaf-cutting ant nests. *Mycologia* 87:407–413.
- Currie CR, Mueller UG, Malloch D (1999) The agricultural pathology of ant fungus gardens. *Proc Natl Acad Sci USA* 96:7998–8002.
- Currie CR, Stuart AE (2001) Weeding and grooming of pathogens in agriculture by ants. *Proc R Soc Lond Ser B* 268:1033–1039.
- Ortius-Lechner D, Maile R, Morgan ED, Boomsma JJ (2000) Metapleural gland secretion of the leaf-cutter ant *Acromyrmex octospinosus*: New compounds and their functional significance. *J Chem Ecol* 26:1667–1683.
- Schildknecht H, Koob K (1971) Myrmicacin, the first insect herbicide. *Angew Chem Int Ed Engl* 10:124–125.
- Bot ANM, Ortius-Lechner D, Finster K, Maile R, Boomsma JJ (2002) Variable sensitivity of fungi and bacteria to compounds produced by the metapleural glands of leaf-cutting ants. *Insect Soc* 49:363–370.
- Fernandez-Marin H, Zimmerman JK, Rehner SA, Wcislo WT (2006) Active use of the metapleural glands by ants in controlling fungal infection. *Proc R Soc London Ser B* 273:1689–1695.
- Currie CR, Scott JA, Summerbell RC, Malloch D (1999) Fungus-growing ants use antibiotic-producing bacteria to control garden parasites. *Nature* 398:701–704.
- Currie CR, et al. (2003) Ancient tripartite coevolution in the attine ant-microbe symbiosis. *Science* 299:386–388.
- Currie CR, Bot ANM, Boomsma JJ (2003) Experimental evidence of a tripartite mutualism: Bacteria protect ant fungus gardens from specialized parasites. *Oikos* 101:91–102.
- Kost C, et al. (2007) Non-specific association between filamentous bacteria and fungus-growing ants. *Naturwissenschaften* 94:821–828.
- Santos AV, Dillon RJ, Dillon VM, Reynolds SE, Samuels RI (2004) Occurrence of the antibiotic-producing bacterium *Burkholderia* sp. in colonies of the leaf-cutting ant *Atta sexdens rubropilosa*. *FEMS Microbiol Lett* 239:319–323.
- Mueller UG, Dash D, Rabeling C, Rodrigues A (2008) Coevolution between attine ants and actinomycete bacteria: A reevaluation. *Evolution* 62:2894–2912.
- Atlas RM (1997) *Handbook of Microbiological Media* (CRC Press, Boca Raton, FL).
- Oroshnik W, Vining LC, Mebane AD, Taber WA (1955) Polyene antibiotics. *Science* 121:147–149.
- Zhang MM, Poulsen M, Currie CR (2007) Symbiont recognition of mutualistic bacteria by *Acromyrmex* leaf-cutting ants. *ISME J* 1:313–320.
- Lechevalier H, Acker RF, Corke CT, Haenseler CM, Waksman SA (1953) Candicidin, a new antifungal antibiotic. *Mycologia* 45:155–171.
- Zielinski J, et al. (1979) Structure of levorin-A2 And candicidin-D. *Tetrahedron Lett* 20:1791–1794.
- Campelo AB, Gil JA (2002) The candicidin gene cluster from *Streptomyces griseus* IMRU 3570. *Microbiology* 148:51–59.
- Chen S, et al. (2003) Organizational and mutational analysis of a complete FR-008/candicidin gene cluster encoding a structurally related polyene complex. *Chem Biol* 10:1065–1076.
- Zhou YJ, et al. (2008) Incomplete beta-ketone processing as a mechanism for polyene structural variation in the FR-008/candicidin complex. *Chem Biol* 15:629–638.
- Kieser T, Bibb MJ, Buttner MJ, Chater KF, Hopwood DA (2000) *Practical Streptomyces Genetics* (Crowes, Norwich, UK), p 169.
- Amann RI, Ludwig W, Schleifer KH (1995) Phylogenetic identification and in-situ detection of individual microbial-cells without cultivation. *Microbiol Rev* 59:143–169.
- Mangone DM, Currie CR (2007) Garden substrate preparation behaviours in fungus-growing ants. *Can Entomol* 139:841–849.
- Zotchev SB (2003) Polyene macrolide antibiotics and their applications in human therapy. *Curr Med Chem* 10:211–223.
- Hammond SM, Kliger BN (1976) Mode of action of polyene antibiotic candicidin-binding factors in wall of *Candida albicans*. *Antimicrob Agents Chemother* 9:561–568.
- Poulsen M, Bot ANM, Boomsma JJ (2003) The effect of metapleural gland secretion on the growth of a mutualistic bacterium on the cuticle of leaf-cutting ants. *Naturwissenschaften* 90:406–409.
- Lee MY, Myeong JS, Park HJ, Han K, Kim ES (2006) Isolation and partial characterization of a cryptic polyene gene cluster in *Pseudonocardia autotrophica*. *J Ind Microbiol Biotechnol* 33:84–87.
- Oh DC, Scott JJ, Poulsen MZ, Currie CR, Clardy J (2008) Discovery of new secondary metabolites mediating insect-microorganism symbioses. *Planta Med* 74:906–906.
- Desai C, Madamwar D (2007) Extraction of inhibitor-free metagenomic DNA from polluted sediments, compatible with molecular diversity analysis using adsorption and ion-exchange treatments. *Bioresour Technol* 98:761–768.
- Bereswill S, Bugert P, Bruchmüller I, Geider K (1995) Identification of the fire blight pathogen, *Erwinia amylovora*, by PCR assays with chromosomal DNA *Appl Environ Microbiol* 61:2636–2642.
- Aittamaa M, et al. (2008) Distinguishing bacterial pathogens of potato using a genome-wide microarray approach. *Molecular Plant Pathology* 9:705–717.
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 24:1596–1599.

ACKNOWLEDGMENTS. We thank the Smithsonian Tropical Research Institute for providing logistical help, and the Autoridad Nacional del Ambiente y el Mar for permission to sample ant colonies in Panama and for issuing export permits. We are grateful to Prof. Dr. Michael Spiteller (Institut für Umweltforschung, Dortmund) for the HR-ESI-MS measurements and to Dr. Ute Möllmann (Hans Knöll Institute, Jena) for a candicidin D standard. D.S. thanks Prof. Dr. Wilhelm Boland for his generous support and the Deutsche Forschungsgemeinschaft for an Emmy Noether Fellowship (SP 1106/3–1). Additional funding by the Verband der Chemischen Industrie and the Max Planck Society is gratefully acknowledged.