

# Isotopic evidence for the diets of European Neanderthals and early modern humans

Michael P. Richards<sup>a,b,1</sup> and Erik Trinkaus<sup>c</sup>

<sup>a</sup>Department of Human Evolution, Max-Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, D-04103 Leipzig, Germany; <sup>b</sup>Department of Anthropology, University of British Columbia, Vancouver, British Columbia V6T 1Z1, Canada; and <sup>c</sup>Department of Anthropology, Campus Box 1114, Washington University, St. Louis, MO 63130

Edited by Richard G. Klein, Stanford University, Stanford, CA, and approved June 23, 2009 (received for review April 7, 2009)

**We report here on the direct isotopic evidence for Neanderthal and early modern human diets in Europe. Isotopic methods indicate the sources of dietary protein over many years of life, and show that Neanderthals had a similar diet through time ( $\approx 120,000$  to  $\approx 37,000$  cal BP) and in different regions of Europe. The isotopic evidence indicates that in all cases Neanderthals were top-level carnivores and obtained all, or most, of their dietary protein from large herbivores. In contrast, early modern humans ( $\approx 40,000$  to  $\approx 27,000$  cal BP) exhibited a wider range of isotopic values, and a number of individuals had evidence for the consumption of aquatic (marine and freshwater) resources. This pattern includes Oase 1, the oldest directly dated modern human in Europe ( $\approx 40,000$  cal BP) with the highest nitrogen isotope value of all of the humans studied, likely because of freshwater fish consumption. As Oase 1 was close in time to the last Neanderthals, these data may indicate a significant dietary shift associated with the changing population dynamics of modern human emergence in Europe.**

Europe | isotopes | collagen | fishing

Isotope evidence is a powerful tool for reconstructing past human diets and subsistence adaptations (1–3), and it has been applied to a number of Neanderthals and early modern humans from Europe (4–12). In 2 earlier studies (10, 11), we argued that Neanderthals had relatively uniform dietary adaptations while early modern humans in Europe had more variable isotope values—and therefore diets—than the Neanderthals. Specifically, we proposed that a number of European early modern humans had higher nitrogen isotope values than any Neanderthal and that these values likely indicated that some of these humans were obtaining much of their protein from aquatic resources, namely freshwater fish (11). This interpretation was largely based on the similarity of these isotope values to those of freshwater fish consumers from Mesolithic Eastern Europe (13, 14). We further postulated that the isotope evidence demonstrated a shift to a broader dietary spectrum by early modern humans (11), which probably included small game, in addition to fish, an inference supported by the faunal evidence (15, 16). Since that study, there have been more isotopic studies of Neanderthals (5–9) and early modern humans (4, 17, 18). Importantly, we present here previously unrecorded isotopic data from an early modern human from the Peçstera cu Oase (4), which overlaps in time with the last Neanderthals (19), allowing us a direct comparison between Neanderthal and modern human diets when they were both present in Europe. Below, we summarize the current isotopic evidence for Neanderthal and modern human diets and suggest that the previously unrecorded data support our original inference that there was a shift in dietary spectra between the Neanderthals and early modern humans in Europe.

**Isotopic Analysis.** Carbon and nitrogen stable isotope ratios of adult human bone collagen are indicators of the main sources of dietary protein consumed over a number of years (20, 21). Carbon isotope ratios ( $\delta^{13}\text{C}$  values) can indicate if the source of

dietary protein was from marine resources or terrestrial resources (22, 23), as there is an  $\approx 7$  per mil (‰) shift between dissolved ocean bicarbonate and atmospheric carbon dioxide (the main respective sources of carbon for plants in each ecosystem) (24, 25). Therefore, exclusively marine consumers, such as seals, have  $\delta^{13}\text{C}$  values of  $-12 \pm 1\text{‰}$ , and this value is relatively constant around the world, including in the oceans surrounding Europe (26). Terrestrial consumers have  $\delta^{13}\text{C}$  values close to  $-20 \pm 2\text{‰}$ , depending on the region (27). Organisms in freshwater ecosystems generally have carbon values similar to terrestrial values, but these values are highly variable because dissolved carbon in rivers and lakes can derive from many geological sources with differing carbon isotope ratios (27, 28). Carbon isotope values can also be used to discriminate between the consumption of  $\text{C}_3$  and  $\text{C}_4$  photosynthetic pathway plants (29); however, there were no edible  $\text{C}_4$  plants in Europe during the Late Pleistocene.

Nitrogen isotope ratios ( $\delta^{15}\text{N}$  values) of human bone collagen are between 3 to 5‰ higher than dietary protein (30, 31). This “trophic level effect” is widely used in archeology and modern ecology as a means to determining the positions of organisms in food webs, particularly for identifying the prey species of carnivores as well as the diets of omnivores, such as humans (27). The basis of this pattern is that plants obtain nitrogen from the soil or atmosphere (24), and in Holocene Europe plants generally had  $\delta^{15}\text{N}$  values of between 0 and 2‰. Herbivores that consume plants have body protein (flesh and bone collagen) that is 3 to 5‰ higher than the plants (i.e., 3 to 7‰). Carnivores that consume those herbivores have bone collagen nitrogen isotope values that are again 3 to 5‰ higher than the herbivores (i.e., 6 to 12‰). Omnivore values can fall between the ranges of carnivore and herbivore values depending on the amount of plant vs. animal protein in their diets. Therefore, by comparing the isotope values of omnivores such as humans to the isotope values of herbivores and carnivores from the same site or region, we can determine whether they obtained their proteins from mainly animal sources (carnivore-like) or plant sources (herbivore-like), or a mix of both. The same process applies in aquatic (freshwater and marine) ecosystems. However, as there are often many more steps in the food chain in aquatic ecosystems, top-level aquatic consumers often have much higher  $\delta^{15}\text{N}$  values [i.e., seals have  $\delta^{15}\text{N}$  values of 18 to 20‰ (26)] than their terrestrial counterparts [i.e., wolves generally have values between 10 and 12‰ (9)].

A confounding factor in the use of stable isotope analysis to reconstruct past diets is that carbon and nitrogen isotope values vary between different geographical regions, especially related


Author contributions: M.P.R. and E.T. designed research; M.P.R. and E.T. performed research; M.P.R. and E.T. analyzed data; and M.P.R. and E.T. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

<sup>1</sup>To whom correspondence should be addressed. E-mail: richards@eva.mpg.de.

This article contains supporting information online at [www.pnas.org/cgi/content/full/0903821106/DCSupplemental](http://www.pnas.org/cgi/content/full/0903821106/DCSupplemental).



**Fig. 1.** Carbon and nitrogen isotope values of bone collagen from Neanderthals and early modern humans from Europe. Errors on the isotope measurements are typically  $\pm 0.2\%$  for both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ .


to temperature and aridity (24, 32, 33). In Europe, the climate is relatively similar so, to date, there have not been any observed large-scale differences in carbon and nitrogen isotopes of herbivores and carnivores across Europe, especially in the Holocene [where most of the application of isotope analysis has occurred (34)]. However, the climate has changed considerably in the past, and it now appears that these climate changes had an effect on isotope values, especially during and just after the Late Glacial Maximum, when herbivores had  $\delta^{15}\text{N}$  values of  $\geq 3\%$  lower than in the preceding marine isotope stage (MIS) 3 and subsequent Holocene period (34–36). Of note here is that these studies did not observe a significant increase in  $\delta^{15}\text{N}$  values in the period from 50,000 to 25,000 cal BP. However, the fauna in these studies was exclusively from Northern Europe, and mostly from the United Kingdom, so it is not a valid comparative framework for Eastern and Southern Europe. Regardless, to best interpret human carbon and nitrogen isotope values we should compare the human isotope values to the isotope values of associated fauna that are as close in time and space as possible, and ideally from the same site and stratigraphic context.

## Results

**Neanderthal Isotope Values.** There are currently 13  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values for European adult Neanderthals [Fig. 1 and [supporting information \(SI Table S1\)](#)]. These Neanderthals range in age from  $\approx 120,000$  BP (Scladina) to  $\approx 37,000$  cal BP (Vindija), but most of them are  $< 50,000$  cal BP. We have excluded the Engis juvenile Neanderthal (8), as there are a number of physiological effects on juvenile isotopic values, such as breastfeeding and weaning (37–39), and therefore juvenile data cannot be directly compared with adult values. Following Bocherens et al. (9), we have excluded 2 of the Neanderthals from Les Pradelles (Marillac) (12), as they apparently have poor collagen preservation (40). Additionally, as we are focusing only on Europe, we do not include the Siberian Okladnikov Neanderthal ( $\delta^{13}\text{C} = -19.1\%$ ,  $\delta^{15}\text{N} = 12.9\%$ ) (41).

The carbon isotope values for all of the Neanderthals (see Fig. 1 and [Table S1](#)) are all  $< -19\%$ , indicating that dietary protein was from terrestrial sources. Therefore, there is no evidence that marine foods (which have higher  $\delta^{13}\text{C}$  values,  $\approx -12 \pm 1\%$ ) were consumed in any significant quantities by these Neanderthals, few of which were in close proximity to coastlines.

Because of the potentially fluctuating faunal isotope values over this time period [for which we do not have the baseline faunal data that we do for later periods (34)], we cannot directly




**Fig. 2.** Carbon and nitrogen isotope data for the Oase 1 human and associated fauna taken from Trinkaus et al. (4) and the present study. Errors on the isotope measurements are typically  $\pm 0.2\%$  for both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ .

compare these Neanderthal nitrogen isotope values to each other. However, in each analysis in which the Neanderthals were compared to contemporary faunal isotope values from the same site or contemporary adjacent sites, the conclusions were the same. Each Neanderthal had  $\delta^{15}\text{N}$  values that were 3 to 5‰ higher than contemporary herbivores and similar to carnivores (or in some cases slightly higher). In each study, the authors concluded that Neanderthals were top-level carnivores and that their main protein source was large herbivores.

**Early Modern Human Isotope Values.** There are 14 modern humans from the European earlier (Early and Mid) Upper Paleolithic (MIS 3) that have carbon isotope values, and 10 which have both carbon and nitrogen isotope values (see Fig. 1, [Table S2](#)). The Oase 1 modern human is the oldest directly dated modern human in Europe [ $\approx 40,000$  cal BP (42)] and the only one in our study that overlaps in time with Neanderthals. The other early modern humans from the Early- to Mid-Upper Paleolithic with isotopic values date to between  $\approx 34,000$  and  $\approx 27,000$  cal BP and are (or are likely to be) associated with late Aurignacian or especially Gravettian technology.

The Oase 1 human carbon and nitrogen values are plotted with isotope results from associated faunal remains in Fig. 2. The human and faunal remains were largely recovered from surface deposits in the cave, and therefore represent a range of time periods dating to between  $\approx 50,000$  (wolf, hyena, and red deer) and  $\approx 20,000$  (ibex) cal BP (43). The herbivore isotope values are similar, despite their likely range of ages. The highest wolf  $\delta^{15}\text{N}$  value is 11.5‰, which is 8.9‰ higher than the *Capra* (ibex) (average  $2.6 \pm 0.5\%$ ) and 6.1‰ higher than the *Cervus* (red deer) (average 5.4‰), while a hyena has a value of 11.1‰, which is 8.5‰ higher than the ibex and 5.7‰ higher than the deer. As there is an enrichment of between 3 to 5‰ in  $\delta^{15}\text{N}$  between prey and consumer, the wolves and the hyena were likely obtaining most of their protein from the red deer and not the ibex at this site. In contrast, Oase 1 has a  $\delta^{15}\text{N}$  value (13.3‰) that is 10.8‰ higher in  $\delta^{15}\text{N}$  than the ibex and 8.0‰ higher than the red deer. The enrichment between both herbivores and Oase 1 is far beyond the 3 to 5‰ trophic level effect in  $\delta^{15}\text{N}$ . The Oase 1  $\delta^{15}\text{N}$  value is also above those of the hyena (11.1‰), and the highest wolf value (11.5‰) from the same site and dating to about the same time. Therefore, Oase 1 must have obtained a significant portion of its protein from a different ecosystem, for which the best candidate is freshwater fish.

Further evidence for the use of aquatic foods by early modern



**Fig. 3.**  $\delta^{13}\text{C}$  bone collagen values of directly radiocarbon-dated Neanderthals and early modern humans compared to the average  $\delta^{13}\text{C}$  values of directly radiocarbon-dated herbivores from Northern Europe (34) over the period 50,000 to 20,000 cal BP.

humans comes from the isotope values of 2 later coastal Mid Upper Paleolithic modern humans, Arene Candide IP from coastal Italy (18) and La Rochette 1 from further inland in France (44). Both of these individuals have carbon isotope values  $> -18\text{‰}$ , which indicate that they consumed significant (i.e., 20–30% of dietary protein) amounts of marine foods, probably high trophic level (carnivorous) fish or marine mammals. These values are evident in a plot of  $\delta^{13}\text{C}$  values of the Neanderthals and early modern humans (Fig. 3) that have direct radiocarbon dates, combined with the  $\delta^{13}\text{C}$  values compiled in a large-scale survey of directly radiocarbon-dated European faunal isotope values (34). These 2 humans have very different carbon isotope values from all of the other early modern humans from Europe and all of the Neanderthals (see Fig. 1, Tables S1 and S2), and the associated faunal data shows that there was no unrecognized shift in terrestrial  $\delta^{13}\text{C}$  values at this time. Therefore, these 2 individuals most likely consumed significant amounts of marine protein. This interpretation is also supported by the associated high  $\delta^{15}\text{N}$  values (12.4‰ for Arene Candide IP, 11.2‰ for La Rochette 1), as the  $\delta^{13}\text{C}$  values of marine food consumers are correlated with an increase in  $\delta^{15}\text{N}$  values (26, 45).


The other early modern humans all have  $\delta^{13}\text{C}$  values  $< -18.5\text{‰}$  (see Fig. 1 and Table S2), which indicate that their protein came from terrestrial  $\text{C}_3$  (or freshwater) foods, yet many of them have high  $\delta^{15}\text{N}$  values, at or above the highest Neanderthal values. To best interpret these high human  $\delta^{15}\text{N}$  values, they should be compared to associated faunal remains. However, in contrast to the Neanderthals, very few of these humans have associated faunal isotope values. This situation is partly because of the nature of their discovery and excavation, as many of them are from deliberate burials, were not found in association with fauna, and were excavated in the 19<sup>th</sup> or early 20<sup>th</sup> century without associated faunal remains or have isotopic data from direct radiocarbon dating. Unfortunately, unlike the  $\delta^{13}\text{C}$  values, we cannot use the average faunal  $\delta^{15}\text{N}$  values in studies of directly radiocarbon dated fauna from this period, as these data are exclusively from Northern Europe and so cannot be compared to humans from Southern and Eastern Europe (contra ref. 46). Nonetheless, a plot of those faunal  $\delta^{15}\text{N}$  data with the human

values (Fig. 4) shows no consistent shift across mammalian families in average  $\delta^{15}\text{N}$  values.

The Eel Point 1 human from South Wales has a  $\delta^{15}\text{N}$  value of 11.4‰. This value can be compared to those of approximately contemporary fauna from the nearby site of Paviland (47). Two bison bones from Paviland have an average  $\delta^{15}\text{N}$  value of 8.4‰. The  $\delta^{15}\text{N}$  value of Eel Point 1 is 3‰ higher than the average  $\delta^{15}\text{N}$  values of these 2 herbivores, which is within the 1 trophic level increase range. Therefore, Eel Point 1 most likely obtained most of its dietary protein from herbivores, such as bison and horse. The human has very different carbon isotope values than a contemporary brown bear ( $\delta^{13}\text{C} = -17.3\text{‰}$ ,  $\delta^{15}\text{N} = 11.3\text{‰}$ ), which likely consumed marine foods such as salmon.

The human from Paviland itself has a lower  $\delta^{15}\text{N}$  value (10.4‰) than Eel Point 1 and also has a more positive  $\delta^{13}\text{C}$  value ( $-18.4\text{‰}$ ). This human was originally interpreted as having a minor, but detectable, contribution of marine protein in its diet (47). Redating of Paviland 1 to  $\approx 33,000$  cal BP (48) means that we can now compare the human isotope values to fauna dating to an earlier period. There are 3 faunal bone samples dating to  $\approx 33,000$  to  $\approx 34,000$  cal BP, a fox ( $\delta^{13}\text{C} = -19.6\text{‰}$ ,  $\delta^{15}\text{N} = 9.4\text{‰}$ ), a bison ( $\delta^{13}\text{C} = -19.9\text{‰}$ ,  $\delta^{15}\text{N} = 6.2\text{‰}$ ), and a brown bear ( $\delta^{13}\text{C} = -18.8\text{‰}$ ,  $\delta^{15}\text{N} = 10.6\text{‰}$ ). The fox and bison have terrestrial  $\delta^{13}\text{C}$  values, so there is no unusual effect at this site and time to explain the more positive  $\delta^{13}\text{C}$  values of Paviland 1. Therefore, based on both the  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of Paviland 1, this human likely had a marine component in its diet, and indeed it has isotope values similar to the contemporary brown bear that was likely to have been a marine consumer.

Several other Early Upper Paleolithic humans have high  $\delta^{15}\text{N}$  values, higher than any of the Neanderthals (see Fig. 1, Table S1 and S2). These high values may, similar to Oase, indicate the consumption of freshwater foods, but without comparative faunal isotope values it is not possible to definitively conclude this inference. Yet, if we compare the 3 other humans from Romania, Muierii 1 and 2, and Cioclovina 1, with  $\delta^{15}\text{N}$  values of 12.3‰, 12.4‰, and 12.7‰, respectively (average: 12.5‰), with the herbivore values from Oase, they are  $\approx 10\text{‰}$  higher than the ibex and 7‰ higher than the deer. Likewise, if we compare the  $\delta^{15}\text{N}$  values of the 2 humans from the Czech



**Fig. 4.**  $\delta^{15}\text{N}$  bone collagen values of directly radiocarbon-dated Neanderthals and early modern humans compared to the average  $\delta^{15}\text{N}$  values of directly radiocarbon dated herbivores from Northern Europe (34) over the period 50,000 to 20,000 cal BP.

republic (Dolní Věstonice 35 and Brno-Francouzská 1), both of which have  $\delta^{15}\text{N}$  values of 12.3‰, with the single, contemporary, herbivore bone from Brno-Francouzská ( $\delta^{15}\text{N}$  value of 5.2‰) (4), the humans are 7.1‰ higher than the herbivore from Brno-Francouzská. In both of these cases the humans have  $\delta^{15}\text{N}$  values that are more than the observed trophic level effect (3–5‰); an explanation may be the inclusion of significant amounts of freshwater resources in their diets. However, these interpretations are preliminary until associated faunal values can be obtained from these 4 sites.

## Discussion

Isotopic analysis provides information about the sources of dietary protein over a number of years, even though it does not measure the caloric contributions of different foods. As the method only measures protein intake, many low-protein foods that may have been important to the diet (i.e., high caloric foods like honey, underground storage organs, and essential mineral and vitamin rich plant foods) are simply invisible to this method. There are high-protein plant foods in Europe that Neanderthals and early modern humans could have consumed, such as hazelnuts [commonly consumed in the Mesolithic (49)] that would have been visible in the isotopic analyses, but they are simply not seen. Another important consideration is that this method tracks the bulk protein consumption over a number of years, and provides an average and proportional measure of the protein sources. Therefore, the occasional consumption of resources like fish or marine mammals by Neanderthals (i.e., once a month, 1 month a year) would be largely invisible to this method. Indeed, it is evident that Neanderthals sometimes consumed aquatic resources (50), although the isotope evidence shows that it was not likely to have been an important part of their diet across Neanderthal populations.

Despite these caveats, isotopic analysis is the only direct measure of the protein sources in past diets, and it allows reconstruction of this aspect of Neanderthal and early modern human diets in Europe. The isotopic evidence for Neanderthal diets is notable for its consistency. Although the Neanderthals that have been studied to date come from different regions of Europe and time periods, the isotopic data show that, in each

case, they have  $\delta^{15}\text{N}$  values between  $\approx 3$  and 5‰ higher than the local herbivores, and plot close to carnivores from the same or nearby sites. These higher  $\delta^{15}\text{N}$  values indicate that European Neanderthals had similar dietary adaptations.

Early modern humans in Europe have a more varied range of isotopic values that indicates that some of them consumed significant quantities of aquatic foods, both from freshwater and marine sources. The only human contemporary with the Neanderthals that currently provides isotopic data, Oase 1, has  $\delta^{15}\text{N}$  values that are the highest of all of the modern humans and higher than all of the Neanderthal values.

The isotope data agree well with results from faunal analyses. Studies of animal remains from Neanderthal sites in Europe have repeatedly shown that Neanderthals consistently hunted large herbivores (51, 52), including seasonal use of reindeer (53). Yet, there is little evidence for the use of small game such as birds or fish (4, 51, 54). Early modern humans also appear to have regularly hunted large herbivores (55–57), but there is also evidence for the use of small game, including fish at some of these sites (15, 16).

The observed isotope difference between Neanderthals and early modern humans may therefore indicate that there were dietary spectra differences between MIS 3 modern humans in Europe and the Neanderthals. Modern humans may well have had a broader dietary spread than Neanderthals, but we need more data from areas where Neanderthals and modern humans overlapped to confirm this inference. It is nonetheless clear that modern humans, in the relatively short period covered by this study (40,000 to 27,000 cal BP), were exhibiting a wide range of diets, while the Neanderthals seem to have had the same general dietary adaptation throughout a much longer time range (120,000 to 37,000 cal BP).

Why do we see these dietary differences between Neanderthals and modern humans in Europe? It is tempting to see the difference as being somehow cognitive, in that Neanderthals were unable to alter their (albeit very successful) subsistence strategies, whereas modern humans were more creative and were able to exploit resources more than the Neanderthals. However, the difference need not be cognitive and instead an increased flexibility in modern human diets could be part of the techno-

logical and organizational package that allowed the widespread expansion of modern humans out of Africa into Europe  $\approx 40,000$  years ago. Indeed, the first clear evidence for the regular use of marine resources comes from modern humans in Africa only after the middle stone age/late stone age transition  $\approx 40,000$  BP (58), although, as in Europe, there is scattered earlier evidence of fish consumption in Africa (59). Therefore, the differences in diets may simply be linked to adaptive and technological changes between these 2 time periods, and their cultural complexes, probably analogous to the clear dietary changes from marine to terrestrial foods at the Mesolithic to Neolithic transition in Northern Europe (60, 61). Moreover, the relatively rapid dispersal of modern humans into most of Europe and the evidence for limited if widespread assimilation of Neanderthals into those populations (62) imply markedly higher effective population sizes of those early modern humans. Larger modern human populations may well have promoted the variable exploitation of a broader range of resources, ones requiring greater effort or technological investment for their acquisition.

Stable isotope analysis is therefore a powerful method for reconstructing aspects of past diets, and it has been especially

useful in determining the protein sources in Neanderthal and early modern human diets in Europe. There are now enough isotopic data to see patterns in the data, and they show that the Neanderthals and early modern humans had similar dietary adaptations, obtaining most of their dietary protein from animals, although some of the early modern humans obtained significant amounts of their protein from aquatic, and not just terrestrial, sources. Ongoing work on Neanderthal and early modern human isotopic profiles will continue to test whether this difference is real; work in other regions with different available resources, such as the Mediterranean, Russia, and China, will test whether this pattern was more widespread. We will also soon be able to apply much needed new isotopic measurements to evaluate the evidence for the consumption of aquatic resources, such as single amino acid carbon isotope analysis (63) and sulfur isotope analysis (64–66).

**ACKNOWLEDGMENTS.** The authors thank the many collaborators who kindly provided material for analysis. This work was funded in part by the Max Planck Society (M.P.R.).

- Lee-Thorp JA (2008) On isotopes and old bones. *Archaeometry* 50:925–950.
- Lee-Thorp JA, Sponheimer M (2006) Contributions of biogeochemistry to understanding hominin dietary ecology. *Yearb Phys Anthropol* 49:131–148.
- Sealy J (2001). Body tissue chemistry and Palaeodiet. In Brothwell DR, Pollard AM (eds.) *Handbook of Archaeological Sciences*. (Chichester, John Wiley and Sons) pp. 269–279.
- Trinkaus E, et al. (2009) Stable isotope evidence for early modern human diet in Southeastern Europe: Peștera cu Oase, Peștera Muierii and Peștera Cioclovina Uscată. *Materiale 'i Cercetări Arheologice*, in press.
- Richards MP, Schmitz R (2008) Isotope evidence for the diet of the Neanderthal type specimen. *Antiquity* 82:553–559.
- Richards MP, et al. (2008) Isotopic dietary analysis of a Neanderthal and associated fauna from the site of Jonzac (Charente-Maritime), France. *J Hum Evol* 55:179–185.
- Beauval C, Lacrampe-Cuyaubere F, Maureille B, Trinkaus E (2006) Direct radiocarbon dating and stable isotopes of the Neanderthal femur from Les Rochers de Villeneuve. *Bulletins et Memoires de la Societe d'Anthropologie de Paris* 18:35–42.
- Bocherens H, et al. (2001) New isotopic evidence for dietary habits of Neanderthals from Belgium. *J Hum Evol* 40(6):497–505.
- Bocherens H, Drucker D, Billiou D, Patou-Mathis M, Vandermeersch B (2005) Isotopic evidence for diet and subsistence pattern of the Saint-Ce'saire I Neanderthal: Review and use of a multi-source mixing model. *J Hum Evol* 49:71–87.
- Richards MP, et al. (2000) Neanderthal diet at Vindija and Neanderthal predation: The evidence from stable isotopes. *Proc Nat Acad Sci USA* 97:7663–7666.
- Richards MP, Pettitt PB, Stiner MC, Trinkaus E (2001) Stable isotope evidence for increasing dietary breadth in the European mid-Upper Paleolithic. *Proc Nat Acad Sci USA* 98:6528–6532.
- Fizet M, et al. (1995) Effect of diet, physiology and climate on carbon and nitrogen stable isotopes of collagen in Late Pleistocene anthropic palaeoecosystem: Marillac, Charente, France. *J Archaeol Sci* 22:67–79.
- Bonsall C, et al. (1997) Mesolithic and early Neolithic in the Iron Gates: A palaeodietary perspective. *J Eur Archaeol* 5:1:50–92.
- Cook GT, et al. (2001). A freshwater diet-derived C-14 reservoir effect at the stone age sites in the iron gates gorge. *Radiocarbon* 43(2A):453–460.
- Muñoz M, Casadevall M. (1998) Fish remains from Arbreda Cave (Serinyà, Girona), northeast Spain, and their palaeoecological significance. *J Quaternary Sci* 12:111–115.
- Stiner MC (1999) Palaeolithic mollusc exploitation at Riparo Mochi (Balzi Rossi, Italy): Food and ornaments from the Aurignacian through Epigravettian. *Antiquity* 73:735–754.
- Schulting RJ, et al. (2005) A Mid-Upper Palaeolithic human humerus from Eel Point, South Wales, UK. *J Hum Evol* 48:493–505.
- Pettitt PB, Richards MP, Maggi R, Formicola V (2003) The Gravettian burial known as the Prince ('Il Principe'): New evidence for his age and diet. *Antiquity* 295:15–19.
- Higham T, Bronk Ramsey C, Karavanic I, Smith FH, Trinkaus E (2006) Revised direct radiocarbon dating of the Vindija G1 Upper Paleolithic Neanderthals. *Proc Nat Acad Sci USA* 103:553–557.
- Ambrose SH, Norr L. (1993) Experimental evidence for the relationship of the carbon isotope ratios of whole diet and dietary protein to those of bone collagen and carbonate. In Lambert J, Grupe G, eds. *Prehistoric Human Bone: Archaeology at the Molecular Level* (Springer-Verlag, New York) pp. 1–37.
- Schwarcz H, Schoeninger M (1991) Stable isotope analyses in human nutritional ecology. *Yearb Phys Anthropol* 34:283–321.
- Schoeninger M, DeNiro M, Tauber H (1983) Stable nitrogen isotope ratios of bone collagen reflect marine and terrestrial components of prehistoric human diet. *Science* 220:1381–1383.
- Chisholm BS, Nelson DE, Schwarcz HP (1982) Stable carbon ratios as a measure of marine versus terrestrial protein in ancient diets. *Science* 216:1131–1132.
- Hoefs J (2009) *Stable Isotope Geochemistry*, 6th Edition (Berlin, Springer-Verlag).
- Peterson BJ, Fry B (1987) Stable isotopes in ecosystem studies. *Annual Reviews of Ecological Systems*, 18:293–320.
- Richards MP, Hedges REM (1999) Stable isotope evidence for similarities in the types of marine foods used by Late Mesolithic humans at sites along the Atlantic coast of Europe. *J Archaeol Sci* 26:717–722.
- Kelly JF (2000) Stable isotopes of carbon and nitrogen in the study of avian and mammalian trophic ecology. *Canadian J Zool* 78:1–27.
- Fry B (1991) Stable isotope diagrams of freshwater food webs. *Ecology* 72:2293–2297.
- O'Leary M (1981) Carbon isotopic fractionation in plants. *Phytochemistry* 20:553–567.
- Hedges REM, Reynard L (2007) Nitrogen isotopes and the trophic level of humans in archaeology. *J Archaeol Sci* 34:1240–1251.
- Schoeninger M, DeNiro M (1984) Nitrogen and carbon isotopic composition of bone collagen from marine and terrestrial animals. *Geochim Cosmochim Acta* 48:625–639.
- van Klinken GJ, van der Plicht H, Hedges R (1994) Bone 13C/12C ratios reflect (palaeo-) climatic variations. *Geophys Res Lett* 21:445–448.
- Heaton THE, Vogel JC, von la Chevallerie G, Collett G (1986) Climatic influence on the isotopic composition of bone nitrogen. *Nature* 322:822–823.
- Hedges REM, Stevens RE, Richards MP (2004) Bone as a stable isotope archive for local climatic information. *Quaternary Sci Rev* 23:959–965.
- Richards MP, Hedges REM (2003) Bone collagen  $\delta^{13}C$  and  $\delta^{15}N$  values of fauna from Northwest Europe reflect palaeoclimatic variation over the last 40,000 years. *Palaeogeogr Palaeoclimatol* 193:261–267.
- Stevens RE, Hedges REM (2004) Carbon and nitrogen stable isotope analysis of north-west European horse bone and tooth collagen, 40,000 BP–present: Palaeoclimatic interpretations. *Quaternary Sci Rev* 23:977–991.
- Herring DA, Saunders SR, Katzenberg MA (1998) Investigating the weaning process in past populations. *Am J Phys Anthropol* 105:425–439.
- Schurr MR (1998) Using stable nitrogen isotopes to study weaning behaviour in past populations. *World Archaeol* 30:327–342.
- Richards MP, Mays SA, Fuller BT (2002) Stable carbon and nitrogen isotope values of bone and teeth reflect weaning at the Mediaeval Wharram Percy Site, Yorkshire, UK. *Am J Phys Anthropol* 199:205–210.
- Ambrose SH (1990) Preparation and characterization of bone and tooth collagen for stable carbon and nitrogen isotope analysis. *J Archaeol Sci* 17:431–451.
- Krause J, et al. (2007) Neanderthals in Central Asia and Siberia. *Nature* 449:902–904.
- Trinkaus E, et al. (2003) An early modern human from the Peștera cu Oase, Romania. *Proc Nat Acad Sci USA* 100:11231–11236.
- Richards MP, et al. (2008) Isotopic evidence for omnivory among European cave bears: Late Pleistocene *Ursus spelaeus* from the Peștera cu Oase, Romania. *Proc Nat Acad Sci USA* 105:600–604.
- Orschiedt J (2002) Datation d'un vestige humain provenant de la Rochette (Saint-Leon-Sur-Vezere) par la methode du carbone 14 en spectrometrie de masse. *Paléo* 14:15.
- Richards MP, Fuller BT, Molleson TI (2006) Stable isotope palaeodiet study of humans and fauna from the multi-period (Iron Age, Viking and Late Medieval) site of Newark Bay, Orkney. *J Archaeol Sci* 33:122–131.
- Drucker D, Bocherens H (2004) Carbon and nitrogen stable isotopes as tracers of change in diet breadth during Middle and Upper Palaeolithic in Europe. *Int J Osteoarchaeol* 14:162–177.
- Richards, M.P. (2000). Human and faunal stable isotope analyses from Goat's Hole and Foxhole Caves, Gower. In Aldhouse-Green, S, ed., *Paviland Cave and the 'Red Lady': A Definitive Report*. (University of Wales College, Newport, and National Museums and Galleries of Wales, Newport, U.K.) pp. 71–75.
- Jacobi RM, Higham TFG (2008) The 'Red Lady' ages gracefully: New ultrafiltration AMS determinations from Paviland. *J Hum Evol* 55:898–907.
- Mithen S, Finlay N, Carruthers W, Carter S, Ashmore P (2001) Plant use in the Mesolithic: Evidence from Staosnaig, Isle of Colonsay, Scotland. *J Archaeol Sci* 28:223–234.

50. Stringer CB, et al. (2008) Neanderthal exploitation of marine mammals in Gibraltar. *Proc Natl Acad Sci USA* 105:14319–14324.
51. Stiner MC (1994) *Honor Among Thieves: A Zooarchaeological Study of Neandertal Ecology* (Princeton Univ. Press, Princeton).
52. Pathou-Mathis M, (2000) Neanderthal subsistence behaviour in Europe. *Int J Osteoarchaeol* 10:379–395.
53. Gaudzinski S, Roebroeks W (2000) Adults only: Reindeer hunting at the Middle Palaeolithic site Salzgitter Lebenstedt, Northern Germany. *J Hum Evol* 38:497–521.
54. Hardy BL, Kay M, Marks AE, Monigal K (2001) Stone tool function at the Paleolithic sites of Starosele and Buran Kaya III, Crimea: Behavioral implications. *Proc Nat Acad Sci USA* 98:10972–10977.
55. Grayson DK, Delpech F (2003) Ungulates and the Middle-to-Upper Paleolithic transition at Grotte XVI (Dordogne, France). *J Archaeol Sci* 30:1633–1648.
56. Grayson DK, Delpech F (2008) The large mammals of Roc de Combe (Lot, France): The Châtelperronian and Aurignacian assemblages. *J Anthropol Archaeol* 27:338–362.
57. Niven L (2006) *The Palaeolithic Occupation of Vogelherd Cave: Implications for the Subsistence Behavior of Late Neanderthals and Early Modern Humans* (Tübingen: Kerns Verlag).
58. Klein RG, et al. (2004) The Ysterfontein 1 Middle Stone Age site, South Africa, and early human exploitation of coastal resources *Proc Nat Acad Sci USA* 101:5708–5715.
59. Yellen JE, Brooks AS, Cornelissen E, Mehlman MJ, Stewart K (1995) A Middle Stone Age worked bone industry from Katanda, Upper Semliki Valley, Zaire. *Science* 268:553–556.
60. Richards MP, Schulting RJ, Hedges REM (2003). Archaeology: Sharp shift in diet at onset of Neolithic. *Nature* 425:366.
61. Tauber H (1981)  $^{13}\text{C}$  evidence for dietary habits of prehistoric man in Denmark. *Nature* 292:332–333.
62. Trinkaus E (2007) European early modern humans and the fate of the Neandertals. *Proc Natl Acad Sci USA* 104:7367–7372.
63. Smith C, et al. (2009) A three-phase liquid chromatographic method for  $\delta^{13}\text{C}$  analysis of amino acids from biological protein hydrolysates using liquid chromatography-isotope ratio mass spectrometry. *Anal Biochem* 390:165–172.
64. Richards MP, Fuller BF, Hedges REM (2001) Sulphur isotopic variation in ancient bone collagen from Europe: implications for human palaeodiet, residence mobility, and modern pollutant studies. *Earth Planet Sci Lett* 191:185–190.
65. Richards MP, Fuller BT, Sponheimer M, Robinson T, Ayliffe L (2003) Sulphur isotope measurements in archaeological samples: some methodological considerations. *Int J Osteoarchaeol* 13:37–45.
66. Nehlich O, Richards MP (2009) Establishing collagen quality criteria for sulphur isotope analysis of archaeological bone collagen. *Archaeol Anthropol Sci* 1:59–75.