

Killing niche competitors by remote-control bacteriophage induction

Laura Selva^a, David Viana^a, Gili Regev-Yochay^b, Krzysztof Trzcinski^b, Juan Manuel Corpa^a, Íñigo Lasa^c, Richard P. Novick^d, and José R. Penadés^{a,e,1}

^aUniversidad Cardenal Herrera-CEU, 46113 Moncada, Valencia, Spain; ^bDepartments of Epidemiology and Immunology and Infectious Diseases, Harvard School of Public Health, Boston; ^cInstituto de Agrobiotecnología, CSIC-Universidad Pública de Navarra, 31006 Pamplona, Navarra, Spain; ^dSkirball Institute, New York University Medical Center, 540 First Avenue, New York, NY 10016; and ^eCentro de Investigación y Tecnología Animal, Instituto Valenciano de Investigaciones Agrarias (CITA-IVIA), Apdo. 187, 12.400 Segorbe, Castellón, Spain

Edited by Sankar Adhya, National Institutes of Health, Bethesda, MD, and approved December 3, 2008 (received for review September 26, 2008)

A surprising example of interspecies competition is the production by certain bacteria of hydrogen peroxide at concentrations that are lethal for others. A case in point is the displacement of *Staphylococcus aureus* by *Streptococcus pneumoniae* in the nasopharynx, which is of considerable clinical significance. How it is accomplished, however, has been a great mystery, because H₂O₂ is a very well known disinfectant whose lethality is largely due to the production of hyperoxides through the abiological Fenton reaction. In this report, we have solved the mystery by showing that H₂O₂ at the concentrations typically produced by pneumococci kills lysogenic but not nonlysogenic staphylococci by inducing the SOS response. The SOS response, a stress response to DNA damage, not only invokes DNA repair mechanisms but also induces resident prophages, and the resulting lysis is responsible for H₂O₂ lethality. Because the vast majority of *S. aureus* strains are lysogenic, the production of H₂O₂ is a very widely effective antistaphylococcal strategy. Pneumococci, however, which are also commonly lysogenic and undergo SOS induction in response to DNA-damaging agents such as mitomycin C, are not SOS-induced on exposure to H₂O₂. This is apparently because they are resistant to the DNA-damaging effects of the Fenton reaction. The production of an SOS-inducing signal to activate prophages in neighboring organisms is thus a rather unique competitive strategy, which we suggest may be in widespread use for bacterial interference. However, this strategy has as a by-product the release of active phage, which can potentially spread mobile genetic elements carrying virulence genes.

hydrogen peroxide | SOS response | *Staphylococcus aureus* | *Streptococcus pneumoniae* | bacterial interference

The interactions among bacteria living communally are highly complex and extremely interesting—illuminating, as they do, a long-ignored but nevertheless critical aspect of microbial biology. One can readily envision interactions such as direct competition for scarce nutrients, mutual cooperation for the conversion of substrates to utilizable metabolites, “borrowing” of quorum-sensing signals, DNA transfer, biofilm formation and maintenance, and interference or inhibition mediated by antibacterial products, including bacteriocins, antibiotics, and low-molecular-weight toxic compounds such as H₂O₂ (1). In this article, we consider a specific case of H₂O₂-mediated bacterial interference, that between pneumococci and *Staphylococcus aureus*, which, although well documented, occurs by an entirely unknown mechanism.

Several epidemiological studies have shown a negative association between carriage of *Streptococcus pneumoniae* and *S. aureus* (2, 3), raising public health concern that mass pneumococcal vaccination may cause an increase in *S. aureus* colonization and infection. As a case in point, it has been reported that children with recurrent otitis media vaccinated with the heptavalent pneumococcal vaccine had increased incidence of *S. aureus*-related acute otitis media and *S. aureus* colonization (3).

Recent in vitro and in vivo studies have demonstrated that the interference between these 2 pathogens is related to hydrogen peroxide production by *S. pneumoniae*, which is bactericidal to *S. aureus* (4, 5). Similar observations have been reported for certain other pairs of bacteria (6). It is highly intriguing how the relatively low levels of hydrogen peroxide produced safely by some bacteria are bactericidal to others, despite the relative abundance of mechanisms protecting bacterial cells from oxidative damage, such as H₂O₂-inactivating enzymes and antioxidants (7) or DNA lesion repair systems (8).

Here, we shed light on the mechanism of interference between H₂O₂-producing bacteria and *S. aureus*. We present data supporting the idea that prophages may have a much greater role in bacterial ecology than has hitherto been suspected—namely, that killing of a target organism by “remote control” prophage induction may represent a major modality of directional bacterial interference. We show also that lysogenic staphylococci are much more sensitive to DNA-damaging antibiotics, such as fluoroquinolones, than nonlysogens, almost certainly for the same reason. Given the high prevalence of lysogeny, we can now predict that small, SOS-inducing molecules, produced in the environment at subinhibitory concentrations, may have strong selective value as effectors of directional interference.

Results

Hydrogen Peroxide Kills Only Lysogenic *S. aureus*. Several species of bacteria have H₂O₂-dependent bactericidal activity toward *S. aureus* (4, 9). However, the mechanism by which the relatively low levels of H₂O₂ produced by these organisms are bactericidal to *S. aureus* remains to be determined. One possibility is that H₂O₂ produced by one organism induces the SOS response in a competing (target) organism, lethally activating resident prophages in the latter. If so, staphylococcal lysogens but not nonlysogens should be sensitive to H₂O₂ and pneumococci should be insensitive, even though they are often or always lysogenic. Accordingly, we tested 8 strains of *S. aureus*, 6 lysogenic, 2 nonlysogenic (RN450 and V329), including a congenic pair in which one (RN10359) was an 80α lysogen of the other (RN450) and a strain producing the phage-carried PVL toxin (strain LUG855), lately implicated in serious staphylococcal infections [10; supporting information (SI) Table S1]. We used H₂O₂ at 0.5 mM, in the range ordinarily seen with pneumococcal cultures, and observed that all of the lysogenic strains were highly

Author contributions: J.R.P. designed research; L.S. and D.V. performed research; G.R.-Y., K.T., J.M.C., and I.L. contributed new reagents/analytic tools; L.S., D.V., I.L., R.P.N., and J.R.P. analyzed data; and G.R.-Y., K.T., R.P.N., and J.R.P. wrote the paper;

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. E-mail: jpenades@ivia.es.

This article contains supporting information online at www.pnas.org/cgi/content/full/0809600106/DCSupplemental.

© 2009 by The National Academy of Sciences of the USA




Fig. 1. Killing of *S. aureus* by H₂O₂. Survival of lysogenic (COL, Mu50, FRI-S6, FR122, LUG855, and RN10359) or nonlysogenic (V329, RN450) *S. aureus* strains in media supplemented with hydrogen peroxide at a concentration of 0.5 mM, and unsupplemented medium (control). Values represent the average of 3 independent experiments. Variation was within $\pm 5\%$ in all cases.

sensitive, whereas the nonlysogens were insensitive (see Fig. 1), consistent with the above prediction.

H₂O₂ Induces SOS in *S. aureus*. It is well known that H₂O₂ induces the SOS response in *Escherichia coli* and other bacteria (8), and it is strongly predicted that it would do so in *S. aureus*, and that this would be responsible for H₂O₂ killing of lysogens, although no clear test of this has been published. Accordingly, we tested the same lysogenic strains as in Fig. 1, for H₂O₂ induction of prophages. Because *recA*-dependent prophage induction is a highly typical feature of the SOS response, we included RN1030, a lysogenic *recA*-defective derivative of RN451. Because cleav-

age of the prophage repressor is also necessary for prophage induction, we also included JP3592, an 80 α lysogen with a noncleavable mutant phage repressor. As shown in Tables 1 and S2, H₂O₂ induced the prophages of all strains tested, in a dose-dependent manner, with the exception of RN1030, and JP3592. However, the response to H₂O₂ was quite variable, perhaps owing to differences in the response to oxidative stress. Phage induction by H₂O₂ was, as one would expect, accompanied by a corresponding decrease in viability, as also shown in Tables 1 and S2. No reduction in viability was seen with the nonlysogen or with the phage repressor mutant, except for a slight reduction in viability owing to direct H₂O₂ toxicity at the highest H₂O₂ concentration used.

As would be expected for the producer of an antibacterial substance, *S. pneumoniae* is entirely resistant to H₂O₂ at the concentrations present in its cultures, and it has been shown that this is because it is insensitive to the toxic products of the Fenton reaction (11). Because these products cause DNA damage, which is responsible for SOS induction, we hypothesized that H₂O₂ would not induce the SOS response in *S. pneumoniae* and therefore that even lysogenic *S. pneumoniae* would be able to use H₂O₂ to interfere with competing species. To test this hypothesis, we compared lysogenic and nonlysogenic *S. pneumoniae* for sensitivity to H₂O₂ and mitomycin C (MC). As shown in Fig. 2, neither was sensitive to H₂O₂; however, the lysogens (strains 623 and 949), but not the nonlysogen (TIGR-4) were sensitive to SOS induction by MC, leading to lysis and, presumably, to the release of active phage, although we have not confirmed this in the present study. Nevertheless, it is clear that neither pneumococcal lysogens nor nonlysogens are sensitive to H₂O₂ at concentrations normally present in pneumococcal cultures and that H₂O₂ does not induce the SOS response in pneumococci, which accounts for the ability of this organism to produce H₂O₂ for use as a weapon of mass destruction with impunity.

Table 1. Phage titer and survival of H₂O₂-induced lysogenic staphylococcal strains

Donor strain	[H ₂ O ₂], mM	CFU*	Phage titer†
RN450	NI	8.4×10^8	–
	0.05	7.1×10^8	–
	0.1	2.9×10^8	–
	0.5	1.8×10^8	–
	1	6.3×10^7	–
RN10359 (RN450 lysogenic for 80 α)	NI	5.9×10^8	8.8×10^5
	0.05	1.2×10^9	1.9×10^6
	0.1	2.9×10^7	9.0×10^7
	0.5	1.8×10^4	5.3×10^9
	1	5.6×10^3	2.0×10^8
JP3592 (RN10359 cl G130E)	NI	6.7×10^8	<10
	0.05	1.8×10^8	<10
	0.1	1.9×10^8	<10
	0.5	1.8×10^8	<10
	1	0.8×10^8	<10
RN451 (RN450 lysogenic for ϕ 11)	NI	5.9×10^8	2.8×10^4
	0.05	4.2×10^8	4.9×10^4
	0.1	4.9×10^7	3.1×10^5
	0.5	3.8×10^5	2.8×10^6
	1	5.6×10^4	8.9×10^5
RN1030 (RN451 <i>recA</i> -mutant)	NI	ND	<10
	0.05	ND	<10
	0.1	ND	<10
	0.5	ND	<10
	1	ND	<10

The means of results from 3 independent experiments are presented. Variation was within $\pm 5\%$ in all cases. NI, not induced; ND, not determined.

*Number of cells recovered after 4 h of incubation.

†Number of plaque-forming phages per milliliter of induced culture, using RN4220 as indicator strain.




Fig. 2. Lysis of *S. pneumoniae* induced by mitomycin C or H₂O₂. Cultures of lysogenic (623 and 949) or nonlysogenic (TIGR4) strains received mitomycin C (0.1 µg/mL) or H₂O₂ (0.5 mM) at time 0, and the OD of cultures was monitored at 600 nm.

Effect of H₂O₂ Production by *S. pneumoniae*. To confirm that H₂O₂ is responsible for the killing of *S. aureus* by *S. pneumoniae*, we cultured a lysogenic, H₂O₂-sensitive, *S. aureus* strain (RN10395) with 2 *S. pneumoniae* strains, TIGR4 and Pn-20. It has reported that these strains naturally produce H₂O₂ (4), with full-grown cultures ($\approx 10^7$ cfu/mL) containing ≈ 1 mM hydrogen peroxide (4). As shown in Table 2, H₂O₂-producing *S. pneumoniae* had the expected effects in coculture with the 80 α WT lysogen—phage induction and loss of viability. We also tested mutants of TIGR4 and Pn-20 with deletions of the pyruvate oxidase gene, *spxB*, required for H₂O₂ production (4). Neither of the Δ *spxB* mutants had any effect on viability or prophage induction with lysogenic *S. aureus*. Further evidence was provided by a test with the phage 80 α mutant defective in the ability to cleave its prophage repressor, and therefore, not SOS inducible. This mutant was as insensitive to H₂O₂ as the nonlysogens tested.

Role of Catalase in Hydrogen Peroxide SOS Induction. Because *S. aureus* produces a potent catalase as a defense against H₂O₂ toxicity, it seemed paradoxical that it would nevertheless be susceptible to SOS induction of prophages by such low concentrations of H₂O₂. To evaluate the role of catalase in SOS induction by H₂O₂, we constructed *katA* mutants of RN450 (JP3852) and RN10359 (JP3853) by transducing the *katA* mu-

tation from strain KS100 (7). The 2 *katA* mutants and their *katA*+ parents were incubated for 6 h in TSB broth with various concentrations of H₂O₂. As shown in Fig. 3, KatA strongly protected the nonlysogenic RN450 at the higher H₂O₂ concentrations, but had a much less dramatic effect on the congenic lysogen, RN10359, both with respect to viability and to phage production, suggesting that KatA is much more strongly protective of general viability in the presence of H₂O₂ than of susceptibility to SOS induction. This may be because peroxide lethality involves a wide range of oxidative damages, whereas SOS induction involves only the RecA-LexA pathway, which is induced by only minimal DNA damage, much less than that necessary to kill the cell. Perhaps catalase does not act rapidly enough or efficiently enough at these lower concentrations to prevent minimal DNA damage. This suggests that general toxicity and SOS induction by H₂O₂ may target different processes, and that KatA can protect a nonlysogen but not a lysogen against H₂O₂ killing at relevant concentrations. Additionally, the lower phage production by the *kat*-mutant at high concentrations of H₂O₂ is probably the result of decreased capacity of the cells to support phage growth.

Hydrogen Peroxide Induces Phage-Mediated SaPI Transfer. A by-product of H₂O₂ killing of lysogens by SOS induction is, of

Table 2. Effects of coculture of *S. aureus* with H₂O₂-producing *S. pneumoniae*

Donor strain	Inducer*	CFU [†]	Phage titer [‡]
RN10359 (RN450 lysogenic for 80 α)	H ₂ O ₂	7.0×10^5	5.0×10^9
	Pn-20	4.6×10^5	8.7×10^7
	Pn-20 Δ <i>spxB</i>	1.4×10^9	3.3×10^5
	TIGR4	4.2×10^6	5.9×10^7
	TIGR4 Δ <i>spxB</i>	1.2×10^9	5.3×10^5
RN450	NI	1.9×10^9	4.6×10^5
	H ₂ O ₂	7.3×10^8	–
	Pn-20	7.8×10^8	–
	Pn-20 Δ <i>spxB</i>	1.1×10^9	–
	TIGR4	1.2×10^9	–
	TIGR4 Δ <i>spxB</i>	1.0×10^9	–
JP3592 (RN10359 cl G130E)	NI	1.1×10^9	–
	H ₂ O ₂	5.3×10^8	<10
	Pn-20	8.8×10^8	<10
	Pn-20 Δ <i>spxB</i>	2.1×10^8	<10
	TIGR4	1.2×10^8	<10
TIGR4 Δ <i>spxB</i>	9.7×10^8	<10	
NI	1.0×10^9	<10	

The means of results from 3 independent experiments are presented. Variation was within $\pm 5\%$ in all cases.

*H₂O₂ (0.5 mM). NI, not induced.

[†]Number of cells recovered after 4 h of incubation.

[‡]Number of plaque-forming phages per milliliter of induced culture, using RN4220 as indicator strain.




Fig. 3. Role of catalase in SOS induction by H₂O₂. (A) Survival of *S. aureus* strains in media supplemented with hydrogen peroxide at different concentrations. Values represent the average of 3 independent experiments. Variation was within $\pm 5\%$ in all cases. (B) Phage titer obtained from the lysogenic strains analyzed in A.

course, the release of viable phage into the environment. This could promote the spread of phage-encoded toxins, such as PVL, and of phage-related pathogenicity islands such as the SaPIs, which encode TSST-1 and other superantigen toxins (12). We have presented data on H₂O₂ induction of a PVL-encoding phage (Table S2); induction of SaPI particle production by coculture with *S. pneumoniae* is presented in Table S3.

Discussion

In this report, we have examined the widespread strategy of bacterial interference by the production of H₂O₂, demonstrating, in the case of *S. pneumoniae* vs. *S. aureus* that it acts through the lethal induction of resident prophages in the target organism. We note that our results play deeply into the overall universe of bacteria and phages. Lysogeny is extremely common among bacteria, representing a major symbiotic strategy in which the prophage, residing stably in the host's chromosome, provides protection against attack by other phages, not only through coimmunity, but also by encoding restriction-modification capabilities or other more general protective mechanisms. And prophages may also enhance the virulence of pathogens. A consequence of lysogeny is, of course, inherent in the definition, and spontaneous lysis, a universal feature of lysogens, probably has an important role in phage-mediated lateral gene transfer. Nevertheless, lysogeny is, after all, an Achilles heel for the organism as the prophage state causes vulnerability to environmental influences such as radiation and certain chemicals that cause (sublethal) DNA damage, invoking the SOS response, a stress response whose objective is to repair damaged DNA. Prophage repressors are generally inactivated during the SOS response, setting in motion the phage lytic cycle; although the biological purpose of this is not entirely obvious, it would provide a wonderful target for aggression by competing organisms, if only they knew how to cause DNA damage and thus to induce prophages. As we have shown here, the production of H₂O₂ at sublethal concentrations serves this purpose admirably. Biochemically, H₂O₂ is easily produced by the oxidation of pyruvate, generates DNA-damaging hyperoxides through the Fenton reaction (13), and thereby induces the SOS response in many bacteria. As is the case with all antibacterial substances, the producing organism is immune. With antibiotics, the immune mechanism is usually obvious. With H₂O₂, the mechanism of immunity is not so obvious because it involves resistance to the toxic products of the Fenton reaction (11), including, as we show here, resistance to induction of the SOS response. Pneumococci do not produce catalase, which inactivates H₂O₂ and could compromise the use of H₂O₂ as a weapon; perhaps this is precisely why the pneumococci lack catalase. And we would predict that other H₂O₂-producing organisms also lack catalase

for the same reason. Given the occurrence of H₂O₂-mediated bacterial interference based on lethal prophage induction in the target organism, we note that DNA-damaging antibiotics have recently been shown to induce prophages and thereby to promote the spread of phage-coded toxins and of phage-related pathogenicity islands (14–16). One would therefore predict that lysogenic bacteria would be more sensitive to the bactericidal




Fig. 4. Killing of lysogenic *S. aureus* cells by antibiotics. (A) Survival *S. aureus* strains in media supplemented with different concentrations of the antibiotic ciprofloxacin. RN450, nonlysogenic; RN10359, RN450 lysogenic for phage 80 α ; RN10359 cI⁻, derivative of RN10359 carrying a non-SOS-inducible phage 80 α ; RN451, RN450 lysogenic for phage 11; RN451 cI⁻, derivative of RN451 carrying a non-SOS-inducible phage 11. Values represent the average of 3 independent experiments. Variation was within $\pm 5\%$ in all cases. (B) Antibiogram showing the different susceptibility between strains RN10359 (lysogenic for phage 80 α) and RN10359 cI⁻ (carrying a non-SOS-inducible phage 80 α) to enrofloxacin discs. Average zone of inhibition: RN10359, 36 mm; RN10359 cI⁻, 30 mm.

effects of DNA-damaging antibiotics than nonlysogens, owing to lysis caused by SOS-induced prophages in the former, as we have demonstrated above for H₂O₂. As shown in Fig. 4, two different lysogenic derivatives of *S. aureus* strain RN450, RN451 and RN10359, lysogenic for ϕ 11 and 80 α , respectively, were considerably more sensitive to ciprofloxacin (Fig. 4A) than their nonlysogenic parent and this sensitivity was eliminated in both cases by the phage repressor cleavage defect. Similarly, the noncleavable repressor mutation significantly reduced sensitivity to enrofloxacin (Fig. 4B). In a more general sense, we would predict that lysogenic bacteria would be more sensitive than nonlysogens to interference by antibiotic producers in the environment, where the antibiotic concentrations would surely be very low. This remains to be tested.

Materials and Methods

Bacterial Strains and Growth Conditions. Bacterial strains used in these studies are listed in Table S1. Bacteria were grown at 37 °C overnight on TSA agar medium (*S. aureus*) or BHI with 1.5% agar (*S. pneumoniae*), supplemented with antibiotics as appropriate. Broth cultures were grown at 37 °C in TSB (*S. aureus*) or Todd Hewitt (*S. pneumoniae*).

Induction of Prophages. Procedures for preparation and analysis of phage lysates, transduction, and transformation in *S. aureus* were performed essentially as described in refs. 17 and 18. In general, bacteria were grown in TSB to

OD₅₄₀ = 0.15 and induced by the addition of H₂O₂ at various concentrations and cultures were continued at 32 °C with slow shaking (80 rpm). Presence of the lysis was evaluated within 3 h by using previously applied criteria (17).

Coculture Assay. Interference between *S. aureus* and *S. pneumoniae* strains was measured in coculture assay as described in ref. 4. In brief, 1 mL of the *S. pneumoniae* cultures (OD₅₄₀ 0.4–0.5) were mixed with 1 mL of the *S. aureus* cultures (OD₅₄₀ 0.3–0.4), and incubated 4 h at 37 °C. To quantify bactericidal activity, the cocultured bacterial strains were plated on selective medium (TSA supplemented with 5 μ g/mL optochin), where only *S. aureus* can grow.

DNA Methods. General DNA manipulations were performed by standard procedures (19, 20). Strain JP3592 was obtained by using plasmid pMAD, as described in ref. 15. The oligonucleotides pairs used were phi11–1cB (5′-cgcgatccAGTGT-TAATGTGTATATGCTC-3′)/phi11–3c (5′-TAATTCCTCCTATCTCAGCACCAGTT-GCACC-3′) and phi80alpha-cl-9mE (5′-CCGGAATTCACAGATTCGTTTTATTTCCC-3′)/phi11–4m (5′-GGTGCAACTGGTCTGAGATAGGAGAAGAATTA-3′).

ACKNOWLEDGMENTS. We thank Simon Foster, Gabriela Bowden, and Pedro García for providing bacterial strains used in this study and Jeff Weiser, Simon Foster, Adela González de la Campa, and Alex Mira for helpful comments on the manuscript. This work was supported by Comisión Interministerial de Ciencia y Tecnología Grants BIO2005-08399-C02-02, BIO2008-05284-C02-02, and BIO2008-00642-E/C; Cardenal Herrera-CEU University Grants PRCEU-UCH25/08 and Copernicus program; and by Conselleria de Agricultura, Pesca i Alimentació (CAPIA), and from the Generalitat Valenciana (ACOMP07/258) (J.R.P.). L.S. and D.V. were supported by Cardenal Herrera-CEU University fellowships.

- Bassler BL, Losick R (2006) Bacterially speaking. *Cell* 125:237–246.
- Regev-Yochay G, et al. (2004) Association between carriage of *Streptococcus pneumoniae* and *Staphylococcus aureus* in Children. *JAMA* 292:716–720.
- Bogaert D, et al. (2004) Colonisation by *Streptococcus pneumoniae* and *Staphylococcus aureus* in healthy children. *Lancet* 363:1871–1872.
- Regev-Yochay G, Trzcinski K, Thompson CM, Malley R, Lipsitch M (2006) Interference between *Streptococcus pneumoniae* and *Staphylococcus aureus*: In vitro hydrogen peroxide-mediated killing by *Streptococcus pneumoniae*. *J Bacteriol* 188:4996–5001.
- Park B, Nizet V, Liu GY (2008) Role of *Staphylococcus aureus* catalase in niche competition against *Streptococcus pneumoniae*. *J Bacteriol* 190:2275–2278.
- Pericone CD, Overweg K, Hermans PW, Weiser JN (2000) Inhibitory and bactericidal effects of hydrogen peroxide production by *Streptococcus pneumoniae* on other inhabitants of the upper respiratory tract. *Infect Immun* 68:3990–3997.
- Cosgrove K, et al. (2007) Catalase (KatA) and alkyl hydroperoxide reductase (AhpC) have compensatory roles in peroxide stress resistance and are required for survival, persistence, and nasal colonization in *Staphylococcus aureus*. *J Bacteriol* 189:1025–1035.
- Goerlich O, Quillardet P, Hofnung M (1989) Induction of the SOS response by hydrogen peroxide in various *Escherichia coli* mutants with altered protection against oxidative DNA damage. *J Bacteriol* 171:6141–6147.
- Strauss E (2000) Microbiology. Fighting bacterial fire with bacterial fire. *Science* 290:2231–2233.
- Labandeira-Rey M, et al. (2007) *Staphylococcus aureus* Panton-Valentine leukocidin causes necrotizing pneumonia. *Science* 315:1130–1133.
- Pericone CD, Park S, Imlay JA, Weiser JN (2003) Factors contributing to hydrogen peroxide resistance in *Streptococcus pneumoniae* include pyruvate oxidase (SpxB) and avoidance of the toxic effects of the fenton reaction. *J Bacteriol* 185:6815–6825.
- Novick RP, Subedi A (2007) The SaPIs: Mobile pathogenicity islands of *Staphylococcus*. *Chem Immunol Allergy* 93:42–57.
- Imlay JA, Chin SM, Linn S (1988) Toxic DNA damage by hydrogen peroxide through the Fenton reaction in vivo and in vitro. *Science* 240:640–642.
- Zhang X, et al. (2000) Quinolone antibiotics induce Shiga toxin-encoding bacteriophages, toxin production, and death in mice. *J Infect Dis* 181:664–670.
- Maiques E, et al. (2006) (beta)-Lactam antibiotics induce the SOS response and horizontal transfer of virulence factors in *Staphylococcus aureus*. *J Bacteriol* 188:2726–2729.
- Ubeda C, et al. (2005) Antibiotic-induced SOS response promotes horizontal dissemination of pathogenicity island-encoded virulence factors in staphylococci. *Mol Microbiol* 56:836–844.
- Novick RP (1991) Genetic systems in staphylococci. *Methods Enzymol* 204:587–636.
- Lindsay JA, Ruzin A, Ross HF, Kurepina N, Novick RP (1998) The gene for toxic shock toxin is carried by a family of mobile pathogenicity islands in *Staphylococcus aureus*. *Mol Microbiol* 29:527–543.
- Ausubel FM, et al. (1990) *Current Protocols in Molecular Biology* (Wiley, New York).
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab Press, Cold Spring Harbor, NY).