

# Fungal infection counters insecticide resistance in African malaria mosquitoes

Marit Farenhorst<sup>a</sup>, Joel C. Mouatcho<sup>b,c</sup>, Christophe K. Kikankie<sup>b,c</sup>, Basil D. Brooke<sup>b,d</sup>, Richard H. Hunt<sup>c</sup>, Matthew B. Thomas<sup>e</sup>, Lizette L. Koekemoer<sup>b,d</sup>, Bart G. J. Knols<sup>a,f,1</sup>, and Maureen Coetzee<sup>b,d</sup>

<sup>a</sup>Laboratory of Entomology, Wageningen University and Research Centre, P.O. Box 8031, 6700 EH Wageningen, The Netherlands; <sup>b</sup>Vector Control Reference Unit, National Institute for Communicable Diseases, Private Bag X4, Sandringham 2131, Johannesburg, South Africa; <sup>c</sup>School of Animal, Plant, and Environmental Sciences, University of the Witwatersrand, Private Bag 3, Wits 2050, Johannesburg, South Africa; <sup>d</sup>Division of Virology and Communicable Disease Surveillance, School of Pathology, National Health Laboratory Service and University of the Witwatersrand, Private Bag X4, Sandringham 2131, Johannesburg, South Africa; <sup>e</sup>Center for Infectious Disease Dynamics, Department of Entomology, Pennsylvania State University, 208 Mueller Lab, University Park, PA 16802; and <sup>f</sup>Division of Infectious Diseases, Tropical Medicine and Aids, Academic Medical Center, F4-217 Meibergdreef 9, 1105 AZ, Amsterdam, The Netherlands

Communicated by Hans R. Herren, Millennium Institute, Arlington, VA, July 30, 2009 (received for review December 18, 2008)

**The evolution of insecticide resistance in mosquitoes is threatening the effectiveness and sustainability of malaria control programs in various parts of the world. Through their unique mode of action, entomopathogenic fungi provide promising alternatives to chemical control. However, potential interactions between fungal infection and insecticide resistance, such as cross-resistance, have not been investigated. We show that insecticide-resistant *Anopheles* mosquitoes remain susceptible to infection with the fungus *Beauveria bassiana*. Four different mosquito strains with high resistance levels against pyrethroids, organochlorines, or carbamates were equally susceptible to *B. bassiana* infection as their baseline counterparts, showing significantly reduced mosquito survival. Moreover, fungal infection reduced the expression of resistance to the key public health insecticides permethrin and dichlorodiphenyltrichloroethane. Mosquitoes preinfected with *B. bassiana* or *Metarhizium anisopliae* showed a significant increase in mortality after insecticide exposure compared with uninfected control mosquitoes. Our results show a high potential utility of fungal biopesticides for complementing existing vector control measures and provide products for use in resistance management strategies.**

biopesticide | DDT | pyrethroids | resistance management | vector control

Increasing incidences of insecticide resistance are reported in the major African malaria vector species *Anopheles gambiae* s.s. (1–3), *Anopheles funestus* (4, 5), and *Anopheles arabiensis* (6, 7), thereby threatening the efficiency of insecticides approved for malaria vector control. As the use of indoor residual spraying and insecticide-treated nets is scaling up (ref. 8 and <http://www.rollbackmalaria.org/gmap/>), so will the selection pressure for insecticide resistance. Resistance management strategies, such as the use of chemical mixtures or rotations, are suggested to improve the sustainability of these approaches (9, 10), but with significant cross-resistance between the currently approved classes of insecticide (9–11) and no new public health insecticides introduced in the last 20 years (12), practical options are few. For sustainable malaria vector control, the requirement for new products with unique modes of action is becoming increasingly evident.

Previous work has suggested that entomopathogenic fungi could play this role (13–15). A range of isolates belonging to the fungal species *Metarhizium anisopliae* and *Beauveria bassiana* have been shown to infect and significantly reduce the longevity of adult *Anopheles* mosquitoes, killing them within 14 days (13, 14, 16). This slow speed of kill is considered to dramatically reduce the selection pressure for fungal resistance development (15, 17); yet, because *Plasmodium* maturation within the mosquito also takes  $\approx 14$  days (18) delayed kill is considered to be sufficient to significantly reduce the mosquito's vectorial capacity. Prelethal effects of fungal infection, including reductions in *Plasmodium* sporozoite formation (13), feeding propensity (13,

19), and fecundity (19), increase the potential impact on malaria transmission even further.

Under appropriate conditions (humidity and pH) and adequate nutrient availability, fungal spores can infect insects via attachment to the epicuticle and subsequent germination (15, 20). Spores penetrate the cuticle via formation of germ tubes and cuticle-degrading enzymes (21) and grow in the host hemocoel where they use nutrients, destroy host cells, and eventually kill the insect (20). For fungus-based biopesticides to play a prominent role in malaria control, an important criterion is that fungal susceptibility will remain unaffected by resistance to insecticides. The mode of action of entomopathogenic fungi (i.e., via external contact and proliferation through the hemocoel) makes direct cross-reactions with insecticides unlikely. However, the indirect effects of insecticide resistance mechanisms have been shown to reduce pathogen proliferation. Enhanced concentrations of esterases in organophosphate-resistant *Culex* mosquitoes have been implicated in limiting growth of filarial worms (22, 23), and enhanced monooxygenase levels in pyrethroid-resistant *Anopheles* species increase oxidative stress to the detriment of *Plasmodium* survival (23). Because enzymatic detoxification of insecticides is also an important resistance mechanism in *Anopheles* mosquitoes (11), enhanced detoxification may interact with fungal metabolites, such as cyclic peptide toxins (15, 20), and could reduce the effect of these virulence factors.

The compatibility of fungus- and insecticide-based control methods also will depend on the effect of fungal infection on insecticide-resistant mosquito mortality and resistance levels. Studies on insect hosts other than mosquitoes have indicated that fungal infection can act synergistically with insecticides, increasing the impact of otherwise sublethal insecticide doses (24–26). Mixtures of *M. anisopliae* and deltamethrin were shown to enhance virulence of both components when tested against ticks, indicating synergistic effects that would enhance the effectiveness of low fungi and insecticide concentrations (27). In contradiction to these findings, however, studies on the wax moth *Galleria mellonella* have indicated that the elevation of detoxifying enzymes in response to infection with *M. anisopliae* increases host resistance to organophosphate insecticides (28). To ensure the compatibility of fungal biopesticides and chemical control tools, such potential adverse effects on resistance levels will have to be excluded for anophelines.

Author contributions: B.D.B., R.H.H., L.L.K., B.G.J.K., and M.C. designed research; M.F., J.C.M., and C.K.K. performed research; M.B.T. contributed new reagents/analytic tools; M.F. analyzed data; and M.F., M.B.T., B.G.J.K., and M.C. wrote the paper.

The authors declare no conflict of interest.

<sup>1</sup>To whom correspondence should be addressed. E-mail: bart@malaria-world.com.


This article contains supporting information online at [www.pnas.org/cgi/content/full/0908530106/DCSupplemental](http://www.pnas.org/cgi/content/full/0908530106/DCSupplemental).

So far, there have been no reports on the effects of insecticide resistance mechanisms on mosquito susceptibility to fungal infection or the effects of fungal infection on mosquito insecticide resistance levels. In this study, therefore, we investigated the effectiveness of two potential fungal biocontrol pathogens, *M. anisopliae* and *B. bassiana*, against insecticide-resistant *Anopheles* mosquitoes. Effects of insecticide resistance status on fungal susceptibility and effects of fungal infection on insecticide resistance levels were tested in a diverse suite of insecticide-resistant *Anopheles* colonies.

## Results

**Mosquito Susceptibility to Fungus Infection.** We studied the infectivity and virulence of *B. bassiana* in a permethrin-resistant *A. funestus* colony (Af<sup>Perm</sup>), two dichlorodiphenyltrichloroethane (DDT)-resistant *A. arabiensis* colonies (Aa<sub>1</sub><sup>DDT</sup> and Aa<sub>2</sub><sup>DDT</sup>), one bendiocarb-resistant *A. gambiae* s.s. colony (Ag<sup>Bend</sup>), and the corresponding baseline colonies from which each was selected. Replicate samples of mosquitoes were infected with the fungus by using a standard exposure bioassay previously shown to provide reliable infection in the laboratory (16). *B. bassiana* caused 100% mortality in all eight colonies within 8–20 days (Fig. 1), with levels of infection exceeding 95% [confirmed by fungal sporulation on mosquito cadavers (16)]. Cox regression analyses showed the effect of *Beauveria* infection on mosquito survival to be significant in all tested colonies ( $P < 0.001$ ). Additionally, a highly significant interaction between fungus treatment and insecticide resistance was found in *A. funestus* ( $P = 0.002$ ) and *A. gambiae* s.s. ( $P < 0.001$ ), indicating quantitative differences in the effect of fungus infection on mosquito survival between the baseline and insecticide-resistant colonies. *Beauveria* infection reduced survival more strongly in permethrin-resistant *A. funestus* mosquitoes [hazard ratio (HR) = 47,241.42;  $P < 0.001$ ] than in its baseline colony (HR = 28.22;  $P < 0.001$ ) yet reduced survival less strongly in bendiocarb-resistant *A. gambiae* s.s. (HR = 5.70;  $P < 0.001$ ) than in the baseline mosquitoes (HR = 71.70;  $P < 0.001$ ). However, similar differences in HRs were observed in the corresponding uninfected control mosquitoes, a significantly lower daily mortality rate in permethrin-resistant *A. funestus* mosquitoes (HR = 0.003;  $P = 0.005$ ) and higher mortality rate in bendiocarb-selected *A. gambiae* s.s. mosquitoes (HR = 28.20;  $P < 0.001$ ) compared with those of their baseline colonies, suggesting that these quantitative differences in the effect of fungus infection on survival are caused by inherent differences in laboratory colony longevity rather than any effect of resistance per se.

**Effect of Fungal Infection on Insecticide Resistance.** To investigate whether fungal infection affects the expression of insecticide resistance, we conducted a series of experiments to examine prelethal effects of fungal infection on insecticide sensitivity in resistant mosquitoes. First, we assessed the effect of *B. bassiana* or *M. anisopliae* infection on the expression of permethrin resistance in three mosquito strains with known levels of resistance to this insecticide, permethrin-resistant *A. funestus* (Af<sup>Perm</sup>), DDT-resistant *A. arabiensis* (Aa<sub>2</sub><sup>DDT</sup>), and a recently established, unselected, multiple-resistant strain of *A. gambiae* s.s. (Ag<sup>MR</sup>). Replicate sets of mosquitoes were infected with either *Beauveria* or *Metarhizium* and exposed to permethrin 3 days later, using standard World Health Organization (WHO) insecticide assay protocols. Mortality was recorded 24 h after insecticide exposure and compared with permethrin-induced mortality in uninfected controls. In all three mosquito strains, permethrin induced significantly higher mortality in mosquitoes preinfected with *Beauveria* ( $P < 0.001$ ) or *Metarhizium* ( $P < 0.001$ ) than that in the uninfected groups (Fig. 2A). Both fungi induced similar increases in susceptibility to permethrin. In *A. arabiensis*, we found significantly higher mortality levels induced



**Fig. 1.** Effect of fungus infection on baseline (susceptible) and insecticide-resistant mosquito survival. Mean ( $\pm$ SEM) cumulative proportional survival of *B. bassiana*-infected (red) and uninfected control mosquitoes (black) of four different mosquito strains, *A. funestus* (Af), two strains of *A. arabiensis* (Aa<sub>1</sub> and Aa<sub>2</sub>), and *A. gambiae* s.s. (Ag). Survival of baseline colonies is shown on the left and colonies selected for resistance to either permethrin (Perm), DDT, or bendiocarb (Bend) on the right. Data represent nine replicates, each containing  $\approx$ 30 females, except for the baseline Af colony where  $n = 6$  for both curves.

by *Beauveria* infection compared with *Metarhizium* infection ( $\chi^2 = 36.04$ ,  $P < 0.001$ ), which may be caused by differences in pathogenicity of the two fungal species or a different insecticide resistance mechanism in this mosquito species.

Using the Af<sup>Perm</sup> line, we also examined the effects of *B. bassiana* infection on permethrin resistance levels 5 days after fungal exposure. This revealed a significant ( $\chi^2 = 18.38$ ,  $P < 0.001$ ) increase in permethrin-induced mortality compared with that of the 3-day group (Fig. 2B). However, mortality of the uninfected control group was also significantly higher at 5 days ( $\chi^2 = 57.22$ ,  $P < 0.001$ ), which is consistent with age-dependent responses to insecticide exposure found in several mosquito species (29, 30) and suggests equivalent relative effects of fungus, at least over the initial time course of infection.

The effect of fungus infection on DDT resistance was assessed by using DDT-resistant colonies of *A. arabiensis* (Aa<sub>2</sub><sup>DDT</sup>) and *A. gambiae* s.s. (Ag<sup>MR</sup>). We preinfected replicate samples of mosquitoes with *B. bassiana* spores and compared susceptibility to DDT between the infected and uninfected groups 3 days later. For both tested species, infection with *B. bassiana* led to signif-



fungus was allowed to proliferate for 3 days in the insect before being exposed to the insecticide. There were no differences in infection percentages between insecticide-exposed and nonexposed groups, but that could be a result of already extensive fungus growth at day 3. Especially for testing the efficacy of fungus–insecticide combinations, testing the effect of neurotoxic and other classes of insecticides on fungal infectivity and virulence would be interesting.

Increased knowledge on fungus–insect interactions will augment options for improving fungus-based applications against mosquitoes. For example, modification of fungal spores to enhance their virulence can be used to improve the commercial effectiveness of fungus-based control methods. Genetic alterations that caused overproduction of a cuticle-degrading protease have been shown to effectively increase the speed of kill by the fungus (34). Furthermore, exploring the effectiveness of fungi against mosquito strains with other resistance mechanisms, such as resistance to microbial agents such as *Bacillus thuringiensis* var. *israelensis* (B.t.i.) or insect growth regulators such as methoprene, would substantiate further the usefulness of fungus-based biological control tools against mosquitoes where other current control measures are failing.

Overall, the significant reductions in mosquito survival and insecticide resistance levels induced by fungal infection support the potential use of fungal biopesticides against mosquito vectors in areas where insecticide resistance levels are increasing, potentially adding new product options to the very limited selection of chemicals currently available. With their relatively slow speed of kill, considered to dramatically reduce the selection pressure for resistance development while killing mosquitoes before being able to transmit the malaria parasite (15), fungal biopesticides may provide a sustainable vector control tool. The susceptibility of insecticide-resistant mosquitoes to fungal pathogens adds weight to the possibility of using biopesticides within insecticide resistance management strategies, such as rotations or mosaics (10), to slow the spread of resistance (15, 17). The use of oil-formulated spores in point sources such as black cotton cloths (14) or African water storage pots (35) have shown potential for field implementation and would allow for the integration of fungi in existing control measures. Like protozoan (36) or nematode infections (37), fungal pathogen infection exerts an additional fitness cost for the insect. Because these costs are associated with a slower spread of resistance (38), the additional burden of a fungal infection may reduce the speed of insecticide resistance formation in anopheline vectors. Moreover, with fungal infection reducing the expression of permethrin and DDT resistance, developing “combination treatments” may enhance the efficacy and effective lifespan of key insecticides where resistance has reached high levels. Together, these findings provide a compelling case for viewing biopesticides and chemical insecticides not as mutually exclusive but as complementary technologies that may improve the efficiency and sustainability of integrated malaria vector control programs.

## Materials and Methods

**Mosquitoes.** An overview of mosquito colony names, abbreviations, resistance selection, and origins is given in Table S1. The *A. funestus* colonies (A<sup>fPerm</sup>) originated from collections in southern Mozambique. Mosquitoes from the baseline colony (FUMOZ) were selected for high levels of resistance to the pyrethroid permethrin in the insectary of the National Institute for Communicable Diseases (Johannesburg, South Africa) for 2 years, resulting in the colony FUMOZ-R, of which adults show 0–1% mortality when exposed to 1% lambda-cyhalothrin for 1 h (31). The two *A. arabiensis* colonies used (Aa<sub>1</sub><sup>DDT</sup> and Aa<sub>2</sub><sup>DDT</sup>) originated from Mamfene, KwaZulu-Natal, South Africa (MBN) and from Sennar, south-central Sudan (SENN), respectively. The SENN baseline colony was selected for DDT resistance for 16 generations, after which SENN-DDT adults showed 12.1% mortality when exposed to 4% DDT and 0% mortality when exposed to 0.75% permethrin for 1 h (30). For this study, adults of the F50–F54 generation were used, which were shown to have lower

baseline resistance levels to DDT and permethrin (Fig. 2 A and C). The *A. gambiae* s.s. colony used in survival assays (Ag<sup>Bend</sup>) originated from Obuasi, Ghana (SOG), of which a bendiocarb-resistant colony (BENROG) was selected in the laboratory. The *A. gambiae* s.s. colony used in insecticide resistance assays (Ag<sup>MR</sup>) originated from Ahafo, Ghana (GAH). This colony has not been selected for resistance to insecticides in the laboratory but carries quantified levels of resistance to all four classes of insecticides. Both SOG and FUMOZ baseline colonies exhibit low levels of resistance, which is increased by orders of magnitude in the selected lines. A summary of mosquito insecticide resistance or susceptibility to insecticides approved by WHO is given in Table S2.

Larvae were reared in plastic bowls filled with distilled water. For *A. funestus*, the water was supplemented with green algae. Larval food was supplied daily and contained a mixture of finely crushed dog biscuit and brewer's yeast (31). Adults were collected daily from the bowls and transferred to holding cages in which cotton wool soaked in 10% glucose solution was provided. All species were maintained at 25 °C and 80% relative humidity with a 12-h day/night photoperiod and artificial 45-min dusk/dawn cycles. For experiments, 2- to 5-day-old mosquitoes were used.

**Fungus.** *M. anisopliae* var. *anisopliae* (Metsch.) Sorokin, isolate ICIPE-30 (14) (courtesy N. Maniania, International Centre of Insect Physiology and Ecology, Nairobi, Kenya) and *B. bassiana* isolate IMI 391510 (13) were used. Both fungi were produced through solid-state fermentation with glucose-impregnated hemp (courtesy F. van Breukelen and M. Jumbe, Wageningen University and Research Centre, Wageningen, The Netherlands) of which conidia were dried and stored in the dark at 4 °C.

The viability of the conidia was assessed by mixing some dry spores of each stock in oil (Shell Ondina Oil 917) and plating a drop on Sabouraud dextrose agar. After 20–26 h of incubation at 27 °C, the proportion of germinated conidia was determined with a light microscope at a magnification of 400×. Stocks showing 85% or higher sporulation were used for experiments.

**Fungal Exposures.** Adult mosquitoes were exposed to 100 mg of dry conidia by using the suspensor setup previously shown to give reliable infections, as described by Scholte et al. (16). Female mosquitoes were exposed to one suspensor in a holding cage for 24 h, after which the suspensor was replaced with clean cotton wool soaked in 10% glucose solution. Control mosquitoes were exposed in the same way but to a suspensor without fungus.

**Survival Bioassays.** The effect of fungal infection on mosquito survival was tested in baseline and insecticide-resistant colonies of *A. funestus* (A<sup>fPerm</sup>), *A. arabiensis* from South Africa (Aa<sub>1</sub><sup>DDT</sup>), *A. arabiensis* from Sudan (Aa<sub>2</sub><sup>DDT</sup>), and *A. gambiae* s.s. from Obuasi, Ghana (Ag<sup>Bend</sup>). For each colony, nine test and nine control replicates were performed on 3 consecutive days, exposing ≈30 mosquitoes per replicate to dry spores of *B. bassiana* or control suspensors for 24 h. For the baseline *A. funestus* colony, six replicates were performed. Mosquito mortality was recorded daily, and cadavers were removed from each holding cage, dipped in 70% ethanol, and placed on moist filter paper in sealed Petri dishes. To verify infection, these were incubated at 25 °C for 3 or more days and assessed for fungus sporulation (i.e., emerging hyphae) by using a dissection microscope (16).

**Permethrin Resistance Assays.** The resistant colonies of *A. funestus* (A<sup>fPerm</sup>), *A. arabiensis* from Sudan (Aa<sub>2</sub><sup>DDT</sup>), and *A. gambiae* s.s. from Ahafo, Ghana (Ag<sup>MR</sup>) were used to test the effect of fungal infection on permethrin resistance. A 3-day waiting period was chosen between fungal exposures and assessments for insecticide resistance to allow for some progression of the fungal infection while not losing large numbers through death. Mosquitoes from the same cohort received either a control, *Beauveria*, or *Metarhizium* treatment, of which 25 females per treatment were exposed 3 days later to a control paper and 25 females to a filter paper treated with 0.75% permethrin for 1 h, according to WHO protocol (39). Mosquitoes were then transferred to clean holding tubes and provided with 10% glucose solution by using cotton balls, and the proportion of dead mosquitoes was scored 24 h after insecticide exposure. Five replicates were performed per mosquito species and for each group; mortality per replicate exposed to insecticide was corrected by using mortality data of counterparts exposed to control papers, according to Abbott's formula (39). After mortality measurements, mosquitoes were removed from the exposure tubes with an aspirator, killed through drowning in 70% alcohol, and checked for fungal infection as described above.

With the same methods, the effect of a more advanced *Beauveria* infection was tested on *A. funestus* (A<sup>fPerm</sup>) by measuring the permethrin-resistance levels of control and *Beauveria*-infected mosquitoes 5 days after fungus exposure. Five replicates of 25 mosquitoes each were performed, and mor-

tality of permethrin-exposed mosquitoes was corrected for control mosquito mortality.

**DDT Resistance Assays.** The effect of *Beauveria* infection on DDT resistance was tested in five separate experiments in DDT-resistant *A. arabiensis* from Sudan (Aa<sub>2</sub><sup>DDT</sup>) and *A. gambiae* s.s. from Ghana (Ag<sup>MR</sup>). Three days after *Beauveria* exposure, control and infected mosquitoes (25 females per group) were exposed to 4% DDT papers or untreated papers, and mortality was measured 24 h later and corrected for control mortality as described for the permethrin assays.

**Data Analysis.** Differences in the computed survival curves of treated and control mosquitoes were analyzed by using Cox regression analyses (40) with SPSS 15.0 software. Mortality data were analyzed separately for each tested mosquito species, including all replicates of the control and fungus-infected

groups for both the baseline and the insecticide-resistant colony. A full model, with all main effects and possible interactions included, was calculated to estimate the effects of *Beauveria* on the hazard ratio. *Beauveria*-infected mosquito survival was analyzed separately to quantify and compare the effect of fungus infection between the resistant and the baseline colonies.

Mortality of the permethrin- and DDT-exposed groups was corrected for the mortality of their corresponding control groups by using Abbott's formula (39). Corrected mosquito mortality was compared by using a  $\chi^2$  goodness-of-fit test with GenStat 9.0 software.

**ACKNOWLEDGMENTS.** We thank S. Koenraadt for assistance with statistical analyses and J. Hemingway and A. F. Read for comments and advice. This work was supported by the Adessium Foundation (Reeuwijk, The Netherlands), Dutch Scientific Organization VIDI Grant 864.03.004 (to B.G.J.K.), and the South African Research Chairs Initiative of the Department of Science and Technology with the National Research Foundation (M.C.).

- Awolola TS, Brooke BD, Hunt RH, Coetzee M (2002) Resistance of the malaria vector *Anopheles gambiae* s.s. to pyrethroid insecticides, in south-western Nigeria. *Ann Trop Med Parasitol* 96:849–852.
- Diabate A, et al. (2004) The spread of the Leu-Phe *kdr* mutation through *Anopheles gambiae* complex in Burkina Faso: Genetic introgression and de novo phenomena. *Trop Med Int Health* 9:1267–1273.
- Tia E, et al. (2006) Pyrethroid and DDT resistance of *Anopheles gambiae* s.s. (Diptera: Culicidae) in five agricultural ecosystems from Cote-d'Ivoire. *Bull Soc Pathol Exot* 99:278–282.
- Hargreaves K, et al. (2000) *Anopheles funestus* resistant to pyrethroid insecticides in South Africa. *Med Vet Entomol* 14:181–189.
- Brooke BD, et al. (2001) Bioassay and biochemical analyses of insecticide resistance in southern African *Anopheles funestus* (Diptera: Culicidae). *Bull Entomol Res* 91:265–272.
- Abdalla H, et al. (2007) Insecticide susceptibility and vector status of natural populations of *Anopheles arabiensis* from Sudan. *Trans R Soc Trop Med Hyg* 102:263–271.
- Hargreaves K, et al. (2003) *Anopheles arabiensis* and *An. quadriannulatus* resistance to DDT in South Africa. *Med Vet Entomol* 17:417–422.
- World Health Organization (2008) *The Global Malaria Action Plan, Roll Back Malaria Partnership* (WHO, Geneva).
- Kelly-Hope L, Ranson H, Hemingway J (2008) Lessons from the past: Managing insecticide resistance in malaria control and eradication programmes. *Lancet Infect Dis* 8:387–389.
- Nauen R (2007) Insecticide resistance in disease vectors of public health importance. *Pest Manag Sci* 63:628–633.
- Hemingway J, Ranson H (2000) Insecticide resistance in insect vectors of human disease. *Annu Rev Entomol* 45:371–391.
- Zaim M, Guillet P (2002) Alternative insecticides: An urgent need. *Trends Parasitol* 18:161–163.
- Blanford S, et al. (2005) Fungal pathogen reduces potential for malaria transmission. *Science* 308:1638–1641.
- Scholte E-J, et al. (2005) An entomopathogenic fungus for control of adult African malaria mosquitoes. *Science* 308:1641–1642.
- Thomas MB, Read AF (2007) Can fungal biopesticides control malaria? *Nat Rev Microbiol* 5:377–383.
- Scholte EJ, Njiru BN, Smallegange RC, Takken W, Knols BGJ (2003) Infection of malaria (*Anopheles gambiae* s.s.) and filariasis (*Culex quinquefasciatus*) vectors with the entomopathogenic fungus *Metarhizium anisopliae*. *Malar J* 2:29.
- Read AF, Lynch PA, Thomas MB (2009) How to make evolution-proof insecticides for malaria control. *PLoS Biol* 7:e1000058.
- Beier JC (1998) Malaria parasite development in mosquitoes. *Annu Rev Entomol* 43:519–543.
- Scholte E-J, Knols BGJ, Takken W (2006) Infection of the malaria mosquito *Anopheles gambiae* with the entomopathogenic fungus *Metarhizium anisopliae* reduces blood feeding and fecundity. *J Invertebr Pathol* 91:43–49.
- Clarkson JM, Charnley AK (1996) New insights to the mechanisms of fungal pathogenesis in insects. *Trends Microbiol* 4:197–203.
- St. Leger RJ, Joshi L, Roberts D (1998) Ambient pH is a major determinant in the expression of cuticle-degrading enzymes and hydrophobin by *Metarhizium anisopliae*. *Appl Environ Microbiol* 64:709–713.
- McCarroll L, et al. (2000) Insecticides and mosquito-borne disease. *Nature* 407:961–962.
- McCarroll L, Hemingway J (2002) Can insecticide resistance status effect parasite transmission in mosquitoes? *Insect Biochem Mol Biol* 32:1345–1351.
- Ericsson JD, Kabaluk JT, Goettel MS, Myers JH (2007) Spinosad interacts synergistically with the insect pathogen *Metarhizium anisopliae* against two exotic wireworms, *Agriotes lineatus* and *Agriotes obscurus* (Coleoptera: Elateridae) *J Econ Entomol* 100:31–38.
- Furlong MJ, Groden E (2001) Evaluation of synergistic interactions between the Colorado potato beetle (Coleoptera: Chrysomelidae) pathogen *Beauveria bassiana* and the insecticides, imidacloprid, and cyromazine. *J Econ Entomol* 94:344–356.
- Pachamuthu P, Kamble ST (2000) In vivo study on combined toxicity of *Metarhizium anisopliae* 0 (Deuteromycotina: Hyphomycetes) strain ESC-1 with sublethal doses of chlorpyrifos, propetamphos, and cyfluthrin against German cockroach (Diptoptera: Blattellidae). *J Econ Entomol* 93:60–70.
- Bahiense TC, Fernandes EKK, Bittencourt VREP (2006) Compatibility of the fungus *Metarhizium anisopliae* and deltamethrin to control a resistant strain of *Boophilum microplus* tick. *Vet Parasitol* 141:319–324.
- Serebrov VV, et al. (2006) Effect of entomopathogenic fungi on detoxification enzyme activity in greater wax moth *Galleria mellonella* L. (Lepidoptera, Pyralidae) and role of detoxification enzymes in development of insect resistance to entomopathogenic fungi. *Biol Bull* 33:581–586.
- Lines JD, Nasser NS (1991) DDT resistance in *Anopheles gambiae* declines with mosquito age. *Med Vet Entomol* 5:261–265.
- Matambo TS, et al. (2007) Insecticide resistance in the malarial mosquito *Anopheles arabiensis* and association with the *kdr* mutation. *Med Vet Entomol* 21:97–102.
- Hunt RH, Brooke BD, Pillay C, Koekemoer LL, Coetzee M (2005) Laboratory selection for and characteristics of pyrethroid resistance in the malaria vector *Anopheles funestus*. *Med Vet Entomol* 19:271–275.
- Amenya DA, et al. (2005) Isolation and sequence analysis of P450 genes from a pyrethroid-resistant colony of the major malaria vector *Anopheles funestus*. *DNA Seq* 16:437–445.
- Djouaka RF, et al. (2008) Expression of the cytochrome P450's, *CYP6P3* and *CYP6M2* are significantly elevated in multiple pyrethroid resistant populations of *Anopheles gambiae* s.s. from Southern Benin and Nigeria. *BMC Genomics* 9:538.
- St. Leger RJ, Joshi L, Bidochka MJ, Roberts DW (1996) Construction of an improved mycoinsecticide overexpressing a toxic protease. *Proc Natl Acad Sci USA* 93:6349–6354.
- Farenhorst M, et al. (2008) African water storage pots for the delivery of the entomopathogenic fungus *Metarhizium anisopliae* to the African malaria vectors *Anopheles gambiae* s.s. and *Anopheles funestus*. *Am J Trop Med Hyg* 78:910–916.
- Hogg JC, Hurd H (1995) Malaria-induced reduction of fecundity during the first gonotrophic cycle of *Anopheles stephensi* mosquitoes. *Med Vet Entomol* 9:176–180.
- Gassmann AJ, Stock SP, Carriere Y, Tabashnik BE (2006) Effect of entomopathogenic nematodes on the fitness cost of resistance to Bt toxin Cry1Ac in pink bollworm (Lepidoptera: Gelechiidae). *J Econ Entomol* 99:920–926.
- Gassmann AJ, Carriere Y, Tabashnik BE (2009) Fitness costs of insect resistance to *Bacillus thuringiensis*. *Annu Rev Entomol* 54:147–163.
- World Health Organization (1998) *Test Procedures for Insecticide Resistance Monitoring in Malaria Vectors, Bio-Efficacy and Persistence of Insecticides on Treated Surfaces* (WHO, Geneva).
- Cox DR (1972) Regression models and life tables. *J R Stat Soc* 34:187–220.