

Corrections

BIOCHEMISTRY

Correction for “Confinement of caspase-12 proteolytic activity to autoprocessing,” by Sophie Roy, Jeffrey R. Sharom, Caroline Houde, Thomas P. Loisel, John P. Vaillancourt, Wei Shao, Maya Saleh, and Donald W. Nicholson, which appeared in issue 11, March 18, 2008, of *Proc Natl Acad Sci USA* (105:4133–4138; first published March 10, 2008; 10.1073/pnas.0706658105).

The authors note that Fig. 6 appeared incorrectly. There was an error in the alignment of the molecular mass markers, and minor adjustments have been made to the assignment of caspase-12 bands. The authors also note that the source of the anti-caspase-12 antibody for Fig. 6 was Sigma (clone 14F7). This error does not affect the conclusions of the article. The corrected figure and its corresponding legend appear below.




Fig. 6. ProCaspase-12 forms a complex with caspase-1 and is partially autoprocessed in the complex. HEK293T cells were cotransfected with expression vectors harboring Flag-tagged procaspase-1 (all lanes) plus either procaspase-12 (lanes 1 and 4) or the catalytically incapacitated C²⁹⁹A mutant (lanes 2 and 5). After 24 h, cells were harvested and lysed. One tenth of the lysate was directly applied to SDS/PAGE (lanes 1 and 2), and the remainder was immunoharvested with antibodies directed against the caspase-1 Flag epitope tag (lanes 4 and 5; lane 3 was processed in the same way, except that only lysis buffer was used). Immunoblotting for the large subunit (p20) of caspase-12 revealed that procaspase-12 and the resulting autocleavage product were both immunoharvested with caspase-1. The asterisk indicates a band of unknown identity that is detected by prebleed control serum.

www.pnas.org/cgi/doi/10.1073/pnas.1302753110

CHEMISTRY

Correction for “Direct asymmetric vinylogous Michael addition of cyclic enones to nitroalkenes via dienamine catalysis,” by Giorgio Bencivenni, Patrizia Galzerano, Andrea Mazzanti, Giuseppe Bartoli, and Paolo Melchiorre, which appeared in issue 48, November 30, 2010, of *Proc Natl Acad Sci USA* (107:20642–20647; first published June 21, 2010; 10.1073/pnas.1001150107).

The authors note that they incorrectly assigned the structure of the reaction product reported in Scheme 4. The published structure represents the γ -aminated adduct **8**, when it should instead be the α -analogue arising from an α -site selective pathway. As a result of this, Scheme 4 and its related comments should be removed from the article.

On page 20642, left column, within the Abstract, lines 16–18, “Finally, we describe the extension of the dienamine catalysis-induced vinylogous nucleophilicity to the asymmetric γ -amination of cyclohexene carbaldehyde” should be removed from the article.

On page 20645, right column, third full paragraph, lines 1–8, to page 20646, left column, first paragraph, lines 1–2, “Finally, to explore the potential of the chiral primary amine-induced vinylogous nucleophilicity, we wondered whether this unique reactivity concept may be translated to an aldehyde derivative adorned with a six-membered ring scaffold, reminiscent of the β -substituted cyclohexanone framework. Although the vinylogous Michael addition of 1-cyclohexene-1-carboxaldehyde **7** to nitrostyrene **2a** did not proceed at all, the combination with *tert*-butylazodicarboxylate under the catalysis of **A** furnished the γ -amination product **8** with perfect regio- and enantioselectivity (Scheme 4)” should be removed from the article.

These errors do not affect the conclusions of the article of the vinylogous Michael addition of cyclic enones to nitroalkenes. The ability of primary amine catalysis to address the synthetic issue connected with the enantioselective carbon–carbon bond formation gamma to a carbonyl group, promoting vinylogous nucleophilicity upon selective activation of unmodified cyclic unsaturated ketones, is fully supported by the separated results presented in Tables 1, 2, and 3, and Schemes 2 and 3.

www.pnas.org/cgi/doi/10.1073/pnas.1302980110

MICROBIOLOGY

Correction for “Commensal bacteria play a role in mating preference of *Drosophila melanogaster*,” by Gil Sharon, Daniel Segal, John M. Ringo, Abraham Hefetz, Ilana Zilber-Rosenberg, and Eugene Rosenberg, which appeared in issue 46, November 16, 2010, of *Proc Natl Acad Sci USA* (107:20051–20056; first published November 1, 2010; 10.1073/pnas.1009906107).

The authors note the following: “The mating frequencies reported in Table 1 of this paper did not follow a multinomial distribution, making the statistical analysis inapplicable. This problem was obviated by considering only the first matings in each experimental unit and computing odds ratios. After submitting the paper, we continued to perform experiments

identical in design to those we reported. In the table below, we combined the results of those additional replicate experiments with those already reported. From the new analysis, we now find that experiment 4, in which flies were infected with a mixture of *Lactobacillus* spp., assortative mating was not restored. Otherwise, the conclusions of the article were not changed by our reanalysis. We acknowledge the statistical advice of Dan Yekutieli and thank Tal Lahav for calculating the odds ratios and their 95% confidence intervals, and for performing the chi-squared tests presented in the corrected Table 1.”

The corrected Table 1 appears below.

Table 1. The role of bacteria in diet-induced mating preference of *D. melanogaster*

Experiment	Fly treatment*	N [†]	OR [‡]	95% CI	P value [‡]
1	Starch-grown × CMY-grown	18	3.21	2.14–4.81	1.8 × 10 ⁻⁸
2	Experiment 1 after antibiotics	11	1.04	0.63–1.71	0.9888
3	Experiment 2 after infection of starch-grown flies with homologous bacteria [§]	6	2.68	1.40–5.11	0.0477
4	Experiment 3 with <i>Lactobacillus</i> spp. replacing homologous bacteria	4	1.76	0.74–4.19	0.2912
5	Experiment 3 with <i>Lactobacillus plantarum</i> replacing homologous bacteria	7	2.14	1.35–3.39	0.0019
6	Infection control (no added bacteria)	4	1.26	0.53–3.00	0.7712

*After all treatments, the flies were grown for one generation in CMY medium before performing the mating preference test.

[†]N is the number of replicate experiments.

[‡]Cochran-Mantel-Haenszel Odds Ratio and P value are from the Cochran-Mantel-Haenszel Chi-squared test (34, 35).

[§]Antibiotic-treated starch- and CMY-grown flies were infected with bacteria isolated from their respective growth medium (before antibiotic treatment).

34. Mantel N, Haenszel W (1959) Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst* 22(4):719–748.

35. Cochran WD (1954) Some methods for strengthening the common χ^2 tests. *Biometrics* 10:417–451.

www.pnas.org/cgi/doi/10.1073/pnas.1302326110

MEDICAL SCIENCES

Correction for “Interaction of intracellular β amyloid peptide with chaperone proteins,” by Virginia Fonte, Vadim Kapulkin, Andrew Taft, Amy Fluet, David Friedman, and Christopher D. Link, which appeared in issue 14, July 9, 2002, of *Proc Natl Acad Sci USA* (99:9439–9444; first published June 27, 2002; 10.1073/pnas.152313999).

The authors note that the author name Vadim Kapulkin should instead appear as Wadim Jan Kapulkin. The corrected author line appears below. The online version has been corrected.

**Virginia Fonte, Wadim Jan Kapulkin, Andrew Taft,
Amy Fluet, David Friedman, and Christopher D. Link**

www.pnas.org/cgi/doi/10.1073/pnas.1302545110

SYSTEMS BIOLOGY

Correction for “Expanded methyl-sensitive cut counting reveals hypomethylation as an epigenetic state that highlights functional sequences of the genome,” by Alejandro Colaneri, Nickolas Staffa, David C. Fargo, Yuan Gao, Tianyuan Wang, Shyamal D. Peddada, and Lutz Birnbaumer, which appeared in issue 23, June 7, 2011, of *Proc Natl Acad Sci USA* (108:9715–9720; first published May 20, 2011; 10.1073/pnas.1105713108).

The authors note that, within the corresponding author footnote on page 9715, the email address “colaneria@niehs.nih.gov” should instead appear as “acolaneri_2000@yahoo.com.ar”.

www.pnas.org/cgi/doi/10.1073/pnas.1302473110

Commensal bacteria play a role in mating preference of *Drosophila melanogaster*

Gil Sharon^a, Daniel Segal^a, John M. Ringo^b, Abraham Hefetz^c, Ilana Zilber-Rosenberg^d, and Eugene Rosenberg^{a,1}

^aDepartment of Molecular Microbiology and Biotechnology, Tel Aviv University, Tel Aviv 69978, Israel; ^bSchool of Biology and Ecology, University of Maine, Orono, ME 04469; ^cDepartment of Zoology, Tel Aviv University, Tel Aviv 69978, Israel; and ^d18 Rachavat Ilan St., Givat Shmuel 51905, Israel

Edited by R. John Collier, Harvard Medical School, Boston, MA, and approved September 28, 2010 (received for review July 12, 2010)

Development of mating preference is considered to be an early event in speciation. In this study, mating preference was achieved by dividing a population of *Drosophila melanogaster* and rearing one part on a molasses medium and the other on a starch medium. When the isolated populations were mixed, “molasses flies” preferred to mate with other molasses flies and “starch flies” preferred to mate with other starch flies. The mating preference appeared after only one generation and was maintained for at least 37 generations. Antibiotic treatment abolished mating preference, suggesting that the fly microbiota was responsible for the phenomenon. This was confirmed by infection experiments with microbiota obtained from the fly media (before antibiotic treatment) as well as with a mixed culture of *Lactobacillus* species and a pure culture of *Lactobacillus plantarum* isolated from starch flies. Analytical data suggest that symbiotic bacteria can influence mating preference by changing the levels of cuticular hydrocarbon sex pheromones. The results are discussed within the framework of the hologenome theory of evolution.

holobiont | hologenome | *Lactobacillus* | microbiota | symbiosis

Mating preference is considered one of the mechanisms for the origin of new species (1–4). Experimentally, mating preference in flies has been shown to occur when populations were divided and reared separately for many generations under different environmental conditions (5), such as temperature and humidity (6), diet (7), and exposure to media at different pH values (8). In one of these studies, Dodd (7) reared *Drosophila pseudoobscura* on starch-based and on maltose-based media for more than 25 generations and discovered that “starch flies” preferred to mate with other starch flies and that “maltose flies” preferred to mate with other maltose flies (i.e., positive assortative mating). These data were surprising because there was no selection for the observed mating preference. Somehow mating preference developed as a correlated response when selection favored novel adaptation to the environment (namely, the different food sources).

Evidence presented here indicates that the “correlated response” driving mating preference is the emergence of different bacterial communities associated with the two fly populations grown on the different food sources: these bacteria, which are part of the normal fly microbiota, are responsible for the mating preference.

Results

An inbred laboratory strain of WT *Drosophila melanogaster* maintained on standard cornmeal-molasses-yeast (CMY) agar medium, was divided in two: flies in one group continued to be grown on CMY medium whereas flies in the other group were transferred to starch medium (Fig. 1A; *Materials and Methods*). After a fixed number of generations, each population was grown for one generation on CMY medium (to avoid the medium itself becoming the cause for mating preference) and then tested for mating preference in mating chambers. Each chamber contained four flies: one male and one female from one population and one male and one female from the other population. Individuals of one population had the tips of their wings clipped to allow

identification. Several replicates of each experiment were performed with the wing clippings alternating between populations (i.e., half of each experiment was done with wing clipping of flies from one treatment, and the other half was the reciprocal). Even though wing clipping has been previously shown not to affect mating preference in *Drosophila* (9, 10), we used a counter-balanced design for mating preference tests to control for any possible wing clipping effects.

In the first experiment, the mating preference test was performed after 11 generations (Fig. 1B). Of the 38 recorded matings, 29 were homogamic, i.e., “starch males” with “starch females” and “CMY males” with “CMY females,” whereas only nine were heterogamic (i.e., “starch” × “CMY”). These results demonstrate a significant positive sexual isolation index (SII) (11): $SII = 0.53 \pm 0.14$ (SEM) and $P = 0.0012$ (binomial test), with the following assumptions:

$$SII = \frac{\text{homogamic matings} - \text{heterogamic matings}}{n(\text{total matings})} \quad [1]$$

and

$$SEM \text{ of } SII = \sqrt{\frac{(1 - SII^2)}{n}} \quad [2]$$

SII values greater than zero indicate positive assortative mating (mating preference).

The second set of experiments was carried out exactly like the first except that the mating preference test was performed periodically, beginning after the second generation (Fig. 2). Homogamic mating occurred at all 15 times examined. Pooling all tests yielded 571 homogamic and 329 heterogamic matings: $SII = 0.27 \pm 0.02$ (SEM), $P < 0.0001$. The finding that mating preference occurred after only two generations in different growth media has not been reported previously.

A control experiment was performed in which two lines of CMY-reared flies, grown separately for 27 generations, were tested for mating preference. Of the 38 matings, 18 were homogamic and 20 were heterogamic ($SII = -0.05$, $P = 0.4357$ by binomial test), demonstrating no mating preference. Thus, difference in diet and not physical separation was the experimental variable responsible for the mating preferences reported here.

To test whether the addition of antibacterials affects mating preference, fly populations were reared on media supplemented with antibiotics (tetracycline, rifampicin, and streptomycin, together or separately; *Materials and Methods*). Following treatment with antibacterial agents, the flies’ mating preference changed from

Author contributions: G.S., J.M.R., I.Z.-R., and E.R. designed research; G.S. performed research; G.S., D.S., J.M.R., A.H., and E.R. analyzed data; and I.Z.-R. and E.R. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. E-mail: eros@post.tau.ac.il.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1009906107/-DCSupplemental.




Fig. 1. (A) Schematic representation of the experimental procedure. A population of flies was divided, serially transferred in two different media, and then examined for mating preference. After rearing the flies for a number of generations on starch or CMY media, each population was grown separately for one generation on CMY medium and then tested for mating preference. The multiple-choice mating tests were performed in 24-well plastic plates; each well contained four flies: one male and one female starch-reared and one male and one female CMY-reared. Matings were recorded every 4 min for 1 h. (B) Mating preference tests of *D. melanogaster* after growing 11 generations on starch or CMY medium.

positive assortative to random: pooling the results from 10 independent experiments yielded 267 homogamic and 263 heterogamic matings (Table 1, experiment 2; SII = 0.01 ± 0.03). Clearly,

there is a significant difference in mating preference between the antibiotic-treated and untreated flies, suggesting that symbiotic bacteria were responsible for the homogamic mating preference.

Table 1. The role of bacteria in diet-induced mating preference of *D. melanogaster*

Experiment	Fly treatment*	Matings	SII, mean \pm SEM	P value [†]
1	Starch-grown \times CMY-grown	18	0.27 ± 0.02	<0.0001
2	Experiment 1 after antibiotics	10	0.01 ± 0.03	0.4483
3	Experiment 2 after infection with homologous bacteria [‡]	4	0.22 ± 0.03	0.0024
4	Experiment 3 with <i>Lactobacillus</i> replacing homologous bacteria in starch-bred flies	4	0.16 ± 0.06	0.0392
5	Experiment 3 with <i>Lactobacillus plantarum</i> replacing homologous bacteria in starch-bred flies	5	0.19 ± 0.05	0.0004
6	Infection control (no added bacteria)	4	-0.04 ± 0.08	0.4052

*After all treatments, the flies were grown for one generation in CMY medium before performing the mating preference test.

[†]P value of the normal approximation to the binomial test. $P < 0.05$ was considered to indicate significant mating preference.

[‡]Antibiotic-treated starch- and CMY-grown flies were infected with bacteria isolated from their respective growth medium (before antibiotic treatment).




Fig. 2. Mating preference tests of *D. melanogaster* after growth for different numbers of generations on either starch or CMY medium. The SII is plotted for each generation examined. Bars indicate the SEM.

Further evidence that bacteria were responsible for the mating preference came from four independent infection experiments. Microbiota, isolated from starch or CMY media upon which the flies had grown, were inoculated into separate vials containing sterile CMY medium. Virgin, antibiotic-treated flies were then introduced into these vials (containing the starch-derived or CMY-derived bacteria on CMY medium) so that each antibiotic-treated fly population was exposed to its specific corresponding microbiota. After one generation of growth with the corresponding microbiota, positive assortative mating was regained (SII = 0.22 ± 0.03 , $P = 0.0024$; Table 1, experiment 3). Thus, the reinfected flies showed similar mating preference to flies before antibiotic treatment.

The most abundant bacteria associated with flies reared on the different media were characterized by analysis of the 16S rRNA

gene sequences of the third-generation flies (Table 2). *D. melanogaster* reared on the standard CMY medium contained the endosymbiont *Wolbachia* plus a diverse bacterial community. However, the flies transferred to starch medium contained *Wolbachia* plus 26% *Lactobacillus plantarum*. CMY-reared flies contained only 3% *L. plantarum*. Viable counts of fly homogenates showed that starch flies contained 2.3×10^5 *Lactobacillus* sp. per fly, whereas CMY flies contained only 2.6×10^4 *Lactobacillus* sp. per fly. Although this is only a limited analysis of the microbiota, it demonstrates that the associated bacteria of the CMY and starch populations are already different by the third generation and points to *L. plantarum* as a likely candidate for inducing the mating preference phenomenon. *Wolbachia* species could not have been responsible for the homogamic mating

Table 2. Bacterial communities in *D. melanogaster* grown on CMY or starch

Closest match (accession no.)	Identity, %	Representation in clone library, %*	
		CMY [†]	Starch [‡]
<i>Acetobacter pomorum</i> strain EW816 (EU096229.1)	89.81	1.49	—
<i>Acetobacter pomorum</i> strain EW816 (EU096229.1)	93.99	1.49	—
<i>Acetobacter pomorum</i> strain EW816 (EU096229.1)	94.00	1.49	—
<i>Acetobacter pomorum</i> strain EW816 (EU096229.1)	100.00	14.93	—
<i>Bacillus firmus</i> strain XJSL2-8 (GQ903397.1)	100.00	11.94	—
<i>Enterococcus faecalis</i> strain 3–12 (GU177628.1)	100.00	1.49	—
<i>Lactobacillus plantarum</i> strain IMAU:10272 (GU138600.1)	100.00	2.99	26.09
Low G+C Gram-positive bacterium T135 (AB116139.1)	99.44	5.97	—
<i>Weissella paramesenteroides</i> strain CTSP5 (EU855224.1)	97.24	1.49	—
<i>Weissella paramesenteroides</i> strain CTSP5 (EU855224.1)	99.64	4.48	—
<i>Wolbachia</i> endosymbiont of <i>Drosophila melanogaster</i> (AB360385.1)	100.00	47.77	73.91

16S rRNA gene analysis was performed on flies (third generation) grown on CMY medium for one generation before mating preference tests.

*Sequences with $\geq 99\%$ identity were clustered by DOTUR (9).

[†]Based on 64 clones.

[‡]Based on 23 clones.

preference for two reasons: (i) both fly populations contained the same *Wolbachia* wMel strain, as verified by sequencing the *wsp* gene, and (ii) *Wolbachia* was not present in flies that were antibiotic-treated and subsequently infected. The presence of *Wolbachia* before antibiotic treatment and its absence after treatment and infection was demonstrated by PCR using primers specific for both its 16S rRNA and *wsp* genes (Fig. S1). Accordingly, we next isolated *Lactobacillus* and other species from starch-reared flies, used them to infect antibiotic-treated, starch-reared flies, and then tested these flies for mating preference against antibiotic-treated, CMY-reared flies infected with CMY-derived bacteria. More than 80% of the culturable bacteria from starch-bred flies were *Lactobacillus* spp. (*Wolbachia* is not culturable). Similar percentages of Lactobacilli were found in the starch medium after fly growth.

Infection with a mixture of isolated *Lactobacillus* spp. caused a significant increase in mating preference (Table 1, experiment 4), compared with antibiotic-treated controls without added bacteria (Table 1, experiment 6). When a pure culture of *L. plantarum*, obtained from starch-bred flies, was used to infect antibiotic-treated flies, significant homogamic mating preference was obtained (Table 1, experiment 5). In the latter experiment, the data clearly demonstrate that a single bacterial species can induce mating preference. Parallel experiments using other bacterial species isolated from starch-bred flies, e.g., *Providencia rettgeri* and a mixture of 41 bacterial strains isolated from starch-bred flies had no effect on mating preference.

To discover the origin of *L. plantarum* in starch-bred flies, CMY-bred flies were treated with antibiotics before dividing them into CMY and starch media. No homogamic mating preference was observed for the three generations that were tested. In the no-antibiotic control, significant homogamic mating preference was observed already in the first generation. As the flies were reared on sterile media, these data indicate that the bacterium responsible for the mating preference, *L. plantarum*, was already present in CMY-bred flies, albeit in low numbers (consistent with the data in Table 2), and, when transferred to starch medium, were amplified and induced mating preference.

To gain some insight at a biochemical level into how the fly microbiota may influence mating behavior, we analyzed the cuticular hydrocarbon (CH) composition of antibiotic-treated and untreated CMY and starch flies (Table 3, Figs. S2 and S3, and Tables S1 and S2). In nontreated female flies, there was a significant difference between the two populations in at least four of the major CHs. In nontreated male flies, there was a significant difference between CMY and starch-bred flies in at least three CHs. In most cases, antibiotic treatment reduced the level of CHs and decreased the differences between the two populations.

These data suggest that symbiotic bacteria can influence the levels of fly sex pheromones.

Discussion

The major findings of this study are (i) diet-induced mating preference occurred in *D. melanogaster* after only one generation on different growth media and was maintained under these conditions for at least 37 generations, (ii) fly-associated commensal bacteria were responsible for the mating preference, (iii) *L. plantarum* was responsible, at least in part, for the mating preference, and (iv) the source of *L. plantarum* was the commensal microbiota of the CMY-bred flies, which were amplified in starch medium. There are abundant data demonstrating that changes in the diet of an animal result in changes in its microbiota, especially microorganisms associated with the digestive tract (13, 14). In the present case of a shift from molasses-based CMY to starch-based media, one would expect that bacteria secreting amylases would come to the fore. This is consistent with the report that *L. plantarum* is amylase-positive (15).

Mating preference has previously been achieved in long-term laboratory experimental populations by exposure to different selection pressures. In one of these studies, *D. melanogaster* populations were maintained for 30 y on media supplemented with either heavy metals or ethanol, and an SII of 0.34 was obtained (16). Part of the mating preference was a result of multiple genetic factors distributed over the chromosomes. The other part was a result of cytoplasmic factors the authors (16) attributed to *Wolbachia*, because tetracycline treatment reduced the mating preference by approximately 50%. However, the cytoplasmic factor could be any of the tetracycline-sensitive bacteria associated with the flies. In the short-term experiments described here, amplification of *L. plantarum* was responsible for mating preference, as demonstrated by direct infection experiments. It is also possible that the diverse microbiota in CMY-bred flies compared with starch-bred flies contributed to mating preference.

The data presented here provide further support for the hologenome theory of evolution (17), which posits that the holobiont (host plus its associated microorganisms) acts as a unit of selection in evolutionary change. The hologenome is defined as the sum of the genetic material of the host and its microbiota. One of the principles of the theory is that variation, an important factor in evolution, can be brought about by modification in either the host or the microbiota genomes. In this study, changes in the fly diet led to a rapid amplification of certain bacteria, especially *L. plantarum*, which was shown to be responsible for mating preference. According to the hologenome theory, microbial amplification is equivalent to gene multiplication and leads to variation in the holobiont. In a review by Andersson and Simmons (18), the authors state: "As experimental evidence

Table 3. Major differences in CH profiles of CMY and starch bred flies

Peak name	Retention time (min)	Identified compound ^a	Mean CH per fly \pm SEM, ng			
			No antibiotic treatment		Treated with antibiotics	
			CMY (n = 3)	Starch (n = 2)	CMY (n = 3)	Starch (n = 3)
Females						
F16	17.92	7-Tricosene	44.7 \pm 10.1	22.6 \pm 0.5	25.8 \pm 2.3	16.2 \pm 0.8
F24	21.08	7-Pentacosene	37.1 \pm 7.7	11.1 \pm 2.3	20.3 \pm 2.5	10.5 \pm 0.8
FF12	23.77	7,11-Heptacosadiene	117.4 \pm 21.8	172.0 \pm 5.6	50.6 \pm 8.1	59.4 \pm 8.8
F40	26.54	2-Methyloctacosane	87.2 \pm 2.5	136.2 \pm 4.3	34.2 \pm 4.3	49.0 \pm 6.1
Males						
M12	16.4	Cis-vaccenyl acetate	9.9 \pm 6.2	51.6 \pm 7.8	16.2 \pm 5.7	32.4 \pm 14.0
M16	17.92	7-Tricosene	259.1 \pm 59	495.4 \pm 39.9	181.2 \pm 15.8	415.0 \pm 10.5
M24	21.08	7-Pentacosene	146 \pm 29.8	61.5 \pm 8.4	93.1 \pm 3.5	69.7 \pm 4.8

^aBased on the GC CH profiles of *D. melanogaster* (12) and on GC-MS analyses.

accumulated, mate choice became widely recognized, but the genetic mechanisms underlying its evolution remain the subject of debate." Clearly, a rapid change in host microbiota is one mechanism for mate choice.

The mechanism by which bacteria induce mating preference remains to be elucidated. In principle, a bacterially induced mating signal could be a volatile compound emitted by the fly holobiont or a detectable compound on its surface. The odor of many animals results from microbial modification of compounds secreted by the host (19, 20) or compounds released by the microorganisms themselves (21); moreover, odor is known to influence sexual behavior (22–24). In the case of *Drosophila*, odorant receptors are located in the antennae and maxillary palps. At least five of the CHs, which have been shown to play a major role in fly mating (23), showed significant differences between starch- and CMY-bred flies (Table 3). These differences were reduced after antibiotic treatment, suggesting that symbiotic bacteria can influence the levels of fly sex pheromones and, by doing so, modify fly behavior.

How can a bacterially induced mating preference, as described here, contribute to speciation and evolution in nature? One possibility is that, in the natural world, multiple environmental factors act synergistically to differentiate the microbiota and strengthen the homogamic mating preference. For example, it is reasonable to assume that fly populations living on different nutrients will be, at least to some extent, geographically separated. The combination of partial geographic separation and diet-induced mating preference would reduce interbreeding of the populations. Slower changes in the host genome could further enhance the mating preference. The stronger the mating preference, the greater the chance that two populations will become sexually isolated, and many evolution biologists (25–28) have argued that the emergence of sexual isolation is the central event in the evolution of species.

Materials and Methods

Drosophila Stocks and Culture Conditions. A WT strain of *D. melanogaster* Oregon R was used in all experiments. Each population was maintained in three 50-mL vials, with 10 mL solid medium, at 25 °C on (i) CMY medium (0.65% agar, 7.6% cornmeal, 7.6% molasses, 5% inactivated yeast, 0.1% methyl-4-hydroxybenzoate, 0.76% ethanol, and 0.4% propionic acid) or (ii) starch medium (3% starch, 5% inactivated yeast, 1.5% agar, and 0.5% propionic acid).

Adaptation to Starch Medium. A population of flies maintained on CMY medium was transferred to a medium that contained one-half concentration CMY medium and one-half concentration of starch medium. After this one-generation adaptation step, the flies were then grown on starch medium for the rest of the experiment. Flies were transferred for one generation on CMY (regardless of their origin) before mating tests were performed.

Antibiotic Treatment. Flies were transferred into 50-mL vials containing 10 mL CMY medium, supplemented with 50 µg/mL tetracycline, 200 µg/mL rifampicin, 100 µg/mL streptomycin, or a mixture of the three antibiotic agents. Random mating data were obtained when testing antibiotic-treated flies (with each individual antibiotic, as well as the mixture) for mating preference. Controls—namely fly lines that were reared on their respective media without antibiotics—showed homogamic mating preference when tested.

Microbial Infection Experiments. Bacteria were extracted from the medium upon which the flies were reared for a single generation (either CMY or starch) by adding 10 mL of sterile PBS solution to the used medium (from one growth vial) and mixing vigorously. After allowing the solids to settle, 5 mL of the fluid was collected and centrifuged at low speed (100 × g) for 10 min to further remove solid debris. The resulting supernatant was then centrifuged at high speed (16,000 × g) for 2 min to pellet bacteria. The bacterial pellet was resuspended and washed twice in PBS solution. The final pellet was resuspended in 5 mL of PBS solution.

Flies treated with antibiotics (either CMY-reared or starch-reared) were allowed to propagate on CMY medium that was supplemented with bacteria extracted from CMY medium, starch medium, or with PBS solution (as

a control). Multiple-choice mating tests were performed using CMY flies reared on CMY medium supplemented with bacteria extracted from CMY, and starch flies reared on starch medium supplemented with bacteria from starch medium. Similar mating preference tests were performed with CMY flies reared on CMY medium with PBS solution and starch flies reared on CMY medium supplemented with PBS solution as controls.

Multiple Choice Mating Tests. For mating tests, virgin male and female flies from each treatment were anesthetized with CO₂, pooled from all relevant vials, and kept separately under a 12 h light/12 h dark cycle for 4 d on CMY medium. To test for assortative mating, a 24-cell plate was used. Each cell had a 2.5-mm-diameter hole that was capped (for fly introduction) and was lined with a 2 × 2 mm no. 3 filter paper (Whatman) soaked with deionized water. The entire apparatus was sealed with Parafilm M (Pechiney). Wings of flies were clipped for identification of source. Previous studies found no clipping effect (9, 10). To control for any clipping effects, a counterbalanced design was used: each cell had two clipped-wing flies (one male and one female) from the same population and two nonclipped flies (one male and one female) from another population. To transfer flies into the mating chamber without anesthesia, an aspirator was used to transfer them one by one. Males were introduced first, and then females, one after the other (the clipped female first).

Analysis of the Bacterial Community from CMY- and Starch-Reared Flies Using 16S rRNA Gene Clone Libraries. DNA was extracted from 10 third-generation flies (five males and five females) with the UltraClean Soil DNA kit (MoBio), according to the manufacturer's protocol. Primers 16S8F and 16S1492R (29) were used for amplification of the 16S rRNA genes from extracted flies DNA. 16S rRNA genes were amplified in a 25-µL reaction mixture consisting of 2.5 µL of 10× buffer, 0.5 µL of a 2.5-mM total deoxynucleoside triphosphate mixture, each primer at 5 µM, 10 ng of template DNA, and 1.25 U of Ex Taq DNA polymerase (TaKaRa). Amplification conditions for the PCR included an initial denaturation step of 94 °C for 3 min, followed by 30 cycles of 94 °C for 35 s, 56 °C for 35 s, and 72 °C for 45 s, and a final extension step of 72 °C for 3 min. Reaction products were checked for size and purity on 1% agarose gel.

Amplified DNA from fly samples was ligated into the pGEM-T Easy vector (Promega) according to the protocol of the manufacturer. The ligated vector and insert were transformed into competent *Escherichia coli* TG1 cells. Each clone was amplified by colony PCR with M13 forward and reverse primers in a 25-µL reaction mixture consisting of 2.5 µL of 10× buffer, 0.5 µL of a 2.5-mM total deoxynucleoside triphosphate mixture, each primer at 5 µM, and 1.25 U of Ex Taq (TaKaRa) or Extensor DNA polymerase (Thermo Fisher Scientific). Amplification conditions for the colony PCR included an initial denaturation step of 95 °C for 4.5 min, followed by 30 cycles of 95 °C for 0.5 min, 59.5 °C for 0.5 min, and 72 °C for 1 min, and a final extension step of 72 °C for 10 min. Reaction products were checked for size and purity on 1% agarose gel. PCR products were cleaned with ExoSAP-IT (USB). DNA sequencing was performed by the chain termination method using an ABI Prism sequencer (model 377, version 2.1.1; Applied Biosystems).

Sequences were aligned with ClustalX (30), and a DNA distance matrix was created with BioEdit. Sequences that had greater than 99% identity were clustered together with DOTUR (31). BLASTN (32) was then used to characterize each sequence cluster.

Lactobacillus Isolation and Infection Experiments. When *L. plantarum* had been identified as the major component of the starch fly microbiota, we isolated bacteria from starch flies, grown for one generation on CMY, on Brain-Heart Infusion (BHI) and de Man–Rogosa–Sharpe media (Difco). Flies were homogenized and the homogenate was serially diluted and plated on BHI agar and de Man–Rogosa–Sharpe agar plates. Plates were then incubated at 30 °C for 48 to 72 h. Morphologically different colonies were then picked and streaked on BHI agar plates and incubated at 30 °C for 48 to 72 h. This process was repeated twice to ensure cultures are pure. The different isolates were then identified according to their 16S rRNA genes, obtained by colony-PCR using 16S8F and 16S1492R primers (29). 16S rRNA genes of the isolated strains were amplified in a 25-µL reaction mixture consisting of 2.5 µL of 10× buffer, 0.5 µL of a 2.5-mM total deoxynucleoside triphosphate mixture, each primer at 5 µM, 10 ng of template DNA, and 1.25 U of Ex Taq DNA polymerase (TaKaRa). Amplification conditions for the PCR included an initial denaturation step of 94 °C for 3 min, followed by 30 cycles of 94 °C for 35 s, 56 °C for 35 s, and 72 °C for 45 s and a final extension step of 72 °C for 3 min. Reaction products were checked for size and purity on 1% agarose gel. PCR products were cleaned with ExoSAP-IT (USB). DNA sequencing was performed by the chain termination method in an ABI Prism sequencer (model 377, version 2.1.1; Applied Biosystems). Of the 25 isolates

that were screened, four were identified as *Lactobacillus* spp., two of which were identified by BLASTN (32) as *L. plantarum* strain IMAU80181 (accession no. GU125601.1).

At the first stage, all *Lactobacillus* spp. isolates (identified as *L. plantarum* and *Lactobacillus brevis*) were used for infection experiments, in which the isolates were grown for 48 h in liquid BHI medium at 30 °C on a rotary shaker (model 4300; New Brunswick Scientific) at 150 rpm. These cultures were then mixed and centrifuged at 16,000 × g. The pellet was resuspended in PBS solution and diluted to approximately 10⁸ cells per mL, which were then applied on the fly medium for infection (as described earlier). A subsequent infection experiment was done in which only one *L. plantarum* isolate was used to infect antibiotic-treated starch flies for subsequent mating preference tests.

Wolbachia Identification. To test whether *Wolbachia* spp. were present in the extracted DNA samples, PCR reactions using specific primers for *Wolbachia* sp. 16S rRNA and for the *Wolbachia* surface protein (*wsp*) genes were performed, using primers and PCR amplification conditions as previously described by Zhou et al. (33). Briefly, 16S rRNA and *wsp* genes were amplified in a 25-μL reaction mixture consisting of 2.5 μL of 10× buffer, 0.5 μL of a 2.5-mM total deoxynucleoside triphosphate mixture, each primer at 5 mM, 10 ng of template DNA, and 1.25 U of Ex-Taq DNA polymerase (TaKaRa). Amplification conditions for the PCR included an initial denaturation step at

94 °C for 35 min, followed by 30 cycles at 94 °C for 1 min, 55 °C for 1 min, and 72 °C for 45 s, and a final extension step at 72 °C for 35 min. Reaction products were checked for size and purity on 1% agarose gels.

Chemical Analysis of CHs. Starch or CMY flies, reared for one generation on CMY, were collected as pupae from culture vials, separated by sex, and placed for 4 d in empty vials lined with no. 3 Whatmann paper soaked in 4% sucrose. Flies were then transferred to empty glass vials and were kept at –20 °C. For CH extraction, frozen flies were divided into three groups of three flies, each in a glass vial, to which 200 μL of pentane containing 100 ng n-octadecane (C-18), as an internal standard, were added. Vials were then shaken at room temperature for 5 min at 150 rpm. One to two microliters of the extract were injected into the gas chromatography apparatus (CP 3900; Varian). Quantitative analyses of CHs were conducted by using a DB-1 fused silica column that was temperature-programmed from 150 °C (1 min of initial hold) at 5 °C/min to 300 °C. Compound quantification was done by peak integration in comparison with the internal standard. Peaks identity was verified by GC-MS as well as by comparison with the retention time data presented by Everaerts et al. (12).

ACKNOWLEDGMENTS. We thank the Ayelet Fishman laboratory for help with the GC/MS analyses and Ralph Martinez for useful comments.

- Mayr E (1963) *Animal Species and Evolution* (Harvard Univ Press, Cambridge, MA), pp xiv, 797 pp.
- Coyne JA, Orr HA (2004) *Speciation* (Sinauer, Sunderland, MA), pp xiii, 545 pp.
- McKinnon JS, et al. (2004) Evidence for ecology's role in speciation. *Nature* 429: 294–298.
- Spieth HT, Ringo JM (1983) Mating behavior and sexual isolation. *The Genetics and Biology of Drosophila*, eds Ashburner M, Carson HL, Thompson JN, Jr. (Academic Press, London), Vol 3c, pp 224–270.
- Rice WR, Hostert EE (1993) Laboratory experiments on speciation: What have we learned in 40 years? *Evolution* 47:1637–1653.
- Kilias G, Alahiotis SN, Pelecanos M (1980) A multifactorial genetic investigation of speciation theory using *Drosophila-melanogaster*. *Evolution* 34:730–737.
- Dodd DMB (1989) Reproductive isolation as a consequence of adaptive divergence in *Drosophila-pseudoobscura*. *Evolution* 43:1308–1311.
- de Oliveira AK, Cordeiro AR (1980) Adaptation of *Drosophila-Willistoni* experimental populations to extreme pH medium. I. Changes in viability and developmental rate. *Heredity* 44:111–122.
- van den Berg MJ, Thomas G, Hendriks H, van Delden W (1984) A reexamination of the negative assortative mating phenomenon and its underlying mechanism in *Drosophila melanogaster*. *Behav Genet* 14:45–61.
- Dodd DMB, Powell JR (1985) Founder-flush speciation: An update of experimental results with *Drosophila*. *Evolution* 39:1388–1392.
- Malogolowkincohen C, Simmons AS, Levene H (1965) A study of sexual isolation between certain strains of *Drosophila paulistorum*. *Evolution* 19:95–103.
- Everaerts C, Farine JP, Cobb M, Ferveur JF (2010) *Drosophila* cuticular hydrocarbons revisited: Mating status alters cuticular profiles. *PLoS ONE* 5:e9607.
- Ley RE, Lozupone CA, Hamady M, Knight R, Gordon JI (2008) Worlds within worlds: Evolution of the vertebrate gut microbiota. *Nat Rev Microbiol* 6:776–788.
- Ley RE, et al. (2008) Evolution of mammals and their gut microbes. *Science* 320: 1647–1651.
- Giraud E, Champailier A, Raimbault M (1994) Degradation of raw starch by a wild amylolytic strain of *Lactobacillus plantarum*. *Appl Environ Microbiol* 60:4319–4323.
- Koukou K, et al. (2006) Influence of antibiotic treatment and *Wolbachia* curing on sexual isolation among *Drosophila melanogaster* cage populations. *Evolution* 60: 87–96.
- Zilber-Rosenberg I, Rosenberg E (2008) Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiol Rev* 32: 723–735.
- Andersson M, Simmons LW (2006) Sexual selection and mate choice. *Trends Ecol Evol* 21:296–302.
- Hunt DWA, Borden JH (1989) Terpene alcohol pheromone production by *dendroctonus-ponderosae* and *lps-paraconfusus* (Coleoptera, Scolytidae) in the absence of readily culturable microorganisms. *J Chem Ecol* 15:1433–1463.
- Leyden JJ, McGinley KJ, Hölzle E, Labows JN, Kligman AM (1981) The microbiology of the human axilla and its relationship to axillary odor. *J Invest Dermatol* 77:413–416.
- Persson S, Edlund MB, Claesson R, Carlsson J (1990) The formation of hydrogen sulfide and methyl mercaptan by oral bacteria. *Oral Microbiol Immunol* 5:195–201.
- Dickson BJ (2008) Wired for sex: The neurobiology of *Drosophila* mating decisions. *Science* 322:904–909.
- Billeter JC, Atallah J, Krupp JJ, Millar JG, Levine JD (2009) Specialized cells tag sexual and species identity in *Drosophila melanogaster*. *Nature* 461:987–991.
- Benton R (2007) Sensitivity and specificity in *Drosophila* pheromone perception. *Trends Neurosci* 30:512–519.
- Coyne JA (1992) Genetics and speciation. *Nature* 355:511–515.
- Mayr E (1942) *Systematics and the Origin of Species from the Viewpoint of a Zoologist* (Columbia Univ Press, New York).
- Dobzhansky TG (1937) *Genetics and the Origin of Species* (Columbia Univ. Press, New York).
- Schluter D (2009) Evidence for ecological speciation and its alternative. *Science* 323: 737–741.
- Lane DJ (1991) 16S/23S rRNA sequencing. *Nucleic Acid Techniques in Bacterial Systematics*, eds Stackebrandt E, Goodfellow M (Wiley, New York), pp 115–175.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL_X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25:4876–4882.
- Schloss PD, Handelsman J (2005) Introducing DOTUR, a computer program for defining operational taxonomic units and estimating species richness. *Appl Environ Microbiol* 71:1501–1506.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410.
- Zhou WG, Rousset F, O'Neil S (1998) Phylogeny and PCR-based classification of *Wolbachia* strains using *wsp* gene sequences. *Proc Biol Sci* 265:509–515.