

# Genetic variation in human telomerase is associated with telomere length in Ashkenazi centenarians

Gil Atzmon<sup>a,b,1,2</sup>, Miook Cho<sup>a,1</sup>, Richard M. Cawthon<sup>c</sup>, Temuri Budagov<sup>b</sup>, Micol Katz<sup>b</sup>, Xiaoman Yang<sup>b</sup>, Glenn Siegel<sup>b</sup>, Aviv Bergman<sup>d</sup>, Derek M. Huffman<sup>a,b</sup>, Clyde B. Schechter<sup>e</sup>, Woodring E. Wright<sup>f</sup>, Jerry W. Shay<sup>f</sup>, Nir Barzilai<sup>a,b</sup>, Diddahally R. Govindaraju<sup>g</sup>, and Yousin Suh<sup>a,b,2</sup>

<sup>a</sup>Departments of Medicine and Genetics, Albert Einstein College of Medicine, Bronx, NY 10461; <sup>b</sup>Institute for Aging Research, Diabetes Research and Training Center, Albert Einstein College of Medicine, Bronx, NY 10461; <sup>c</sup>Department of Human Genetics, University of Utah, Salt Lake City, UT 84112; <sup>d</sup>Department of Systems and Computational Biology, Albert Einstein College of Medicine, Bronx, NY 10461; <sup>e</sup>Department of Family and Social Medicine, Albert Einstein College of Medicine, Bronx, NY 10461; <sup>f</sup>Department of Cell Biology, University of Texas Southwestern Medical Center, Dallas, TX 75390; and <sup>g</sup>Department of Neurology, Boston University School of Medicine, Boston, MA 02118

Edited by Stephen Curtis Stearns, Yale University, New Haven, CT, and accepted by the Editorial Board October 7, 2009 (received for review July 20, 2009)

**Telomere length in humans is emerging as a biomarker of aging because its shortening is associated with aging-related diseases and early mortality. However, genetic mechanisms responsible for these associations are not known. Here, in a cohort of Ashkenazi Jewish centenarians, their offspring, and offspring-matched controls, we studied the inheritance and maintenance of telomere length and variations in two major genes associated with telomerase enzyme activity, *hTERT* and *hTERC*. We demonstrated that centenarians and their offspring maintain longer telomeres compared with controls with advancing age and that longer telomeres are associated with protection from age-related diseases, better cognitive function, and lipid profiles of healthy aging. Sequence analysis of *hTERT* and *hTERC* showed overrepresentation of synonymous and intronic mutations among centenarians relative to controls. Moreover, we identified a common *hTERT* haplotype that is associated with both exceptional longevity and longer telomere length. Thus, variations in human telomerase gene that are associated with better maintenance of telomere length may confer healthy aging and exceptional longevity in humans.**

longevity | heritability | aging | biomarker

Telomeres consist of the TTAGGG tandem repeats at the ends of chromosomes and are known to protect these regions from degradation and DNA repair activities (1). Telomeres progressively shorten with each cell division in cultured primary human cells (2) until a critically shortened length is achieved, upon which the cells enter replicative senescence (3). Although the relevance of replicative senescence to in vivo aging remains poorly understood, numerous reports suggest that telomere shortening may be associated with organismal aging, with concomitant metabolic decline and increased risk for disease and death (4, 5). For example, several cross-sectional studies in humans have shown that telomere length in white blood cells is inversely related to the age of the cell donor (6–9). Likewise, shorter telomere length has been shown to be associated with age-related disease including coronary heart disease, hypertension, and dementia, as well as general risk factors for disease such as insulin resistance and obesity (10). Furthermore, oxidative stress and inflammation, two major postulated causal factors of aging, are known to accelerate telomere shortening, suggesting that telomere length may be an important biomarker of aging because it reflects the cumulative burden of oxidative stress and inflammation (4, 11). In addition, most (5, 12, 13) but not all studies (14, 15) have shown a positive association between telomere length and overall survival in humans. These results indicate that telomere shortening could be used as a biomarker of disease risk and progression as well as early mortality. However, biological mechanisms responsible for these associations are not known.

Telomere length varies among individuals and families and follows the polygenic mode of inheritance pattern typical of most

quantitative traits (16). Heritability estimates for telomere length vary from 35 to 80% (9, 13, 17). Although several candidate genes have been identified as potential modulators of telomere length in humans (17, 18), none of these genes seem to play a direct role in maintenance of telomere length (19). Recently, one of the most obvious candidate genes of telomere maintenance, telomerase, has been shown to play a direct role in the maintenance of telomere length in humans (20). Telomerase is a specialized ribonucleoprotein enzyme complex that adds telomere repeats to the ends of chromosomes and has two essential components: a catalytic component encoded by the human telomerase reverse transcriptase (*hTERT*) and a human telomerase RNA component (*hTERC*). The latter component provides the template for nucleotide addition by *hTERT*. Heterozygote mutations in the *hTERT* and *hTERC* genes lead to short telomeres and are the major risk factors for rare hematopoietic disorders of bone marrow failure, including aplastic anemia and dyskeratosis congenital. These results indicate that the levels of functional telomerase are critical for telomere maintenance (21). Telomerase is expressed at high levels in specific germline cells, proliferating stem-like cells, and many cancers, whereas in normal adult cell types, it is either not expressed or is expressed at very low levels that are not sufficient to maintain telomere length. However, telomerase can be unregulated in these cells under certain conditions to maintain telomere length (22). This suggests that efficient regulation of telomerase gene expression in response to stresses that are known to reduce telomere length such as oxidative damage or inflammation would lead to better telomere maintenance.

We have previously shown that individuals in Ashkenazi families with exceptional longevity have generally been spared major age-related diseases such as cardiovascular disease and diabetes mellitus, which are largely responsible for mortality in the elderly, and that these features are heritable (23). Because studies on individuals with a normal life span suggest that

This paper results from the Arthur M. Sackler Colloquium of the National Academy of Sciences, "Evolution in Health and Medicine" held April 2–3, 2009, at the National Academy of Sciences in Washington, DC. The complete program and audio files of most presentations are available on the NAS web site at [www.nasonline.org/Sackler\\_Evolution\\_Health\\_Medicine](http://www.nasonline.org/Sackler_Evolution_Health_Medicine).

Author contributions: G.A., R.M.C., N.B., and Y.S. designed research; G.A., M.C., R.M.C., T.B., X.Y., G.S., D.M.H., and Y.S. performed research; G.A., M.C., M.K., A.B., D.M.H., C.B.S., W.E.W., J.W.S., and Y.S. analyzed data; and G.A., M.C., M.K., A.B., J.W.S., N.B., D.R.G., and Y.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. S.C.S. is a guest editor invited by the Editorial Board.

<sup>1</sup>G.A. and M.C. contributed equally to this work.

<sup>2</sup>To whom correspondence may be addressed. E-mail: [gil.atzmon@einstein.yu.edu](mailto:gil.atzmon@einstein.yu.edu) or [yousin.suh@einstein.yu.edu](mailto:yousin.suh@einstein.yu.edu).


This article contains supporting information online at [www.pnas.org/cgi/content/full/0906191106/DCSupplemental](http://www.pnas.org/cgi/content/full/0906191106/DCSupplemental).

common age-related diseases and a shorter life span are associated with shorter telomeres (24–28), we assessed the telomere length in blood leukocytes among subjects with exceptional longevity (centenarians) to investigate if centenarians survived with long telomeres, an indicator of better telomere length maintenance, or short telomeres reflecting chronologically old age. In addition, in view of a strong family history of exceptional longevity and genetic basis of telomere length maintenance, we tested if telomere length as a trait would be inherited by the offspring of centenarians and associated with the status of aging-related diseases. Finally, to study the genetic basis of telomere length maintenance, we took a candidate gene approach to discover all possible variations in two main genes associated with telomerase enzyme activity, *hTERT* and *hTERC*, in centenarians and younger controls for genetic association analysis with human exceptional longevity and telomere length. Our overall hypothesis is that maintenance of telomere length may indicate generalized genomic integrity, which, in turn, may have a profound influence on health and aging in humans.

## Results

**Telomere Length, Heritability, and Longevity.** Phenotypes of centenarians, their offspring (approximate age of 70 years), and offspring-matched controls without a family history of unusual longevity, all of Ashkenazi Jewish descent, are presented in Table 1. The control and offspring groups have similar characteristics of lipid profiles, except that controls have a slightly smaller percentage of large LDL particle size, but these differences are not significant (Table 1). The centenarian group displays a lower mean body mass index and higher HDL levels as compared with controls (Table 1).

When controls and centenarians are combined to represent cross-sectional age groups of unrelated individuals from middle age to oldest old (range: 43–105 years), telomere length declines with age until the age of 85 years (Fig. 1). However, individuals older than 86 years of age as well as the centenarians show longer telomere lengths compared with unrelated individuals younger than 85 years of age, with a mean difference of 0.17 ( $P = 0.04$ ) in the linear regression model adjusting for age and gender. The telomere length of offspring exceeds that of controls in a hierarchical linear model, on average by 0.082 [95% confidence interval (CI): 0.017–0.148;  $P = 0.014$ ]. Because centenarians and their offspring are appreciable singleton families, we used the regression approach for nonindependencies of the observed subjects. To account for parametric model bias as well as for unknown



**Fig. 1.** Comparison of telomere length among Ashkenazi Jewish centenarians ( $n = 86$ ), their offspring ( $n = 175$ ), and controls ( $n = 93$ ). Values are adjusted for age at recruitment and gender in the offspring and control groups and for gender alone in the centenarian group. \* $P < 0.05$ .

intrafamily correlation, we used a sandwich variance estimator equation and a generalized estimating equation (GEE), respectively. Both analyses repeated our first estimation supporting the observation of significant telomere length differences between offspring and controls, on average, by 0.081 (95% CI: 0.022–0.139;  $P = 0.007$ ) for the sandwich estimation and by 0.082 (95% CI: 0.017–0.147;  $P = 0.013$ ) for the GEE. Moreover, unlike controls, offspring of centenarians do not show an appreciable decline in telomere length with age as compared with controls (Fig. 1). However, the difference in age-related decline did not differ significantly between offspring and controls (adjusted mean difference = 0.002, 95% CI: –0.007 to +0.010;  $P = 0.682$ ) by the same hierarchical linear model described earlier. High heritability (86%;  $P = 0.005$ ) for telomere length between centenarians and their offspring was observed, suggesting that telomere length has a strong genetic component in families with exceptional longevity

**Table 1. Phenotypic characteristics of study subjects for telomere length measurements**

	Controls ( $n = 93$ )	Offspring ( $n = 175$ )	Centenarians ( $n = 86$ )	P values	
				Offspring vs. controls	Centenarians vs. controls
Age range, years	43–94 (71.8)	44–85 (67.8)	95–105 (97.4)	0.0001	0.0001
Female, %	55	36	63	0.001	0.24
Body mass index, kg/m <sup>2</sup>	25 (0.4)	26 (0.3)	22 (0.4)	0.28	0.0001
Waist circumference, inch	34 (0.4)	35 (0.3)	35 (0.5)	0.22	0.57
Cholesterol, mg/dL	197 (3.8)	202 (2.9)	191 (4)	0.28	0.26
Triglyceride, mg/dL	154 (8.9)	146 (6.7)	151 (9.6)	0.52	0.81
LDL, mg/dL	107 (3.5)	112 (2.6)	106 (3.6)	0.22	0.76
HDL, mg/dL	58 (1.7)	61 (1.3)	51 (1.8)	0.21	0.003
LDL particle size, nm	20.9 (0.12)	21.1 (0.08)	21.2 (0.1)	0.29	0.06
HDL particle size, nm	9.23 (0.07)	9.15 (0.05)	9.36 (0.06)	0.37	0.17
Large LDL particle size, % of total	52 (4.4)	62 (3.0)	62 (3.9)	0.07	0.1
Large HDL particle size, % of total	53 (2.3)	50 (1.6)	55 (2.1)	0.26	0.42

The phenotypic characteristics were adjusted for age and gender when controls were compared with offspring and for gender when controls were compared with centenarians. All values are mean (SE).


and that these families have better telomere length maintenance than controls.

**Telomere Length, Age-Associated Diseases, and Lipid Profiles of Healthy Aging.** Association analysis between telomere length and major age-related diseases among centenarians, their offspring, and controls indicated that significantly shorter telomere lengths (adjusted for age, gender, and group) are present in subjects with hypertension ( $P = 0.006$ ), the metabolic syndrome ( $P = 0.03$ ), or diabetes ( $P = 0.03$ ) compared with subjects without these disorders (Fig. 2). Moreover, when we tested the relation between telomere length and cognitive function among centenarians, centenarians with impaired cognitive function [Mini-Mental State Examination (MMSE) score  $\leq 25$ ] have significantly shorter telomeres as compared with centenarians with normal cognitive function after adjustment for age and gender (Fig. 2;  $P = 0.02$ ). We then tested whether telomere length correlated with the lipid profiles, which are known predictors of aging-related diseases such as coronary artery disease and metabolic syndromes (29). These analyses revealed that telomere length is associated with the lipid profiles of healthy aging, showing a positive association with increased particle sizes of LDL ( $P = 0.0001$ ) and HDL ( $P = 0.03$ ), percentage of large particle sizes of LDL ( $P = 0.0002$ ) and HDL ( $P = 0.038$ ), and levels of Apo-A1 ( $P = 0.005$ ) and HDL ( $P = 0.04$ ) (Table 2), whereas there is a negative association with very LDL levels ( $P = 0.008$ ). Total cholesterol, triglycerides, LDL, and Apo-B levels show no significant correlation (Table 2). These results suggest that longer telomeres are positively correlated with the lipid profiles of healthy aging.

**Variation in the Telomerase Genes, Longevity, and Telomere Length.** We performed a comprehensive sequence analysis of *hTERT* and *hTERC* genes to detect all possible genetic variants, including the rare variants that may be enriched in centenarians, throughout the coding exons and exon-intron flanking regions in centenarians ( $n = 100$ ) and controls ( $n = 80$ ) using 2D gene scanning and DNA sequencing (Fig. 3). We found a total of 15

sequence variants in the *hTERT* gene, among which 5 were previously unknown unique variants that were not reported in various SNP databases (Table 3). The locations of these variants are shown in Fig. 4. Nine variants in the *hTERT* gene are in the coding region, including two nonsynonymous variants, 893 G > A (Ala-279 Thr) and 3242 G > A (Ala-1062 Thr), which are rare and found only in controls (Table 3). In silico analysis using SIFT (sorts intolerant from tolerant substitutions) (30) and PolyPhen (31), which predict the effects of amino acid changes on protein function, indicates that these changes are likely to be benign. Interestingly, rare synonymous or intronic variants in *hTERT* are enriched in centenarians ( $n = 19$ ) compared with controls ( $n = 3$ ) (Table 2;  $P = 0.041$ ). In addition, three previously undescribed intronic variants in *hTERC* were also found, two of which were rare and found only in centenarians (Table 3).

We genotyped the four common *hTERT* variants, with a minor allele frequency (MAF) of >5% [IVS1-187 T > C, 973 G > A (Ala-305 Ala), 3097 C > T (His-1013 His), IVS16+99 C > T], in 73 centenarians and 49 controls. Information on telomere length was available for all these individuals. We did not find an association of the IVS1-187 T > C, 973 G > A (Ala-305 Ala), and IVS16+99 C > T with longevity. However, significant associations were detected with the 3097 C > T (His-1013 His) variant: the T allele is significantly enriched in centenarians (21.8%) compared with controls (8.2%;  $P = 0.026$ ) (Table 3), and both CT and TT genotypes are significantly enriched in centenarians compared with controls (26.8% and 8.5% vs. 12.2% and 2%, respectively;  $P = 0.035$ ) (Table 4). In contrast, no significant association between the *hTERC* common variant (IVS+63 T > C) and longevity was found (Table 4). Haplotype analysis [see SI Fig. S1 for a linkage disequilibrium (LD) plot derived from the four common variants] revealed that of the four most common haplotypes, Hap 1 is significantly depleted in centenarians compared with controls (34.9% vs. 53.1%;  $P = 0.0056$ ), whereas Hap 3 is significantly enriched in centenarians compared with controls (13.7% vs. 8.2%;  $P = 0.007$ ) when adjusted for age and gender (Table 5). Interestingly, a rare haplotype in controls, Hap 6, is found significantly more frequently in centenarians (1% vs. 6.8%;  $P = 0.024$ ) (Table 5). Association analysis



**Fig. 2.** Telomere length (adjusted for age, gender, and group) according to the absence (gray bars) or presence (white bars) of various age-related diseases in the study subjects. Number of individuals analyzed in each group: hypertension ( $n = 125$ ), without hypertension ( $n = 142$ ), with the metabolic syndrome ( $n = 59$ ), without the metabolic syndrome ( $n = 286$ ), diabetes ( $n = 34$ ), without diabetes ( $n = 276$ ). The MMSE was performed in centenarians only: MMSE score  $\leq 25$  (gray bars,  $n = 46$ ) or  $>25$  (white bars,  $n = 32$ ). \* $P < 0.05$ .

**Table 2. Telomere length and lipid profiles**

	<i>r</i>	<i>P</i> value
Cholesterol, mg/dL	0.06	0.22
Triglyceride, mg/dL	-0.02	0.64
HDL, mg/dL	0.11	0.04
Large HDL, %	0.15	0.038
HDL size, nm	0.15	0.03
Apo-A, mg/dL	0.16	0.005
LDL, mg/dL	0.026	0.62
Large LDL, %	0.26	0.0002
LDL size, nm	0.29	0.0001
Apo-B, mg/dL	-0.003	0.95
Very LDL, mg/dL	-0.19	0.008

Telomere length was adjusted for group, gender, and age at recruitment.

of common *hTERT* or *hTERC* variants with telomere length indicates no significant association (Table S1). However, of the two haplotypes that are enriched in centenarians compared with controls, Hap 3 shows a positive association with telomere length after adjusting for age and gender ( $P = 0.007$ ), whereas Hap 6 does not reach a significant threshold ( $P = 0.59$ ) because of its low frequencies in the tested population (Table 6). In contrast, a centenarian-depleted haplotype, Hap 1, shows no association with telomere length (Table 5).


### Discussion

Several studies have demonstrated a close relation between telomere length and life span in humans, including long-lived humans (4). However, interpretations of results from these studies are often confounded because of a lack of adequate controls. We employed a unique study design to overcome this shortcoming in a cohort of Ashkenazi Jewish individuals with exceptional longevity (centenarians), their offspring (approximate age of 70 years), and age- and gender-matched controls without a family history of unusual longevity. The use of offspring of individuals with exceptional longevity and their matched controls provides a powerful approach to identify ge-

netically controlled longevity traits. This approach has previously led to the identification of longevity phenotypes such as lipoprotein sizes and the subsequent discovery of corresponding longevity genotypes (32–34). In the present investigation, we employed the same study design to delineate the relation between telomere lengths, longevity, and diseases of aging as well as to gain insights into the potential role(s) of genetic variations in the *hTERT* and *hTERC* genes on these phenotypes.

This study demonstrates that centenarians and their offspring have significantly longer telomeres than unrelated controls and that this trait is strongly heritable. Moreover, we have demonstrated that offspring of centenarians do not show an appreciable age-related decline in telomere length as is observed in our unrelated control population as well as in other cross-sectional studies (5). Because telomere length in younger persons (<75 years of age) is not significantly different between offspring of centenarians and unrelated controls, these results suggest that families with exceptional longevity have superior telomere length maintenance. Interestingly, unrelated controls older than 86 years of age exhibit longer telomeres than younger individuals (controls), suggesting that the rate of telomere attrition may be an important determinant of overall survival in the general population. The relatively older group may be affected by survival bias that selected out individuals with aging-related disease or those who would have died before reaching the age of 85 years, leaving the survivors with relatively longer telomeres.

Because shorter telomere length is associated with diseases of aging, including hypertension, the metabolic syndrome, and dementia (10), we studied the association between telomere length, major age-related diseases, and lipid profiles in centenarians, their offspring, and controls. Lipid profiles are known predictors of age-related diseases, and we have previously demonstrated that centenarians and their offspring have significantly larger HDL and LDL particle sizes and that these are heritable phenotypes of healthy aging associated with a lower prevalence of morbidity. We found that longer telomeres are indeed associated with lower prevalence of hypertension, the metabolic syndrome, type 2 diabetes, and better cognitive function as well as with healthier lipid profiles.



**Fig. 3.** 2D gene scanning of human telomerase gene (*hTERT* and *hTERC*). The coding regions and exon–intron junctions of the *hTERT* and *hTERC* were amplified by 2-step PCR. Eighteen PCR fragments were displayed in a 2D gel according to their size and melting temperature. (A) 2D gene scanning pattern from a centenarian subject with the fragment identification number and a heteroduplex band in exon 14 of the *hTERT*. (B) Common genetic variation in exon 14 was identified as 3097 C > T (His-1013 His) by nucleotide sequencing.


**Table 3. Telomerase gene variants in centenarians and controls**

	Nucleotide change	Protein change	No. of Heterozygotes	
			Centenarians (n = 100)	Controls (n = 80)
<b><i>hTERT</i></b>				
Non-synonymous	835 G > A	Ala 279 Thr	0	1
	3148 G > A	Ala 1050 Thr	0	1
Synonymous	915 G > A	Ala 305 Ala	43	27
	1812 A > G	Ala 604 Ala	1	1
	1849 C > T	Leu 618 Leu	1	1
	<b>2355 C &gt; G</b>	<b>Ser 785 Ser</b>	1	0
	<b>2481 G &gt; A</b>	<b>Thr 827 Thr</b>	3	0
	<b>2739 C &gt; T</b>	<b>His 913 His</b>	4	1
	3097 C > T	His 1013 His	28	11
	Intronic	<b>IVS1-211G &gt; A</b>		2
	IVS1-187 T > C		52	36
	IVS4+10 C > T		4	0
	<b>IVS4+21 C &gt; T</b>		1	0
	IVS16+64 C > T		2	0
	IVS16+99 C > T		30	12
<b><i>hTERC</i></b>				
Intronic	<b>IVS-99 C &gt; G</b>		1	0
	<b>IVS+12 A &gt; G</b>		1	0
	<b>IVS+63 T &gt; C</b>		30	27

Previously unknown unique variants are indicated in bold.

Telomere length, as a quantitative trait, is fairly well studied (19), but genetic factors that influence it are not well understood. Recent studies suggest that heterozygote mutation in *hTERT* and *hTERC* genes, which are the essential components of telomerase, show defective phenotypes in several diseases, indicating that half the usual dose of telomerase is inadequate for maintenance of telomeres with normal length (35, 36). Because living to 100 years of age is a rare phenotype in humans, with a prevalence of 1 in 10,000 individuals in the general population (37), we hypothesized that centenarians may harbor rare gain-of-function mutations in the telomerase genes that may also influ-

ence the length of telomeres. Sequence analysis of *hTERT* and *hTERC* genes revealed that rare synonymous or intronic variants in *hTERT* are enriched in centenarians ( $n = 19$ ) compared with controls ( $n = 3$ ) (Table 2;  $P = 0.041$ ). In contrast, centenarians are completely devoid of nonsynonymous variants, whereas there are two control individuals who carry heterozygote nonsynonymous variants, 893 G > A (Ala-279 Thr) and 3148 G > A (Ala-1050 Thr). Although these nonsynonymous changes are predicted to be functionally benign, they are also found in idiopathic pulmonary fibrosis (IPF) patients with a MAF >5% (38), raising the possibility that these variants may have negative



**Fig. 4.** Schematic representation of exons, introns, UTRs, and variants discovered. (A) *hTERT*. (B) *hTERC*. Green blocks, coding exons; white blocks, 5' and 3' UTRs; \*previously unknown variants.

**Table 4. Association of hTERT common alleles/genotypes with exceptional longevity**

Nature of variation			Controls (n = 49)	Centenarians (n = 73)	P value <sup>†</sup>	
			Allele, %			
TERT	Exon 1	IVS1-187 T > C	C	36.7	42.5	0.3717
	Exon 2	973 G > A (A 305 A)	A	30.6	34.9	0.4834
	Exon 14	3097 C > T (H 1013 H)	T	8.1	21.9	0.0049
	Exon 16	IVS16+99 C > T	T	11.2	16.4	0.2556
TERC		IVS+63 T > C	C	18.5	18.8	0.9402
			Genotype, %			
TERT	Exon 1	IVS1-187 T > C	TT	40.8	31.5	0.4494
			CT	44.9	52.1	
			CC	14.3	16.4	
Exon 2	973 G > A (A 305 A)	GG	53.1	43.8	0.5313	
		GA	32.7	42.5		
		AA	14.3	13.7		
Exon 14	3097 C > T (H 1013 H)	CC	85.7	63.9	0.035	
		CT	12.2	26.8		
		TT	2	8.5		
Exon 16	IVS16+99 C > T	CC	81.6	68.5	0.0972	
		CT	14.3	30.1		
		TT	4.1	1.4		
TERC		IVS+63 T > C	TT	64.4	66.2	0.5653
			CT	34.2	29.9	
			CC	1.4	3.9	

<sup>†</sup>Adjusted for age and gender.

functional effects. In addition, although we did not find a single coding variation in the *hTERC* gene, we found two rare intronic variants only in centenarians. The enrichment of both synonymous and intronic variants of *hTERT* and *hTERC* genes in centenarians is intriguing because these variants are known to play a functional role in the regulation of gene expression through modulation of mRNA stability, mRNA secondary structure, alternative splicing, or translational efficiency (39, 40).

The expression level of hTERT is a major determinant of telomerase activity (41). Recent studies indicate that telomerase activation can be regulated by environmental interventions such as lifestyle changes or stress management (42). There are functional hTERT promoter variants known to influence telomerase expression and telomere length (43). However, because there is very little LD in the *hTERT* gene, it is not likely that the centenarian-enriched rare variants are proxies of functional variants in the *hTERT* promoter. It is tempting to speculate that the rare variants might influence expression of *hTERT* in response to environmental stresses such as inflammation or oxidative stress, which then promotes better maintenance of telomere length. Indeed, mean telomere length is much greater among the carriers of the rare variant compared with the

noncarriers after adjusting for age and gender [telomere repeat copy number to single-copy gene copy number (T/S) ratio of 0.62:0.41, crude difference = 0.21;  $P = 0.02$  by Student *t* test,  $P = 0.037$  by Mann-Whitney *U* test], raising the possibility that these rare variants may have a positive functional impact on telomere maintenance. Our study also revealed that common genetic variations in hTERT may influence the telomere length maintenance associated with longer telomeres in families with exceptional longevity.

Haplotypes of the *hTERT* gene derived from four common variants showed associations with longevity and/or telomere length. These are two intronic variants, IVS1-187 T > C and IVS16+99 C > T, and two synonymous variants, 973 G > A (Ala-305 Ala) and 3097 C > T (His-1013 His). These common variants may contribute to the regulation of *hTERT* gene expression. A common intronic *hTERT* variant shown to be associated with susceptibility to IPF may also affect the expression levels of hTERT (44). Taken together, our study demonstrates that centenarians may harbor individually rare but collectively more common genetic variations in genes involved in the telomere maintenance pathway, implicating a role of regulatory variants in

**Table 5. hTERT haplotypes and association with exceptional longevity**

Hap	IVS1-187 T > C	973 G > A (A305A)	3097 C > T (H1013H)	IVS16+99 C > T	Haplotype frequency, %		P value*
					Centenarians (n = 74)	Controls (n = 49)	
1	T	G	C	C	34.9	53.1	0.008
2	C	A	C	C	21.9	23.5	0.322
3	T	G	C	T	13.7	9.2	0.007
4	C	A	T	C	10.3	5.1	0.392
5	C	G	C	C	6.2	5.1	0.749
6	T	G	T	C	6.8	1.0	0.024
7-9			Others		6.2	3.1	0.203

\*Adjusted for age and gender.



parameter is set to 0 with the model in which the parameter is estimated. In addition, we compared telomere length between offspring and controls using a hierarchical linear regression model to account for clustering within families. The regression equation adjusted for age and gender included an indicator variable distinguishing offspring from controls and an offspring vs. controls  $\times$  age interaction term. We tested the hypotheses that offspring have higher mean telomere length than controls and that the age-related decline in telomere length is less in offspring than in controls. To determine the effect of lipid profile on telomere length, a general linear model adjusted for group, gender, and age at recruitment was used. Correlations of telomere length and lipid profile adjusted for group, gender, and age at recruitment were evaluated using Spearman rank correlation coefficients. Heritability of telomere length was assessed in two ways: First, it was computed in the ASSOC module of SAGE (v6.0.1) (<http://darwin.cwru.edu/sage/>). Second, narrow sense heritability was estimated from the slope of the linear regression of the traits of each parent on the mean value of the offspring group (16). Data are expressed as mean (SE) as appropriate. Statistical analyses were performed using SAS 9.1 (SAS Institute).

Comparison of the observed numbers of each genotype with those expected under Hardy-Weinberg equilibrium was tested using  $\chi^2$  tests for each of the three groups: centenarians, their offspring, and controls. Hap-

lotypes were constructed using the PHASE algorithm (50). The haplotype trend regression analysis for logistical (case-control), linear (telomere length), and proportional hazard with telomere length was performed using JMP genomics 4 (SAS Inc., Cary, NC).

**ACKNOWLEDGMENTS.** We are indebted to all participants and their families for their dedication and enthusiasm to enroll in the longevity study. We are also grateful to the Hebrew Home for the Aging (Riverdale, NY), Kittay House (Bronx, NY), Hebrew Home Hospital (West Hartford, CT), and Jewish Home for the Aged (New Haven, CT), all under the aegis of the association for the Jewish Aging Services (Washington, DC). This work was supported by grants from the Paul Beeson Physician Faculty Scholar in Aging Award, Ellison Medical Foundation Senior Scholar Award, Glenn Award for Research in Biological Mechanisms of Aging, and the Resnick Gerontology Center, and by National Institutes of Health Grants RO1 AG-18728-01A1, RO1 AG024391, PO1 AG027734, PO1 AG17242 and RO1 AG7992, General Clinical Research Center Grant MO1-RR12248, and Diabetes Research and Training Center Grant DK 20541 at the Albert Einstein College of Medicine. Some of the results of this paper were obtained by using the program package S.A.G.E., which is supported by a U.S. Public Health Service Resource Grant (RR03655) from the National Center for Research Resources.

- de Lange T (2005) Shelterin: The protein complex that shapes and safeguards human telomeres. *Genes Dev* 19:2100–2110.
- Wong JM, Collins K (2003) Telomere maintenance and disease. *Lancet* 362:983–988.
- Campisi J, d'Adda di Fagnana F (2007) Cellular senescence: When bad things happen to good cells. *Nat Rev Mol Cell Biol* 8:729–740.
- Finch CE (2007) *The Biology of Aging* (Academic Press, Burlington, MA).
- Njajou OT, et al. (2007) Telomere length is paternally inherited and is associated with parental lifespan. *Proc Natl Acad Sci USA* 104:12135–12139.
- Benetos A, et al. (2001) Telomere length as an indicator of biological aging: The gender effect and relation with pulse pressure and pulse wave velocity. *Hypertension* 37:381–385.
- Lindsey J, McGill NI, Lindsey LA, Green DK, Cooke HJ (1991) In vivo loss of telomeric repeats with age in humans. *Mutat Res* 256:45–48.
- Nawrot TS, Staessen JA, Gardner JP, Aviv A (2004) Telomere length and possible link to X chromosome. *Lancet* 363:507–510.
- Slagboom PE, Droog S, Boomsma DI (1994) Genetic determination of telomere size in humans: a twin study of three age groups. *Am J Hum Genet* 55:876–882.
- Aviv A (2006) Telomeres and human somatic fitness. *J Gerontol A Biol Sci Med Sci* 61: 871–873.
- von Zglinicki T, Martin-Ruiz CM (2005) Telomeres as biomarkers for ageing and age-related diseases. *Curr Mol Med* 5:197–203.
- Cawthon RM, Smith KR, O'Brien E, Sivatchenko A, Kerber RA (2003) Association between telomere length in blood and mortality in people aged 60 years or older. *Lancet* 361:393–395.
- Bischoff C, et al. (2005) The heritability of telomere length among the elderly and oldest-old. *Twin Res Hum Genet* 8:433–439.
- Bischoff C, et al. (2006) No association between telomere length and survival among the elderly and oldest old. *Epidemiology* 17:190–194.
- Martin-Ruiz CM, Gussekloot J, van Heemst D, von Zglinicki T, Westendorp RG (2005) Telomere length in white blood cells is not associated with morbidity or mortality in the oldest old: A population-based study. *Aging Cell* 4:287–290.
- Falconer DS, Mackay TFC (1996) *Introduction to Quantitative Genetics* (Addison Wesley Longman, Essex, UK).
- Vasa-Nicotera M, et al. (2005) Mapping of a major locus that determines telomere length in humans. *Am J Hum Genet* 76:147–151.
- Andrew T, et al. (2006) Mapping genetic loci that determine leukocyte telomere length in a large sample of unselected female sibling pairs. *Am J Hum Genet* 78: 480–486.
- Blasco MA (2007) Telomere length, stem cells and aging. *Nat Chem Biol* 3:640–649.
- Bodnar AG, et al. (1998) Extension of life-span by introduction of telomerase into normal human cells. *Science* 279:349–352.
- Garcia CK, Wright WE, Shay JW (2007) Human diseases of telomerase dysfunction: Insights into tissue aging. *Nucleic Acids Res* 35:7406–7416.
- Shay JW, Wright WE (2007) Hallmarks of telomeres in ageing research. *J Pathol* 211: 114–123.
- Atzmon G, et al. (2004) Clinical phenotype of families with longevity. *J Am Geriatr Soc* 52:274–277.
- Benetos A, et al. (2004) Short telomeres are associated with increased carotid atherosclerosis in hypertensive subjects. *Hypertension* 43:182–185.
- Brouillette SW, et al.; West of Scotland Coronary Prevention Study Group (2007) Telomere length, risk of coronary heart disease, and statin treatment in the West of Scotland Primary Prevention Study: A nested case-control study. *Lancet* 369:107–114.
- Panosian LA, et al. (2003) Telomere shortening in T cells correlates with Alzheimer's disease status. *Neurobiol Aging* 24:77–84.
- Samani NJ, Boulty R, Butler R, Thompson JR, Goodall AH (2001) Telomere shortening in atherosclerosis. *Lancet* 358:472–473.
- von Zglinicki T, et al. (2000) Short telomeres in patients with vascular dementia: An indicator of low antioxidative capacity and a possible risk factor? *Lab Invest* 80: 1739–1747.
- Govindaraju DR, et al. (2008) Genetics of the Framingham Heart Study population. *Adv Genet* 62:33–65.
- Ng PC, Henikoff S (2003) SIFT: Predicting amino acid changes that affect protein function. *Nucleic Acids Res* 31:3812–3814.
- Sunyaev S, et al. (2001) Prediction of deleterious human alleles. *Hum Mol Genet* 10: 591–597.
- Atzmon G, et al. (2002) Plasma HDL levels highly correlate with cognitive function in exceptional longevity. *J Gerontol A Biol Sci Med Sci* 57:M712–M715.
- Barzilai N, Atzmon G, Derby CA, Bauman JM, Lipton RB (2006) A genotype of exceptional longevity is associated with preservation of cognitive function. *Neurology* 67:2170–2175.
- Barzilai N, et al. (2003) Unique lipoprotein phenotype and genotype associated with exceptional longevity. *J Am Med Assoc* 290:2030–2040.
- Artandi SE (2006) Telomeres, telomerase, and human disease. *N Engl J Med* 355: 1195–1197.
- Verma S, Slutsky AS (2007) Idiopathic pulmonary fibrosis—New insights. *N Engl J Med* 356:1370–1372.
- Perls TT, Bochen K, Freeman M, Alpert L, Silver MH (1999) Validity of reported age and centenarian prevalence in New England. *Age Ageing* 28:193–197.
- Tsakiri KD, et al. (2007) Adult-onset pulmonary fibrosis caused by mutations in telomerase. *Proc Natl Acad Sci USA* 104:7552–7557.
- Kimchi-Sarfaty C, et al. (2007) A "silent" polymorphism in the MDR1 gene changes substrate specificity. *Science* 315:525–528.
- Qu HQ, Lawrence SG, Guo F, Majewski J, Polychronakos C (2006) Strand bias in complementary single-nucleotide polymorphisms of transcribed human sequences: Evidence for functional effects of synonymous polymorphisms. *BMC Genomics* 7:213.
- Ducrest AL, Szutorisz H, Lingner J, Nabolz M (2002) Regulation of the human telomerase reverse transcriptase gene. *Oncogene* 21:541–552.
- Ornish D, et al. (2008) Increased telomerase activity and comprehensive lifestyle changes: A pilot study. *Lancet Oncol* 9:1048–1057.
- Matsubara Y, et al. (2006) Telomere length of normal leukocytes is affected by a functional polymorphism of hTERT. *Biochem Biophys Res Commun* 341:128–131.
- Mushiroda T, et al.; Pirfenidone Clinical Study Group (2008) A genome-wide association study identifies an association of a common variant in TERT with susceptibility to idiopathic pulmonary fibrosis. *J Med Genet* 45:654–656.
- Lancaster JM, Carney ME, Futreal PA (1997) BRCA 1 and 2—A genetic link to familial breast and ovarian cancer. *Medscape Womens Health* 2:7.
- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (2001) Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *J Am Med Assoc* 285: 2486–2497.
- Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state." A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 12:189–198.
- Cawthon RM (2002) Telomere measurement by quantitative PCR. *Nucleic Acids Res* 30:e47.
- Vijg J, Suh Y (2005) In *Molecular Diagnostics*, eds Patrinos GP, Ansong W (Elsevier, Amsterdam), pp 95–105.
- Stephens M, Donnelly P (2003) A comparison of bayesian methods for haplotype reconstruction from population genotype data. *Am J Hum Genet* 73:1162–1169.