

rnaset2 mutant zebrafish model familial cystic leukoencephalopathy and reveal a role for RNase T2 in degrading ribosomal RNA

Noémie Haud^a, Firat Kara^b, Simone Diekmann^c, Marco Henneke^c, Jason R. Willer^d, Melissa S. Hillwig^d, Ronald G. Gregg^d, Gustavo C. MacIntosh^e, Jutta Gärtner^{c,1}, A. Alia^{b,1}, and Adam F. L. Hurlstone^{a,1}

^aFaculty of Life Sciences, University of Manchester, Manchester M13 9PT, United Kingdom; ^bLeiden Institute of Chemistry, Leiden University, 2300 RA, Leiden, The Netherlands; ^cDepartment of Pediatrics and Pediatric Neurology, Georg August University, 37075 Göttingen, Germany; ^dDepartment of Biochemistry and Molecular Biology, University of Louisville, Louisville, KY 40202; and ^eDepartment of Biochemistry, Biophysics and Molecular Biology, Iowa State University, Ames, IA 50011

Edited by Melissa J. Moore, Howard Hughes Medical Institute, Worcester, MA, and accepted by the Editorial Board October 28, 2010 (received for review July 7, 2010)

T2-family acidic endoribonucleases are represented in all genomes. A physiological role for RNase T2 has yet to be defined for metazoa. RNASET2 mutation in humans is linked with a leukoencephalopathy that arises in infancy characterized by cortical cysts and multifocal white matter lesions. We now show localization of RNASET2 within lysosomes. Further, we demonstrate that loss of *rnaset2* in mutant zebrafish results in accumulation of undigested rRNA within lysosomes within neurons of the brain. Further, by using high field intensity magnetic resonance microimaging, we reveal white matter lesions in these animals comparable to those observed in RNASET2-deficient infants. This correlates with accumulation of Amyloid precursor protein and astrocytes at sites of neurodegeneration. Thus we conclude that familial cystic leukoencephalopathy is a lysosomal storage disorder in which rRNA is the best candidate for the noxious storage material.

T2 endoribonucleases are distributed throughout the kingdoms of life. They perform diverse biological roles in bacteria, fungi, and plants, including phosphate scavenging following nutritional stress, cleaving tRNA in response to oxidative stress, as well as mediating gametophyte self-incompatibility (reviewed in ref. 1). However, little is known about their functions in metazoa. Recently we uncovered that RNASET2 deficiency in man results in a leukoencephalopathy that arises in infancy characterized by cortical cysts and multifocal white matter lesions (2). Patients manifest with psychomotor impairment, micro- or normocephaly, spasticity, and epilepsy. However, neither the physiological role of RNASET2 nor the disease mechanism was apparent from this clinical presentation. Also, to our knowledge, no animal model has yet been generated to study this disease.

Of all the ribonuclease families, the RNase T2 family is the only one active at acidic pH (1) and consequently, the human RNASET2 was described as a putative lysosomal hydrolase (3, 4). Lysosomes are membrane bound organelles containing 60 or more hydrolytic enzymes active at an acidic pH and responsible for degrading all classes of macromolecules upon fusion of lysosomes with phagosomes and endosomes. Lysosomes play an essential role in autophagy and phagocytosis, being responsible for the degradation of molecules, whole organelles, and microorganisms and subsequent salvaging of metabolites (4). Approximately 50 genetic lysosomal diseases, collectively termed lysosomal storage disorders (LSDs), have now been recognized, which affect one in every 5,000 to 7,000 live births in the United States (5, 6). Inborn lysosomal defects result in disease through the accumulation of nondegraded macromolecules. The severity of disease symptoms depends generally on the nature of the accumulated substrate, giving a unique clinical and pathological diagnostic for each LSD. Generally, multiple organs are affected, leading to severe symptoms and premature death. CNS involvement is very common, with neuronal degeneration being apparent (7). Although the

deficient gene and the nature of the accumulating product are known in the majority of LSDs, the events that lead from macromolecule accumulation to disease are still frequently unclear.

In zebrafish we have previously described two RNase T2 encoding paralogues: *rnaset2l* (ribonuclease T2 like, previously referred to as *RNase Dre1*), which is conserved only in fish, on chromosome 15, with 35% identity at the amino acid level to RNASET2; and *rnaset2* (previously referred to as *RNase Dre2*) on chromosome 13, with 47% identity, that appears to be the orthologue of RNASET2 (8). In this study, we have identified zebrafish mutant in the *rnaset2* gene and shown that these animals accumulate undigested rRNA in lysosomes within neurons of the brain. In adults this leads to neuronal degeneration that can be detected by both magnetic resonance microimaging (μ MRI) and immunohistochemical analysis of Amyloid precursor protein (APP) and astrocyte accumulation. Thus we propose that the familial cystic leukoencephalopathy arising in RNASET2-deficient humans is a manifestation of an LSD in which rRNA is the best candidate for the noxious storage material.

Results and Discussion

Identification of an *rnaset2* Mutation in Zebrafish. In a screen of *N*-ethyl-*N*-nitrosourea (ENU)-mutagenized zebrafish for altered acridine orange (AO) uptake, we identified mutant embryos with increased AO staining (Fig. 1A, compare WT with *AO127* mutant). The gene affected in *AO127* mutants was identified by positional cloning (Fig. S1A). Sequencing of a positional candidate *zgc:113369*, equivalent to *rnaset2*, which we previously described (8), identified a T-to-A transversion present only in mutant samples (T304A mutation; Fig. 1B). *rnaset2* is expressed ubiquitously in zebrafish embryos and adults from conception (8). The mutation within codon 118 of the *rnaset2* ORF creates a premature stop codon truncating the protein before its second conserved catalytic site, CAS II, resulting in loss of essential catalytic residues (Fig. S1B). Moreover, as shown by RT-PCR, this mutation results in a dramatic down-regulation of the *rnaset2* messenger RNA (Fig. 1C), presumably through nonsense-mediated decay (9). Capped mRNA was produced from full-length WT

Author contributions: R.G.G., J.G., A.A., and A.F.L.H. designed research; N.H., F.K., S.D., M.H., J.R.W., M.S.H., G.C.M., and A.F.L.H. performed research; N.H., F.K., S.D., M.H., J.R.W., M.S.H., R.G.G., G.C.M., J.G., A.A., and A.F.L.H. analyzed data; and N.H. and A.F.L.H. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. M.J.M. is a guest editor invited by the Editorial Board.

¹To whom correspondence may be addressed. E-mail: gaertnj@med.uni-goettingen.de, a.alia@chem.leidenuniv.nl, or adam.hurlstone@manchester.ac.uk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1009811107/-DCSupplemental.




Fig. 1. *AO127* mutant animals are deficient in *rnaset2*. (A) *AO127* mutants displayed a higher uptake of AO compared with WT embryos, quantifiable by flow cytometry. (Note the intense fluorescence common to both animals is a result of AO within the lumen of the pharynx and intestine, autofluorescence of residual yolk, and ready uptake of AO by olfactory neurons and hair cells within neuromasts.) (Scale bars: 200 μ m.) (B) Sequencing of the gene *zgc:113369*, coding for the zebrafish orthologue of human RNASET2, identified a T-to-A transversion in the mutant sample. (C) RT-PCR indicates a dramatic down-regulation of *rnaset2* mRNA in *AO127* mutants compared with WT siblings. β -Actin serves as a control for RNA integrity and equal input cDNA. (D) Suppression of excess AO staining in *AO127* mutants by injection of WT *rnaset2* mRNA but not mutant mRNA.

rnaset2 cDNA and injected into one-cell stage *AO127* mutant embryos. Flow cytometry analysis confirmed restoration of normal AO staining (Fig. 1D). In contrast, injection with the mutant version of *rnaset2*, *rnaset2*^{T304A}, did not alter AO staining. Furthermore, knockdown of *rnaset2* in *AO127* heterozygotes using an antisense oligonucleotide induced increased AO staining (Fig. S2). Thus, we have identified a mutant harboring a null allele of *rnaset2*, an orthologue of RNASET2, and confirmed that this mutation is responsible for increased AO binding.

RNASET2 Localizes to Lysosomes. Although certain reports imply that RNase T2 molecules are localized to lysosomes (3, 4), others have suggested that they are secreted enzymes (1). To corroborate a lysosomal localization, immunofluorescence staining was performed on HEK 293 cells transfected with WT RNASET2. This revealed partial overlap in the subcellular distribution of RNASET2 with either a marker for the endoplasmic reticulum (ie, PDI) or Golgi apparatus, and significantly more overlap with a marker of late endosomes and lysosomes (LAMP-1; Fig. 2A). This is consistent with a protein that transits through the endoplasmic reticulum and Golgi apparatus to the lysosomes. In contrast, in cells transfected with RNASET2^{C184R}, harboring a missense mutation as identified from a patient with cystic leukoencephalopathy (2), signal was retained within the endoplasmic reticulum (Fig. 2B), presumably as a result of activation of the misfolded protein response (10).

***rnaset2*-Deficient Zebrafish Display Lysosomes Engorged with rRNA.** The localization of RNASET2 to lysosomes, combined with the increased AO staining observed in *AO127* mutants and the fact that within cells AO preferentially stains lysosomes (11), led us to hypothesize the existence of enlarged or supernumerary lysosomes in our mutants. To address this, we examined sections of zebrafish embryos by using electron microscopy (EM). Indeed, enlarged electron-dense lysosomes were readily apparent in the brains of mutant embryos, whereas they were not apparent in WT siblings (Fig. 3). Further, in situ hybridization using a probe for 28S rRNA (12) revealed rRNA aggregates again specifically in *AO127* mutants (Fig. 3; Fig. S3 shows control sense probe) presumably corresponding to the electron-dense material observed by EM. This suggests that the engorged lysosomes might be ac-

cumulating undigested rRNA. Enlarged lysosomes were not readily detected outside the central nervous system in *AO127* mutant zebrafish embryos. Additionally, knockdown of RNA-




Fig. 2. Subcellular localization of RNASET2. HEK293 cells were transfected with myc-epitope tagged WT RNASET2 (A) and the mutated version, RNASET2 C184R (B). Immunofluorescence for endoplasmic reticulum (PDI), Golgi, and late endosome (LAMP-1) markers show partial colocalization with WT RNASET2, suggesting that RNASET2 enters the secretory pathway. The mutated version colocalized completely with the PDI, suggesting ER retention. (Scale bar: 10 μ m.)




Fig. 3. Brain neurons of *AO127* mutants contain lysosomes engorged with rRNA. Brain transverse sections from embryos 5 d after fertilization. EM revealed abundant electron-dense vesicles in the neurons of *AO127* mutants (some of these are indicated by red arrowheads; *Inset*, electron-dense material is bounded by a limiting membrane). (Scale bars: 5 μ m). In situ hybridization revealed accumulation of rRNA aggregates (white arrows) in the cytoplasm of *AO127* mutant neurons. (Scale bars: 10 μ m.)

SET2 in HEK 293 cells resulted in increased lysosome numbers but not the appearance of enlarged lysosomes (Fig. S4 A–D). Also, we failed to detect accumulation of rRNA in RNASET2-depleted HEK293 cells (Fig. S4E). We assume, therefore, that the increased AO uptake detected in most other cells within *AO127* mutants was caused, as for HEK 293 cells, by increased lysosome numbers. Indeed, increased lysosome biogenesis has previously been documented as a compensatory mechanism when lysosome function is compromised (13).

rnaset2-Deficient Zebrafish Develop White Matter Lesions. As previously described, the brains of RNASET2-deficient humans present a diagnostic pattern of brain abnormalities on MRI (2) (Fig. S5A). T₂-weighted μ MRI (14) was also used to analyze adult zebrafish. White matter lesions adjacent to brain ventricles were observed in all *AO127* mutant animals analyzed ($n = 7$) and ventricles also appeared dilated; frequent focal white matter anomalies were scattered throughout brains (Fig. 4). Analysis of T₂ relaxation time suggested that white matter lesions might be filled with fluid similar to CSF in ventricles (Fig. S5B). Moreover, immunohistochemical staining with an antibody recognizing APP indicated APP accumulation and possible axonal damage (15) specifically in mutant brains, particularly evident in white matter regions bordering ventricles (Fig. 4). Likewise, staining for GFAP revealed an increase of astrocytes around sites of putative damage (Fig. 4).

Notwithstanding the neurodegeneration described, animals homozygous for the *AO127* allele appear morphologically normal, reach a normal adult size and weight, are fertile, and do not display obvious gross motor defects. Clearly, this is in contrast to the debilitating psychomotor impairment observed in RNASET2-deficient humans. Previous RT-PCR and in situ hybridization analyses revealed extensive overlap between *rnaset2l* and *rnaset2* expression in embryonic and adult tissues, with both isoforms being expressed in the central nervous system (8). Potentially, *rnaset2l* can partly compensate for *rnaset2* deficiency. Indeed, by in gel RNase activity assay, we observed only partial reduction in RNase activity in the 20- to 30-kDa range at an acidic pH in protein

extracts from *AO127* mutant embryos compared with extracts from WT animals (Fig. S6A). RNase activity in this size range, although not affected by antisense oligonucleotide-mediated knockdown of *rnaset2l* alone, was completely abolished by depleting *rnaset2l* in an *AO127* mutant background (Fig. S6B). Further, knockdown of *rnaset2l* by itself induced only a modest increase in AO staining in WT fish and failed to augment the increased AO staining observed in *AO127* mutants (Fig. S6C). Thus, although *rnaset2* is not an essential gene product in zebrafish and *rnaset2l* is likely to partly compensate for its deficiency, *AO127* mutants nonetheless recapitulate aspects of the leukoencephalopathy observed in RNASET2-deficient humans, supporting their use as a preclinical animal model of this disease.

In this study, we present a zebrafish model for a childhood-onset leukodystrophy. Furthermore, we demonstrate that familial cystic leukoencephalopathy is most likely an LSD in which rRNA accumulates in lysosomes in brain neurons. We hypothesize that RNase T2 is a factor in the specific ribosome recycling pathway termed “ribophagy,” previously described in yeast and through which ribosomes are targeted selectively to the vacuole to be salvaged (16). Significantly, absence of the RNase T2 encoded by *rns2* in *Arabidopsis* also results in constitutive accumulation of undigested RNA in vacuoles, and rRNA acquires a longer half-life in *ms2* mutants (17), so this seems to be a conserved housekeeping function for RNase T2 in multicellular organisms. Depletion of RNASET2 from HEK 293 cells did not result in the appearance of engorged lysosomes or accumulation of rRNA, but rather increased lysosome numbers, indicative of lysosomal dysfunction (13). Potentially, excess undigested rRNA could be “bailed” from these cells through exocytosis (18) and digested by secreted RNase A-like enzymes in the extracellular environment. The problem may be less easily resolved in neurons if rRNA levels were significantly higher or exocytosis by lysosomes limited.

Concerning the disease mechanism leading from RNase T2 deficiency to neurodegeneration, then, potentially, impaired ribophagy could compromise ribogenesis, reducing cell functionality/viability. Alternatively, as with other LSDs, macroautophagy might be impaired as an indirect consequence of lysosomal dysfunction, which in turn appears to harm cells (5, 19). Conversely, accumulated RNA may itself be cytotoxic, as mutations within another ribonuclease, RNASEH2—which neither targets rRNA nor is a lysosomal hydrolase—underlie a very similar leukoencephalopathy, Aicardi–Goutieres syndrome (20). Moreover, the resemblance of RNASET2 deficiency and Aicardi–Goutieres syndrome to in utero CMV infection (2, 20) implies that accumulation of certain RNA species may be recognized by the organism as nonself and thereby elicit an inflammatory reaction. This in turn may be a potential cause of the disease symptoms. Indeed, another causative mutation in Aicardi–Goutieres syndrome affects the *SAMHD1* gene, which encodes a regulator of the innate immune system (21). None of the disease mechanisms suggested here are, however, mutually exclusive.

Future Perspectives. *rnaset2*-deficient zebrafish could serve as preclinical animal models for further elucidating disease pathophysiology linking RNA accumulation with axonal degeneration and white matter alterations. The early manifestation of the phenotype in transparent zebrafish embryos as well as easy detection by AO staining suggests that therapeutic targets could be readily validated in this system. To assess disease course as well as the impact of any candidate therapeutics, longitudinal studies in adult fish could be performed using μ MRI on living fish as previously described (14).

Experimental Procedures

Detailed methods are included in [SI Experimental Procedures](#).

a 28S rRNA Cy3 tagged antisense oligodeoxynucleotide probe (Invitrogen; *SI Experimental Procedures*). Images were collected on a Leica TCS SP5 AOB5 inverted confocal using a 40× objective. When acquiring 3D optical stacks, the confocal software was used to determine the optimal number of Z sections. Only the maximum intensity projections of these 3D stacks are shown.

MRI. MRI studies of the human brain were performed using a 3T MRI system (Magnetom Trio; Siemens) with eight-channel phased-array head coils. Appropriate informed consent was obtained from the patients' parents. MRI was performed on adult zebrafish as previously described (14) using a 400 MHz (9.4 T) vertical bore system and a transmit/receive birdcage radiofrequency coil with an inner diameter of 10 mm and a 1 Tm⁻¹ gradient insert from Bruker. *SI Experimental Procedures* provides further details on MRI.

Paraffin Sections and Immunostaining. Paraffin-embedded tissue sections (5 μm) were used for immunohistochemistry (IHC). The primary antibodies used were mouse anti-APP (1:500 dilution; Zymed) and mouse anti-GFAP (1:300

dilution; Dako), incubated overnight at 4 °C. The secondary antibody used was Vectastain Biotinylated horse anti-pan-Ig antibody. Following incubation with the secondary antibody, the sections were incubated with the ABC kit (Vectorlabs) and stained with DAB (Vectorlabs) before being mounted with Pertex. Images were captured using a Zeiss Axioplan microscope and Axiocam MR camera, and processed using Axiovision software.

ACKNOWLEDGMENTS. We are indebted to Barry McColl, Stuart Allan, Paul Shore, Martin Lowe, and Philip Woodman for reagents and advice (all University of Manchester, Manchester, UK). In addition, we thank staff of the University of Manchester Biological Services Unit for animal caretaking, Ann Warhurst in the histology facility, Mike Jackson in the flow cytometry facility, and Samantha Forbes in the Electron Microscopy facility. This work was supported by Grants C11876/A4495 from Cancer Research UK (to A.H.), Centre for Medical Systems Biology (to A.A.); Deutsche Forschungsgemeinschaft Grant GA354/6-1 (to J.G. and M.H.); the Bundesministerium für Bildung und Forschung through the German Leukodystrophy Network (M.H.); and the Research Program, Faculty of Medicine, Georg August University Göttingen (M.H.).

1. Luhtala N, Parker R (2010) T2 Family ribonucleases: Ancient enzymes with diverse roles. *Trends Biochem Sci* 35:253–259.
2. Henneke M, et al. (2009) RNASET2-deficient cystic leukoencephalopathy resembles congenital cytomegalovirus brain infection. *Nat Genet* 41:773–775.
3. Campomenosi P, et al. (2006) Characterization of RNASET2, the first human member of the Rh/T2/S family of glycoproteins. *Arch Biochem Biophys* 449:17–26.
4. Lübke T, Lobel P, Sleat DE (2009) Proteomics of the lysosome. *Biochim Biophys Acta* 1793:625–635.
5. Ballabio A, Gieselmann V (2009) Lysosomal disorders: From storage to cellular damage. *Biochim Biophys Acta* 1793:684–696.
6. Futerman AH, van Meer G (2004) The cell biology of lysosomal storage disorders. *Nat Rev Mol Cell Biol* 5:554–565.
7. Walkley SU (2009) Pathogenic cascades in lysosomal disease—why so complex? *J Inherit Metab Dis* 32:181–189.
8. Hillwig MS, et al. (2009) Zebrafish RNase T2 genes and the evolution of secretory ribonucleases in animals. *BMC Evol Biol* 9:170.
9. Chang YF, Imam JS, Wilkinson MF (2007) The nonsense-mediated decay RNA surveillance pathway. *Annu Rev Biochem* 76:51–74.
10. Anelli T, Sitia R (2008) Protein quality control in the early secretory pathway. *EMBO J* 27:315–327.
11. Zelenin AV (1966) Fluorescence microscopy of lysosomes and related structures in living cells. *Nature* 212:425–426.
12. Rouquette J, Choismel V, Gleizes PE (2005) Nuclear export and cytoplasmic processing of precursors to the 40S ribosomal subunits in mammalian cells. *EMBO J* 24:2862–2872.
13. Sardiello M, et al. (2009) A gene network regulating lysosomal biogenesis and function. *Science (New York)* 325:473–477.
14. Kabli S, Alia A, Spaink HP, Verbeek FJ, De Groot HJ (2006) Magnetic resonance microscopy of the adult zebrafish. *Zebrafish* 3:431–439.
15. Suenaga T, et al. (1994) Bundles of amyloid precursor protein-immunoreactive axons in human cerebrovascular white matter lesions. *Acta Neuropathol* 87:450–455.
16. Kraft C, Deplazes A, Sohrmann M, Peter M (2008) Mature ribosomes are selectively degraded upon starvation by an autophagy pathway requiring the Ubp3p/BreSp ubiquitin protease. *Nat Cell Biol* 10:602–610.
17. Hillwig MS, et al. (2011) RNS2, a conserved member of the RNase T2 family, is necessary for ribosomal RNA decay in plants. *Proc Natl Acad Sci USA*, 10.1073/pnas.1009809108.
18. Luzzio JP, Pryor PR, Bright NA (2007) Lysosomes: Fusion and function. *Nat Rev Mol Cell Biol* 8:622–632.
19. Settembre C, Fraldi A, Rubinsztein DC, Ballabio A (2008) Lysosomal storage diseases as disorders of autophagy. *Autophagy* 4:113–114.
20. Crow YJ, et al. (2006) Mutations in genes encoding ribonuclease H2 subunits cause Aicardi-Goutières syndrome and mimic congenital viral brain infection. *Nat Genet* 38:910–916.
21. Rice GI, et al. (2009) Mutations involved in Aicardi-Goutières syndrome implicate SAMHD1 as regulator of the innate immune response. *Nat Genet* 41:829–832.
22. Westerfield M (2000) *The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish (Danio rerio)* (Univ of Oregon Press, Eugene, OR).
23. van Eeden FJ, Granato M, Odenthal J, Haffter P (1999) Developmental mutant screens in the zebrafish. *Methods Cell Biol* 60:21–41.
24. Bahary N, et al. (2004) The Zon laboratory guide to positional cloning in zebrafish. *Methods Cell Biol* 77:305–329.
25. Michailidou C, et al. (2009) Dissecting the roles of Raf- and PI3K-signalling pathways in melanoma formation and progression in a zebrafish model. *Dis Model Mech* 2:399–411.