

# A seismologically consistent compositional model of Earth's core

James Badro<sup>a,b,1</sup>, Alexander S. Côté<sup>a,c</sup>, and John P. Brodholts<sup>c</sup>

<sup>a</sup>Institut de Physique du Globe de Paris, Sorbonne Paris Cité–Université Paris Diderot, Unité Mixte de Recherche 7154, Centre National de la Recherche Scientifique, 75005 Paris, France; <sup>b</sup>Earth and Planetary Sciences Laboratory, École Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland; and <sup>c</sup>Department of Earth Sciences, University College London, London WC1E 6BT, United Kingdom

Edited by Ho-kwang Mao, Carnegie Institution of Washington, Washington, DC, and approved April 14, 2014 (received for review September 4, 2013)

**Earth's core is less dense than iron, and therefore it must contain "light elements," such as S, Si, O, or C. We use ab initio molecular dynamics to calculate the density and bulk sound velocity in liquid metal alloys at the pressure and temperature conditions of Earth's outer core. We compare the velocity and density for any composition in the (Fe–Ni, C, O, Si, S) system to radial seismological models and find a range of compositional models that fit the seismological data. We find no oxygen-free composition that fits the seismological data, and therefore our results indicate that oxygen is always required in the outer core. An oxygen-rich core is a strong indication of high-pressure and high-temperature conditions of core differentiation in a deep magma ocean with an FeO concentration (oxygen fugacity) higher than that of the present-day mantle.**

mineral physics | first principles | geophysics

From the analysis of iron meteorites and the observation of Earth's moment of inertia, we know that the primary constituent of Earth's core is an iron alloy (1) with Fe/Ni~16 (2, 3). Comparing seismic travel times in the core with experimental shockwave measurements, Birch (1) proposed that the core is lighter than pure iron. Shockwave and static diamond anvil cell (DAC) experiments have further constrained the core's density deficit (with respect to pure iron) to be between 5 and 10% (4). This requires lower atomic weight elements to be present as additional constituents—so-called light elements. Moreover, the density jump at the inner core boundary (ICB) between the solid inner core and liquid outer core is ~4.5% (5), too large to be due to just the solid–liquid phase transition, and indicates that the outer core contains more light elements (~5–10%) than the inner core (~2–3%). The prime light-element candidates for the core, taking into account cosmochemical and petrological constraints, are silicon, sulfur, carbon, oxygen, and hydrogen (6). Models for core composition allow in principio a mixture of several light elements, and many arguments have been put forward over the years for and against each of the elements (2, 7, 8).

Silicon, sulfur, and carbon are rather soluble in iron at all conditions and were originally quite sensibly proposed as the most valid candidates. They are compatible with low-pressure core formation models, either in a shallow magma ocean or in the differentiated accretionary material. The solubility of these elements in molten iron coexisting with silicate melt would be several percent (9), even at low pressures. On the other hand, oxygen solubility is much more limited at low pressures, and DAC experiments show that oxygen can be introduced in the core by reaction with the molten mantle at high pressures and temperatures (10, 11). Oxygen thus became a natural candidate with the introduction of the "deep magma ocean" models (12–15) of core formation. Additional support for oxygen in the core comes from the fact that oxygen is the only light element to be highly incompatible in solid iron; therefore most of the oxygen would be expelled from the growing inner core and remain in the outer core (7, 8), hence elegantly accounting for the problem of the large density contrast between the inner and outer core. Hydrogen is extremely volatile and is thought to have been

brought to Earth during late accretion (16, 17), after the core had formed. In this case, it would be essentially nonexistent in the proto-Earth during core formation and not a likely candidate for the light element in the core.

The literature offers a wide range (3, 2, 7, 8, 10, 11, 18, 19) of plausible estimates for the light-element composition of the core (*SI Appendix, section 1*). To constrain these further, we need to assess whether the compositional model for the core matches the seismically observed density and sound velocity of the core. As the core is 95% molten, this analysis has not been possible due to the lack of density and velocity data on (Fe–Ni)–C–O–Si–S liquid alloys under core conditions. Measuring bulk sound velocities and densities in molten Fe alloys at core conditions lies currently beyond the capability of experimentation. An alternative is to use ab initio simulations to interpret seismic observations (20) in terms of outer core composition. We therefore calculated the density and bulk sound velocity of liquid alloys in the (Fe–Ni)–C–O–Si–S system using ab initio molecular dynamics. We then compared the properties of the molten alloys directly with the primary geophysical observations [e.g., density and bulk sound velocity obtained (21) from radial seismic models]. This allowed us to identify the subset of compositions that match the constraints and, finally, to propose a seismologically constrained compositional model of Earth's core.

The simulations were performed on liquid iron binaries (Fe<sub>1-x</sub>Ni<sub>x</sub>; Fe<sub>1-x</sub>C<sub>x</sub>; Fe<sub>1-x</sub>O<sub>x</sub>; Fe<sub>1-x</sub>Si<sub>x</sub>; Fe<sub>1-x</sub>S<sub>x</sub>) at two different concentrations ( $x = 8.3$  and  $16.7$  mol%) at the pressure and temperature conditions of the core–mantle boundary (CMB) and the ICB (on the outer core side). Details about the simulations can be found in the *SI Appendix, section 2*.

We calculated the densities with a statistical uncertainty (1 $\sigma$ ) of 0.15% and bulk sound velocities with a statistical uncertainty

## Significance

It is well known that Earth's core is made primarily of iron, alloyed with ~5% nickel and some lighter elements, such as carbon, oxygen, silicon, or sulfur. The amount as well as the chemistry of the light elements is poorly known and still a matter of considerable debate. In this paper we calculate the seismic signature of iron-rich light-element alloys and compare them to the seismic properties of Earth's core. We find that oxygen is required as a major light element in the core, whereas silicon, sulfur, and carbon are not required. We also find that silicon concentration in the core cannot be higher than 4.5%, and sulfur concentration cannot be higher than 2.4%.

Author contributions: J.B., A.S.C., and J.P.B. designed research; J.B., A.S.C., and J.P.B. performed research; J.B., A.S.C., and J.P.B. analyzed data; and J.B. and J.P.B. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

<sup>1</sup>To whom correspondence should be addressed. E-mail: badro@ipgp.fr.


This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1316708111/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1316708111/-DCSupplemental).

(1 $\sigma$ ) of 0.8%. These are reported in Fig. 1. We combined the binary data assuming ideal mixing to obtain the density and bulk sound speed for any composition in the (Fe–Ni)–C–O–Si–S system as  $\rho = \sum x_i \rho_i$  and  $V_\phi = \sqrt{\frac{K}{\rho}}$  where  $\frac{1}{K} = \sum \frac{x_i}{K_i}$ .  $\rho$  is the density of the mixture,  $K$  its bulk modulus,  $V_\phi$  its bulk sound velocity, and  $x_i$ ,  $\rho_i$ , and  $K_i$  are the volume fraction, density, and bulk modulus of the Fe– $X_i$  component, respectively. Ideal mixing has been the standard working hypothesis in this kind of study (6, 19, 22) and will need to be verified by future work. However, our study reinforces this hypothesis by showing that (i) the binary systems are perfectly ideal (as can be seen by the perfectly linear fits of density versus concentration) and (ii) our calculations were compared with existing shockwave data (19, 22–24) on molten Fe, Fe–O, and Fe–S alloys and found them to be in excellent agreement (SI Appendix, section 3). It should also be noted that high-pressure experiments have shown that miscibility gaps vanish at high pressures (25–28), hence also indicating that high-density liquids tend to have a simpler thermodynamic behavior than their low-pressure counterpart.

We calculated  $\rho_{\text{CMB}}$ ,  $\rho_{\text{ICB}}$ ,  $V_{\phi,\text{CMB}}$ , and  $V_{\phi,\text{ICB}}$  for various outer core compositional models in the literature, derived from both experimental and theoretical models (2, 7, 8, 10, 11). These are reported in Fig. 1, alongside the binary data. Except for the ab initio model of Alfè et al. (7), all of the models overestimate the concentration of light elements, yielding densities that are too low. The velocities for the various core compositions are generally higher than observed at the CMB, another indication that the light-element concentration was overestimated.

Assuming a chemically homogeneous outer core, we can constrain its composition by finding all possible combinations of light-element concentrations for which their densities and velocities match those of the Preliminary Reference Earth Model simultaneously at the CMB and ICB. The Fe/Ni ratio in chondrites shows very little variance, so we fix Fe/Ni at 16 (2, 3). The CMB temperature is fixed at  $T_{\text{CMB}} = 4,300$  K so that the ICB temperature (calculated along the isentrope) is  $T_{\text{ICB}} = 6,300$  K (SI Appendix, section 2), which is consistent with iron melting at the ICB (6, 29). The results for other temperature profiles are also tested. We generated over 100 million combinations of ( $x_{\text{O}}$ ,  $x_{\text{Si}}$ ,  $x_{\text{S}}$ ,  $x_{\text{C}}$ ), never exceeding a threshold of 25 mol% for any single light element, and calculated their densities and bulk sound velocities. We kept the compositions that satisfy the four seismological constraints ( $\rho_{\text{CMB}}$ ,  $\rho_{\text{ICB}}$ ,  $V_{\phi,\text{CMB}}$ ,  $V_{\phi,\text{ICB}}$ ) while propagating all uncertainties (0.15% on calculated densities, 0.5% on seismic densities, 0.8% on calculated velocities, and 0.2% on seismic velocities) in our multicomponent model to obtain a seismologically constrained core compositions.

The first striking observation is that all of our solutions contain oxygen, and there are no solutions in an oxygen-free system. Second, there is a valid core composition with oxygen being the only light element ( $5.4 \pm 0.4\%$ ) [all percentages are in weight (wt%) except where otherwise noted], alloyed with Fe–Ni. No other element is able to satisfy the constraints alone. Finally, the maximum concentrations permissible for silicon and sulfur concentrations are rather low, 4.5 and 2.4%, respectively. To visualize the complex solution space, we first plotted the ternary solution spaces: (Fe–Ni)–O–Si, (Fe–Ni)–O–C, and (Fe–Ni)–O–S in



**Fig. 1.** Density (Left) and bulk sound velocity (Right) of molten Fe–Ni, Fe–C, Fe–O, Fe–Si, and Fe–S alloys as a function of concentration at CMB (Upper) and ICB (Lower) conditions. The calculations are represented by full symbols, and the lines are fits to the data (density, linear; bulk sound velocity, quadratic). Note that the densities of C and O at the ICB overlap and are indistinguishable. The horizontal dashed line represents the seismological “target value,” and the shaded area represents its uncertainty. The half-filled circular points are the calculated density and bulk sound velocity for various core compositional models proposed in the literature—black (2), red (10), blue (11), green (8), purple, Si from ref. 7; light blue, S from ref. 7.



