

Hominids adapted to metabolize ethanol long before human-directed fermentation

Matthew A. Carrigan^{a,b,1}, Oleg Uryasev^b, Carole B. Frye^b, Blair L. Eckman^b, Candace R. Myers^c, Thomas D. Hurley^c, and Steven A. Benner^b

^aDepartment of Natural Sciences, Santa Fe College, Gainesville, FL 32606; ^bFoundation for Applied Molecular Evolution, Gainesville, FL 32604; and ^cDepartment of Biochemistry and Molecular Biology, Indiana University School of Medicine, Indianapolis, IN 46202

Edited by Robert Dudley, University of California, Berkeley, CA, and accepted by the Editorial Board October 28, 2014 (received for review March 4, 2014)

Paleogenetics is an emerging field that resurrects ancestral proteins from now-extinct organisms to test, in the laboratory, models of protein function based on natural history and Darwinian evolution. Here, we resurrect digestive alcohol dehydrogenases (ADH4) from our primate ancestors to explore the history of primate–ethanol interactions. The evolving catalytic properties of these resurrected enzymes show that our ape ancestors gained a digestive dehydrogenase enzyme capable of metabolizing ethanol near the time that they began using the forest floor, about 10 million y ago. The ADH4 enzyme in our more ancient and arboreal ancestors did not efficiently oxidize ethanol. This change suggests that exposure to dietary sources of ethanol increased in hominids during the early stages of our adaptation to a terrestrial lifestyle. Because fruit collected from the forest floor is expected to contain higher concentrations of fermenting yeast and ethanol than similar fruits hanging on trees, this transition may also be the first time our ancestors were exposed to (and adapted to) substantial amounts of dietary ethanol.

experimental paleogenetics | alcohol dehydrogenase | ethanol | primates | evolution

One trend in modern medicine attributes diseases in humans to an incomplete adaptation of the human genome to new challenges presented by our changing cultural and demographic environment (1). This attribution is especially convincing for some “lifestyle” diseases. For example, the recent increase in sugar consumption (including sucrose and fructose) is associated with the emergence of obesity, diabetes, and hypertension (2). Under an evolutionary paradigm, an organism fully adapted to a sugar-rich diet would not be expected to become diseased by consuming sugars, suggesting that humankind has not had enough time to adapt to a modern diet rich in such sugars.

It is unclear whether the human genome has had more time to adapt to dietary ethanol (“alcohol” in the vernacular), which also produces a disease spectrum (“alcoholism”) common today in many societies (3). In one historical model, ethanol was not a significant part of the hominin “Paleolithic diet” (4) and was also absent from the diets of earlier ancestors. Rather, the model holds that ethanol entered our diets in significant amounts only after humans began to store surplus food (possibly because of the advent of agriculture) and subsequently developed the ability to intentionally direct the fermentation of food (~9,000 y ago (5), perhaps as a means of preservation (6). In this model, alcoholism as a disease reflects insufficient time since humans first encountered ethanol for their genome to have adapted completely to ethanol. As such, the allelic variants of enzymes in the ethanol metabolic pathway that disfavor ethanol consumption (e.g., *ADH1B*47His* and *ALDH2*487Lys*, both of which lead to an accumulation of acetaldehyde—a toxic intermediate that causes headache, nausea, and general discomfort) represent an early stage of adaptation, possibly in association with pathogenic infections (7–10).

In an alternative model, primates may have ingested ethanol via frugivory as early as 80 million y ago (Ma), a time corresponding to the origin and diversification of primates (11) and

when angiosperm plants first produced fleshy fruits that can become infected by yeast capable of the accumulating ethanol via fermentation (12). In one version of this model, small amounts of ethanol present in slightly fermenting fruit attached to trees attracted arboreal primates foraging in the trees. In this version, our contemporary attraction to ethanol is an “evolutionary hang-over” that ceased to be beneficial once that attraction became redirected to beverages with high concentrations of ethanol (13), made possible only after humans developed the tools allowing them to intentionally direct fermentation (and enhanced with the advent of technology to distill ethanol to higher concentrations). Another version of the “ethanol early” model for ethanol exposure recognizes that ethanol itself, as well as the food naturally containing it, can be a significant source of nutrition. This model posits that any organism with metabolic adaptations that permit the exploitation of ethanolic food would have access to a specialized niche or important fallback foods unavailable to organisms without this metabolic capacity.

Paleogenetics is an emerging field designed to address such natural historical hypotheses and, in particular, to distinguish between competing historical models (14). Here, to gain a genetic perspective on the natural history of the interaction between our human ancestors and ethanol, we examined the evolution of Class IV alcohol dehydrogenases (ADH4) (see *SI Text* for a discussion of the various synonyms used within the ADH family). These digestive enzymes are abundant in the stomach, esophagus, and tongue of primates and are active against a wide range of alcohols. Thus, ADH4 is the first alcohol-metabolizing enzyme to

Significance

Many modern human diseases are attributed to incompatibility between our current environment and the environment for which our genome is adapted. It is unclear whether this model applies to alcoholism. We investigated this possibility by studying alcohol dehydrogenase class IV (ADH4), the first enzyme exposed to ethanol in the digestive tract that is capable of metabolizing ethanol. We resurrected ancestral ADH4 enzymes from various points in the ~70 million y of primate evolution and identified a single mutation occurring ~10 million y ago that endowed our ancestors with a markedly enhanced ability to metabolize ethanol. This change occurred approximately when our ancestors adopted a terrestrial lifestyle and may have been advantageous to primates living where highly fermented fruit is more likely.

Author contributions: M.A.C. and S.A.B. designed research; M.A.C., O.U., C.B.F., and B.L.E. performed research; C.R.M. and T.D.H. contributed new reagents/analytic tools; M.A.C., O.U., and C.B.F. analyzed data; and M.A.C. and S.A.B. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. R.D. is a guest editor invited by the Editorial Board.

Data deposition: The sequences reported in this paper have been deposited in the GenBank database (accession nos. [KM972566](https://doi.org/10.1093/ncbi/kmu972566)–[KM972576](https://doi.org/10.1093/ncbi/kmu972576)).

See Commentary on page 308.

¹To whom correspondence should be addressed. Email: matthew.carrigan@sfcollge.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1404167111/-DCSupplemental.

encounter ethanol that is imbibed (15), and several studies indicate that ADH4 contributes significantly to the first-pass metabolism of ethanol in humans (16).

ADH4 is also active against retinol (in vitro), and ADH4's high catalytic efficiency for retinol (as defined by its k_{cat}/K_M ratio) suggests it may play a role in retinoic acid biosynthesis (17). Mice with inactivated ADH4 genes, however, display few complications associated with retinoid metabolism except under extreme conditions of dietary retinol excess or dietary retinoid deficiency (18, 19). Further, dietary retinoids occur in the form of retinyl esters (from animal foods) or carotenoids (from plant foods); these forms of provitamin A are not substrates for ADH4 present in the upper gastrointestinal track and are converted into retinol only after entering the small intestines. Geraniol, however, is a monoterpenoid that is structurally similar to retinol and is commonly found in plants as an antifungal, making it a physiologically relevant substrate for ADH4 in herbivorous primates.

We therefore tested alternative models for the history of primate exposure to ethanol by comparing the enzymatic efficiencies of modern and ancestral ADH4 enzymes toward geraniol and ethanol. These comparisons identified a dramatic evolutionary transition from an ethanol-inactive ADH4 to an ethanol-active ADH4 in our hominin ancestors ~10 million y ago.

This study focuses on the evolution of one component of ethanol metabolism, ADH4. Ethanol metabolism is complex and involves other ethanol-metabolizing enzymes [e.g., ADH1, ADH2, and the microsomal ethanol oxidizing system (MEOS)], enzymes involved in the downstream metabolism of by-products from ethanol metabolism (e.g., ALDH2, which oxidizes acetaldehyde created from ethanol), and enzymes indirectly affected by the by-products of ethanol metabolism (e.g., ALDH1). A more nuanced understanding of primate adaptation to ethanol will develop as future work examines these related enzymes.

Results

A maximum-likelihood analysis (20) was used to infer the sequences of nine ancestral ADH4 proteins spanning the ~70-million-y history of the primate Order (Fig. 1). These ancestral proteins were synthesized and purified, and their kinetic properties were examined. The evolution of the behaviors of these ancestral enzymes is striking. Nearly all digestive ADH4 enzymes from our primate ancestors were largely inactive against ethanol (Table 1). They did, however, oxidize other alcohols (Table S1), including terpenoid alcohols (such as geraniol) (Table 1) that are abundant in the leaves of plants (21). These results suggest that the last common ancestor of humans and orangutans (node 53 in Fig. 1) was unable to efficiently metabolize dietary ethanol




Fig. 1. The reference phylogeny of the alcohol dehydrogenase 4 (ADH Class IV, ADH4) sequences used in this study. Amino acid changes in the evolution of ADH4 proteins are shown along tree branches. Ancestral ADH4 proteins examined in this study are indicated by numbered nodes within the tree. Ambiguities at nodes are indicated within gray boxes; for example, the sequence at node 44 contained one ambiguity at site 117 (the posterior probabilities for this site were 86% for valine and 14% for isoleucine). All ambiguous versions of the ancestral proteins were resurrected and found to have similar kinetic properties. Blue and red text below branches indicate the V117I and A294V homoplasies shared by the aye aye and Homininae lineages. Sequences indicated by * were, to our knowledge, determined for the first time in the research described here. Sections of gene sequences that are missing in the genomic data are indicated by the corresponding amino acid numbers in italics (except in the case of *T. belangeri*, where the missing region, indicated in bold, was substituted with the corresponding region of the closely related *T. glis*). Some of the polymorphisms found in humans and chimpanzees are indicated on the tree below each species' name. Branches of the tree in red indicate enzymes active against ethanol. The branch leading to chimpanzee is pink to indicate that the R38H polymorphism is frequent among *Pan troglodytes* (based on the limited sequences in current genomic databases), and the impact of this polymorphism is not known.

(at least as part of its first-pass metabolism) but was able to metabolize dietary geraniol and other “long-chain” alcohols. This primate population lived ~13–21 million y ago (based on various estimates of divergence times) (11, 22, 23).

However, this situation changed dramatically in the time after the divergence of orangutans from the lineage leading to humans, chimpanzees, and gorillas (HCGs). In the last common ancestor of humans, chimpanzees, and gorillas (HCG ancestor, node 54 in Fig. 1), an ADH4 emerged that was able to oxidize ethanol 40-fold better than the enzyme at node 53, as measured by its k_{cat}/K_M ratio (this difference is statistically significant, $P = 6 \times 10^{-5}$). Fig. 2 shows the impact of this change on net flux at physiological concentrations of ethanol and geraniol. The dramatic increase in catalytic activity was specific to small molecules like ethanol; activity toward large, physiologically relevant substrates such as vanillyl alcohol decreased twofold and increased less than twofold for cinnamyl alcohol, coniferyl alcohol, and anisyl alcohol. The HCG ancestral ADH4 represented by node 54 is also the predominant form of ADH4 in extant humans and gorillas, and one of two common polymorphs present in extant chimpanzees.

This dramatic increase in catalytic activity arises from a single amino acid replacement at position 294, an impact consistent with protein engineering experiments (24). The increase is due primarily to a decrease in the Michaelis constant for ethanol ($K_{M(ethanol)}$), which drops to 43 mM (± 5.8 mM) (Table 1) in the HCG ancestral ADH4 (node 54) from values greater than 1,000 mM in all earlier ancestors. For comparison, the concentration of ethanol in the blood of a legally intoxicated human is ~10–20 mM, and the concentration of ethanol in fermenting fruits ranges from 85 to 750 mM (25). The changing kinetic properties suggests that the HCG ancestral ADH4 (but not its immediate ancestor, node 53 ADH4) was optimized to metabolize ethanol in fermenting food.

A sample of 13 ancestral and modern ADH4 proteins from outside the Homininae lineage identified only one other ADH4 with increased activity toward ethanol (Table S2). This ethanol-active ADH4 was found in the aye aye (a lemur that diverged from humans ~70 Ma) and possesses the same A294V transition that increased activity in the HCG ancestor (node 54). Our survey of extant primates also identified an ADH4 in a vervet from the Caribbean Island of St. Kitts, *Chlorocebus aethiops sabaues*, with greatly reduced activity toward both ethanol and geraniol (discussed in *SI Text*).

In 2008, the Ensembl database reported nine polymorphisms of the predominant human ADH4 (which is the same as the HCG ancestral ADH4 represented by node 54). This variation

Table 1. Kinetic parameters of various ADH4 enzymes with either ethanol or geraniol substrate (with SDs)

Node	Ethanol		Geraniol	
	K_M , mM	k_{cat}/K_M , $mM^{-1} \cdot min^{-1}$	K_M , μM	k_{cat}/K_M , $\mu M^{-1} \cdot min^{-1}$
54	43 \pm 5.8*	32 \pm 6.2*	17 \pm 3.5	126.1 \pm 8.7
53	3,570 \pm 1,208	0.87 \pm 0.17	160 \pm 23	55.9 \pm 13.7
52	4,822 \pm 5,754	0.45 \pm 0.05	256 \pm 31	42.3 \pm 20.0
49	3,912 \pm 1,876	0.57 \pm 0.14	148 \pm 46	56.7 \pm 19.4
45	3,200 \pm 1,058	0.69 \pm 0.04	183 \pm 20	53.1 \pm 18.7
44 [†]	7,811 \pm 4,387	0.45 \pm 0.06	205 \pm 77	35.8 \pm 5.3

Note that the units for geraniol differ from ethanol by 1,000-fold.

*Comparable to values reported in the literature [e.g., $K_M = 37$ mM; $k_{cat}/K_M = 32$ $mM^{-1} \cdot min^{-1}$ (76)].

[†]An alternative sequence representing node 44 with an isoleucine at position 117 (instead of valine, see Fig. 1) showed similar kinetic properties, with $k_{cat}/K_M \sim 0.87$ and 72 for ethanol and geraniol, respectively. Large SDs are observed in kinetic parameters obtained with ethanol in these enzymes in large part because the K_M values are on the order of several molar (e.g., the K_M of 7.8 M ethanol for node 44 corresponds to ~45% ethanol, a concentration at which many proteins precipitate).




Fig. 2. Substrate oxidation rates by ancestral ADH4 enzymes in assays with ethanol (50 mM; 37.5 ng/mL protein) (Left) or geraniol (0.25 mM; 6.25 ng/mL protein) (Right). Dashed lines are from multiple independent tests. Colors of the lines correspond to colors of the nodes in Fig. 1.

within the human population might produce important differences in alcohol processing, and thus we examined the three polymorphisms closest to the enzyme active site (H52R, R120H, and K226E). Only one of these polymorphisms, H52R, differed catalytically from the predominant form. The catalytic efficiency of the H52R polymorph was reduced approximately sixfold toward ethanol and geraniol (as judged by k_{cat}/K_M), primarily as a result of a lower k_{cat} (Table S2). Amino acid 52 is near the NAD^+ cofactor binding site, and the H52R polymorphism likely alters the binding and/or release of NAD^+ . The frequency of the H52R polymorphism in the human population is not reported in the Ensembl database but is presumably very low.

Discussion

ADH4 Evolution in the Hominoid Lineage. Why did a dietary ADH4 able to metabolize ethanol emerge in our last common ancestor with the chimpanzee and gorilla, an ancestor that lived between 7 and 21 million y before humans developed fermentative technology (11, 22, 23)? Of course, it is always possible that the novel kinetic properties of this ancestral ADH4 served no function in the last common ancestor of humans, chimpanzees, and gorillas. In this case, the ability of node 54 ADH4 to efficiently oxidize ethanol was a “preadaptation” whose utility became important for fitness only after humans developed the process and tools for directing fermentation (6) (see analogous cases of preadaptation in refs. 26 and 27).

However, it is also possible that the novel ADH4 became fixed within the ancestral population in which it arose because it was adaptive. If so, the fixation of an ethanol-active ADH4 would indicate that ethanol became an increasingly important part of the hominid diet after orangutans diverged from the lineage leading to modern human, but before gorillas and chimpanzees diverged. This adaptive model might be preferred if paleontological and paleoclimatic evidence indicate that the ethanol-active ADH4 first emerged in our ancestors during a time when our ancestors experienced selective pressures that increased exposure to dietary ethanol. The available evidence, outlined in the following paragraphs, supports this adaptive model.

The appearance of an ethanol-active ADH4 in the ancestors of HCG (along the branch connecting node 53 and node 54) (Fig. 1) occurred at approximately the same time as a major climatic shift known as the middle Miocene climatic transition (MMCT, ~16 Ma) (28). This rapid environmental change brought about many large-scale ecological transitions, including replacement of Miocene forest ecosystems of East Africa with

fragmented forests and grassland ecosystems (29–32), and coincided with a wave of extinctions (33, 34). Among other global extinctions, hominoid fossils once abundant in the early Miocene forests of East Africa become rare in the middle and late Miocene grassland ecosystems (31, 35, 36).

The major ecosystem changes and mass extinctions seen after the middle Miocene climatic transition suggest changing selective pressures that are associated with adaptations to novel ecological niches (37), including increased terrestriality in the HCG ancestor. Our hominoid ancestors living in the early Miocene and Oligocene (before the MMCT), including the last common ancestor shared with cercopithecoids, were almost certainly arboreal frugivores (38–40). Early Miocene (17–22 Ma) hominoids, such as *Proconsul* (41), and several hominoids of the middle Miocene, including *Nacholapithecus*, were also arboreal frugivores (42–45). Although modern humans, chimpanzees, and gorillas have maintained a frugivorous diet, they have since evolved terrestrial locomotor adaptations and (at least occasionally) consume food collected from the ground (46, 47). Indicators of knuckle-walking in the fossils of *Kenyanthropus* provide evidence of semiterrestriality in hominoids as early as 14–16 Ma (48–50). Fossil evidence suggests that terrestrial adaptations were present in early hominids [e.g., *Orrorin*, 6 Ma (51, 52), and *Sahelanthropus*, 7 Ma (53, 54)], close to the estimated divergence of the human–chimp/gorilla ancestor. Thus, the appearance of an ethanol-active ADH4 in the HCG ancestor occurred at approximately the same time these ancestors were adapting to terrestrial life (presumably) in response to ecological changes brought about by the MMCT.

Saps, nectar, and fruit (in situ) also ferment naturally, thereby exposing both terrestrial and arboreal animals to ethanol (see Wiens et al. for two species of treeshrew that consume significant levels of ethanol from the fermented nectar of bertam palms) (55). Overripe fruit that has fallen to the ground is generally older (and more damaged) than ripe fruit picked directly from trees and, as such, has had more time to ferment, potentially leading to higher concentrations of ethanol (ref. 25, and reviewed in ref. 56). Therefore, the transition to an increasingly terrestrial life would likely have exposed the HCG ancestor to fruit with higher ethanol content. In this context, the increased activity of ethanol-metabolizing enzymes (e.g., ADH4) could provide a selective advantage, particularly during a time of large-scale ecological transitions and extinctions brought about by climate change.

Not all terrestrial organisms are expected to consume ethanolic food. For example, some cercopithecoids independently adapted to a terrestrial life after the hominoid–cercopithecoid split (traits that persist in extant lineages such as *Papio* and *Erythrocebus*), but, unlike the hominoid lineages that remained highly frugivorous, cercopithecoids developed a generalist diet including leaves with high amounts of antifeedants (such as the monoterpene geraniol). The cercopithecoid transition to a generalist, herbaceous diet provided these monkeys a less geographically patchy and seasonally fluctuating diet.

There is a critical distinction between the adaptive strategies adopted by hominoids and cercopithecoids, best exemplified by studies of fallback foods among chimpanzees and three extant cercopithecines of the Kibale Forest in Uganda (57). When fruit was available, chimpanzee diet was confined almost exclusively to ripe fruit; when ripe fruit was not available, chimpanzees relied on alternative terrestrial food sources, such as piths and stems that have low antifeedant levels. The three cercopithecines, however, maintained a diverse diet, including leaves even when ripe fruit was abundant. As a consequence, cercopithecines consumed much higher levels of antifeedants (e.g., geraniol). When ripe fruit was not available, the diet of these cercopithecines consisted primarily of unripe fruits, seeds, and leaves (and thus still contained high antifeedant levels). It is possible that competition between terrestrial middle Miocene monkeys and apes may have been the precursor for the digestive differences between these two groups that persist today, with monkeys

adapted to exploit unripe fruit and African apes adapted to exploit over-ripe ones.

Although ripe fruit is generally the preferred food of all hominoids (58–61), those without an ethanol-oxidizing ADH4 (orangutans and gibbons) differ from chimpanzees and gorillas in several important regards. Unlike chimpanzees and gorillas, gibbons and orangutans rely primarily on arboreal fallback foods, and, whereas gibbons and orangutans have unique strategies to buffer against ripe-fruit scarcity, these strategies bear important similarities to the cercopithecines. During times of food scarcity, gibbons increase consumption of unripe fruit, as well as figs, leaves, and flowers (61, 62). Orangutans make greater use of lower quality food regardless of food availability (63), relying heavily and equally on both ripe and unripe fruit even during periods of high food availability (61), and then increasing dietary diversity during food shortages (primarily by increasing consumption of figs, bark, pith, and leaves) (61, 63, 64). Orangutans also store large amounts of energy as fat and therefore can take advantage of periods of fruit superabundance to overcome energy deficits during periods of fruit scarcity (63, 65). The gibbons and orangutans extensive utilization of unripe fruits (the former primarily during times of food shortage) exposes them to higher levels of terpenoid antifeedants relative to ripe and overripe fruit (66). The need to metabolize large terpenoid antifeedants may preclude ADH4 from simultaneously adapting to small substrates like ethanol. If orangutans or gibbons were to exploit fermented fruit, it might require the recruitment of an enzyme not already involved in terpenoid metabolism.

Having adapted to a more terrestrial lifestyle than its predecessors, it is possible that the ancestor of HCG had less access to the fallback food strategies used by arboreal gibbons or orangutans, likely leading to greater selective pressures to exploit alternative fallback food strategies. Our study indicates that the HCG ancestor possessed the capacity for a novel strategy: an ethanol-active digestive ADH4 enzyme that would permit exploitation of food containing increased concentrations of ethanol. Furthermore, because frugivorous vertebrates generally avoid “rotting” (and presumably more ethanolic) fruit when other options are available (reviewed in ref. 6), there may have been little competition for this resource.

If the A294V mutation endowing ADH4 with enhanced ethanol-oxidizing abilities was adaptive in the HCG ancestor, the exploitation of ethanolic food resources might still persist in modern gorillas or chimpanzees under certain conditions, such as food scarcity. Despite numerous anecdotal accounts of primates consuming ethanol-laden food, we are aware of no published accounts that describe such events in detail (much less quantify the amount of ethanol consumed or discern whether the consumption was inadvertent or intentional). However, a report by Ohashi (67) briefly documents wild chimpanzees in Bossou, Guinea consuming significant amounts of ethanol (repeatedly and apparently intentionally) from fermenting palm wine collected from reservoirs placed in the trees by humans. These findings suggest that ethanol consumption, albeit rare, may nonetheless be a significant part of the natural history of modern chimpanzees and support our conclusion that ethanol consumption was important to the common ancestor of humans, chimpanzees, and gorillas.

Primate Adaptation to Ethanol Beyond Hominoid ADH4. Thirteen extant and ancestral ADH4 enzymes outside the Homininae lineage were also examined and only one—from the aye aye (an arboreal lemur)—had increased activity toward ethanol (Table S2). The absence of an ethanol-active ADH4 enzyme in cercopithecoids is expected, given their generalist diet (including fallback foods) containing high levels of antifeedants (from grasses, leaves, and/or unripe fruit) (57) rather than overripe and possibly fermented fruit collected from the ground. Further, some of the most terrestrial cercopithecoids, *Erythrocebus*, *Theropithecus*, and *Papio*, live in semiarid savannahs and grasslands where ethanol-containing fruit is rare.

Although ADH4 is involved in the first-pass metabolism of ethanol, other enzymes also contribute to ethanol metabolism (e.g., ADH1 and ADH2, homologs of ADH4 that are abundantly expressed in the liver, and the MEOS). It is therefore possible that species without an ethanol-active ADH4 are adapted to consume significant amounts of ethanol, as seems to be the case for the common treeshrew (55); in such cases, we expect to see adaptations in other metabolic pathways that mitigate the effects of ethanol consumption.

Remarkably, the only primate in our sample with an ethanol-active ADH4 similar to the HCG ancestor, the aye aye, shares the A294V transition that accounts for the increased activity in the HCG ancestor [aye ayes also share a second homoplasy (V117I) with the lineage leading to the HCG ancestor; see *SI Text* for a discussion of its possible significance]. The branch leading to the aye aye is poorly articulated in the phylogeny used to reconstruct ancestral ADH4 sequences (Fig. 1), and more ADH4 sequences must be added to this model to determine when in the past ~45 million y the A294V mutation occurred along the lineage leading to the modern aye aye. Given this uncertainty about the timing of the A294V mutation in the aye aye lineage and limited information about the diet and life history of aye ayes, it is difficult to speculate whether the independently evolved ethanol-active ADH4 in aye aye is also an adaptation to dietary ethanol. However, aye ayes are known to consume arboreal fruit, sap, and nectar (68) (all of which ferment naturally), so it is reasonable to surmise that aye ayes may have consumed dietary ethanol at some point in their natural history. If so, ethanol consumption may persist in extant aye ayes.

Human ADH4 Variation. The Ensembl database contained nine human ADH4 missense polymorphisms at the beginning of this study (ca. 2008), but the number of ADH4 polymorphisms reported has since grown to over 50, including several relatively frequent variants. Given the likelihood that human ADH4 is adapted to ethanol, and the observation that one of the three polymorphisms we tested possessed significantly altered activity toward ethanol (*Results*), it is possible that some of the many uncharacterized ADH4 polymorphisms may play an important role in the variability of human ethanol metabolism and risk for developing alcoholism or ethanol-related cancers of the upper gastrointestinal tract.

Summary

Ancestral reconstructions of ADH4 demonstrate that the ancestor of humans, chimpanzees, and gorillas possessed a novel enzyme with dramatically increased activity toward ethanol, and we suspect that this novel metabolic capacity was adaptive to this hominin ancestor. This transition implies that the genomes of modern human, chimpanzee, and gorilla began adapting at least 10 million y ago to dietary ethanol present in fermenting fruit—a source of ethanol that is remarkably similar in concentration and form (i.e., with food) to the moderate ethanol consumption now recognized to be healthy for many humans (69). This conclusion contrasts with the relatively short amount of time (~9,000 y) (5) since fermentative technology enabled humans to consume beverages (devoid of food bulk) with higher ethanol content than fruit fermenting in the wild. Vestiges of this history may persist in the extant descendants of the HCG ancestor. This history has implications, not only for understanding the forces that shaped hominin terrestrial adaptations, but also for

understanding the medical complexities of human interactions with ethanol today.

Methods

The evolutionary history of the ADH4 family was reconstructed using ADH4 genes from 28 different mammals, including 17 primates, collected from public databases [National Center for Biotechnology Information (NCBI) and Ensembl] or were generated de novo by RT-PCR of mRNA or genomic DNA extracted from well-preserved tissue samples (Table S3). The genomic sequence data for *Tupaia belangeri* in the Ensembl database were lacking exon 6; this gap was filled using sequence data from a closely related species, *Tupaia glis*. The last exon, encoding just eight amino acids, was missing from a few primates (Table S4). Because this exon is invariant among primates, the conserved sequence was used when the databases did not include data for this exon.

In the acquisition of tissues, ethical standards for the treatment of animals were ensured by the institutions providing samples according to each institution's ethical review board. In general, tissues were collected at necropsy after natural death or were collected initially for research other than that described here.

Genetic material was extracted from these tissues using TRI reagent with minor modifications to the manufacturer's protocol (Sigma). Recovered mRNA was reverse-transcribed, gel-purified, cloned into the TOPO-TA cloning vector (Invitrogen), and then sequenced using Big Dye technology using the service provided by BioBasic.

All protein sequences used in this study are displayed in FASTA format in Table S4. These sequences were placed on a tree that represents the standard phylogeny of primate species (Fig. 1). The reference phylogeny used in this study was compiled from various published phylogenies. For nonprimate outgroups, the reference phylogeny was based on the phylogeny of Prasad et al. (70). The phylogeny of Prasad et al. did not include treeshrew, so our reference phylogeny placed treeshrew sister to primates (relative to rodents) according to the phylogeny of Liu et al. (71). The placement of lemurs in our reference phylogeny was based on the phylogeny of Horvath et al. (72). Our reference phylogeny for nonlemur primates was based on the phylogeny of Purvis (73).

Ancestral ADH4 protein sequences were inferred from the extant species sequence data and phylogeny using a maximum-likelihood analysis implemented within the PAML software package (20) by applying a codon model and a discrete gamma distribution for rate variation among sites. When the posterior probability that a particular amino acid occupied a particular site was $\geq 94\%$, we assigned that amino acid at that site in the ancestral ADH4. When the posterior probability was $< 94\%$ at a particular position (as was the case for amino acid 171 at node 44), both forms of the protein were synthesized and examined to determine whether the interpretation of the kinetic results was robust with respect to the ambiguity.

Synthetic genes encoding various ancestral and extant ADH4 proteins were cloned into the pET21 vector (BioBasic) and expressed in the *Escherichia coli* TUNER cell line. The heterologously expressed proteins were isolated following the procedures of Niederhut et al. (74). The activity and kinetic parameters were characterized by measuring the formation of NADH using various alcohols as substrates as in Yin et al. (75), with the exception that BSA was included in the assays at 0.2 mg/mL. The kinetic values we obtained for human ADH4 were similar to those reported in the literature by others (76). Statistical significance was determined using a modification of a two-tailed Student *t* test that does not assume equal variances between the two populations being tested (also known as Welch's *t* test).

ACKNOWLEDGMENTS. We acknowledge the collaboration of the organizations that provided the tissue that made this work possible (Table S3). M.A.C. thanks Brenda Benefit and three anonymous reviewers whose helpful and thoughtful critiques improved the manuscript quality and clarity. This work was supported by National Institute on Alcohol Abuse and Alcoholism Grant R01AA017723 and National Aeronautics and Space Administration Grant NNX08A023G. This is Duke Lemur Center publication no. 1277.

- Nesse RM, Stearns SC, Omern GS (2006) Medicine needs evolution. *Science* 311(5764):1071.
- Lustig RH, Schmidt LA, Brindis CD (2012) Public health: The toxic truth about sugar. *Nature* 482(7383):27–29.
- Heilig M, Egli M (2006) Pharmacological treatment of alcohol dependence: Target symptoms and target mechanisms. *Pharmacol Ther* 111(3):855–876.
- O'Keefe JH, Jr, Cordain L (2004) Cardiovascular disease resulting from a diet and lifestyle at odds with our Paleolithic genome: How to become a 21st-century hunter-gatherer. *Mayo Clin Proc* 79(1):101–108.
- McGovern PE, et al. (2004) Fermented beverages of pre- and proto-historic China. *Proc Natl Acad Sci USA* 101(51):17593–17598.
- Levey DJ (2004) The evolutionary ecology of ethanol production and alcoholism. *Integr Comp Biol* 44(4):284–289.
- Goldman D, Enoch MA (1990) Genetic epidemiology of ethanol metabolic enzymes: a role for selection. *World Rev Nutr Diet* 63:143–160.
- Luo HR, et al. (2009) Origin and dispersal of atypical aldehyde dehydrogenase ALDH2487Lys. *Gene* 435(1–2):96–103.
- Lin YP, Cheng TJ (2002) Why can't Chinese Han drink alcohol? Hepatitis B virus infection and the evolution of acetaldehyde dehydrogenase deficiency. *Med Hypotheses* 59(2):204–207.
- Peng Y, et al. (2010) The ADH1B Arg47His polymorphism in east Asian populations and expansion of rice domestication in history. *BMC Evol Biol* 10:15.

11. Steiper ME, Young NM (2006) Primate molecular divergence dates. *Mol Phylogenet Evol* 41(2):384–394.
12. Thomson JM, et al. (2005) Resurrecting extinct proteins from ancient yeast at the origin of fermentation. *Nat Genet* 37(6):630–635.
13. Dudley R (2002) Fermenting fruit and the historical ecology of ethanol ingestion: Is alcoholism in modern humans an evolutionary hangover? *Addiction* 97(4):381–388.
14. Benner SA (2007) The early days of paleogenetics: Connecting molecules to the planet. *Experimental Paleogenetics*, ed Liberles DA (Academic, New York), pp 3–19.
15. Vaglenova J, et al. (2003) Expression, localization and potential physiological significance of alcohol dehydrogenase in the gastrointestinal tract. *Eur J Biochem* 270(12):2652–2662.
16. Haber PS, et al. (1996) Metabolism of alcohol by human gastric cells: Relation to first-pass metabolism. *Gastroenterology* 111(4):863–870.
17. Parés X, Farrés J, Kedishvili N, Duester G (2008) Medium- and short-chain dehydrogenase/reductase gene and protein families: Medium-chain and short-chain dehydrogenases/reductases in retinoid metabolism. *Cell Mol Life Sci* 65(24):3936–3949.
18. Deltour L, Foglio MH, Duester G (1999) Impaired retinol utilization in Adh4 alcohol dehydrogenase mutant mice. *Dev Genet* 25(1):1–10.
19. Deltour L, Foglio MH, Duester G (1999) Metabolic deficiencies in alcohol dehydrogenase Adh1, Adh3, and Adh4 null mutant mice: Overlapping roles of Adh1 and Adh4 in ethanol clearance and metabolism of retinol to retinoic acid. *J Biol Chem* 274(24):16796–16801.
20. Yang Z (1997) PAML: A program package for phylogenetic analysis by maximum likelihood. *Comput Appl Biosci* 13(5):555–556.
21. Chen W, Viljoen AM (2010) Geraniol: A review of a commercially important fragrance material. *S Afr J Bot* 76:643–651.
22. Schrago CG, Menezes AN, Moreira MA, Pissinatti A, Seuánez HN (2012) Chronology of deep nodes in the neotropical primate phylogeny: Insights from mitochondrial genomes. *PLoS ONE* 7(12):e51699.
23. Prado-Martinez J, et al. (2013) Great ape genetic diversity and population history. *Nature* 499(7459):471–475.
24. Crossas B, et al. (2000) Molecular basis for differential substrate specificity in class IV alcohol dehydrogenases: A conserved function in retinoid metabolism but not in ethanol oxidation. *J Biol Chem* 275(33):25180–25187.
25. Dudley R (2004) Ethanol, fruit ripening, and the historical origins of human alcoholism in primate frugivory. *Integr Comp Biol* 44(4):315–323.
26. Hartley CJ, et al. (2006) Amplification of DNA from preserved specimens shows blowflies were preadapted for the rapid evolution of insecticide resistance. *Proc Natl Acad Sci USA* 103(23):8757–8762.
27. Bock WJ (1959) Preadaptation and multiple evolutionary pathways. *Evolution* 13:194–211.
28. Flower BP, Kennett JP (1994) The middle Miocene climatic transition: East Antarctic ice sheet development, deep ocean circulation and global carbon cycling. *Palaeogeogr Palaeoclimatol* 108:537–555.
29. Andrews P, Van Couvering JAH (1975) Paleoenvironments in the East African Miocene. *Approaches to Primate Paleobiology*, ed Szalay FS (Karger, Basel).
30. Bonnefille R (1984) Cenozoic vegetation and environments of early hominids in East Africa. *Paleobotany, Paleozoology and Paleoanthropology*. The Evolution of the East Asian Environment (Centre of Asian Studies, University of Hong Kong, Hong Kong), ed Whyte RO, Vol II, pp 579–612.
31. Retallack GJ (1992) Middle Miocene fossil plants from Fort Ternan (Kenya) and evolution of African grasslands. *Paleobiology* 18(4):383–400.
32. Retallack GJ, Wynn JG, Benefit BR, McCrossin ML (2002) Paleosols and paleoenvironments of the middle Miocene, Maboko Formation, Kenya. *J Hum Evol* 42(6):659–703.
33. Pickford M (1981) Preliminary Miocene mammalian biostratigraphy for western Kenya. *J Hum Evol* 10:73–97.
34. Leakey M, Grossman A, Gutiérrez M, Fleagle JG (2011) Faunal change in the Turkana Basin during the late Oligocene and Miocene. *Evol Anthropol* 20(6):238–253.
35. Retallack GJ (1991) *Miocene Paleosols and Ape Habitats in Pakistan and Kenya* (Oxford Univ Press, New York).
36. Fortelius M, Hokkanen A (2001) The trophic context of hominoid occurrence in the later Miocene of western Eurasia: A primate-free view. *Hominoid Evolution and Climatic Change in Europe*, eds de Bonis L, Koufos G, Andrews P (Cambridge Univ Press, Cambridge), pp 19–47.
37. Potts R (2004) Paleoenvironmental basis of cognitive evolution in great apes. *Am J Primatol* 62(3):209–228.
38. Fleagle JG, Simons EL (1982) The humerus of *Aegyptopithecus zeuxis*: A primitive anthropoid. *Am J Phys Anthropol* 59(2):175–193.
39. Ankel-Simons F, Fleagle JG, Chatrath PS (1998) Femoral anatomy of *Aegyptopithecus zeuxis*, an early oligocene anthropoid. *Am J Phys Anthropol* 106(4):413–424.
40. Fleagle JG (1983) Locomotor adaptations of Oligocene and Miocene hominoids and their phyletic implications. *New Interpretations of Ape and Human Ancestry*, eds Ciochon RL, Corruccini RS (Plenum, New York), pp 301–324.
41. Teaford MF, Maas MC, Simons EL (1996) Dental microwear and microstructure in early oligocene primates from the Fayum, Egypt: Implications for diet. *Am J Phys Anthropol* 101(4):527–543.
42. Rose MD (1997) Functional and phylogenetic features of the forelimb in Miocene hominoids. *Function, Phylogeny and Fossils: Miocene Hominoid Evolution and Adaptations*, eds Begun DR, Ward CV, Rose MD (Plenum, New York), pp 79–100.
43. Kay RF (1977) Diets of early Miocene African hominoids. *Nature* 268:628–630.
44. Kay RF, Ungar PS (1997) Dental evidence for diets in some Miocene catarrhines with comments on the effects of phylogeny on the interpretation of adaptations. *Function, Phylogeny and Fossils: Miocene Hominoid Evolution and Adaptations*, eds Begun DR, Ward CV, Rose MD (Plenum, New York), pp 131–151.
45. Nakatsukasa M, Kunimatsu Y (2009) *Nacholapithecus* and its importance for understanding hominoid evolution. *Evol Anthropol* 18:103–119.
46. McNeillage A (2001) Diet and habitat use of two mountain gorilla groups in contrasting habitats in the Virungas. *Mountain Gorillas: Three Decades of Research at Karisoke*, eds Robbins MM, Sicotte P, Stewart KJ (Cambridge Univ Press, Cambridge), pp 265–292.
47. Hockings KJ, Sousa C (2012) Differential exploitation of cashew—a low conflict crop—by sympatric humans and chimpanzees. *Oryx* 46:375–381.
48. McCrossin ML, Benefit BR, Gitau SN, Palmer AK, Blue KT (1998) Fossil evidence for the origins of terrestriality among Old World higher primates. *Primate Locomotion: Recent Advances*, eds Strasser E, Fleagle J, Rosenberger AL, McHenry H (Plenum, New York), pp 353–396.
49. McCrossin ML, Benefit BR (1997) On the relationships and adaptations of *Kenyapithecus*, a large-bodied hominoid from the middle Miocene of eastern Africa. *Function, Phylogeny, and Fossils: Miocene Hominoid Evolution and Adaptation*, eds Begun DR, Ward CV, Rose MD (Plenum, New York), pp 241–267.
50. Sherwood RJ, et al. (2002) Preliminary description of the *Equatorius africanus* partial skeleton (KNM-TH 28860) from Kipsaramon, Tugen Hills, Baringo District, Kenya. *J Hum Evol* 42(1-2):63–73.
51. Senut B, et al. (2001) First hominid from the Miocene (Lukeino Formation, Kenya). *C R Acad Sci Paris, Sciences de la Terre et des planètes/Earth Planet Sci* 332:137–144.
52. Richmond BG, Jungers WL (2008) *Orrorin tugenensis* femoral morphology and the evolution of hominin bipedalism. *Science* 319(5870):1662–1665.
53. Brunet M, et al. (2002) A new hominid from the Upper Miocene of Chad, Central Africa. *Nature* 418(6894):145–151.
54. Zollikofer CPE, et al. (2005) Virtual cranial reconstruction of *Sahelanthropus tchadensis*. *Nature* 434(7034):755–759.
55. Wiens F, et al. (2008) Chronic intake of fermented floral nectar by wild treeshrews. *Proc Natl Acad Sci USA* 105(30):10426–10431.
56. Milton K (2004) Ferment in the family tree: Does a frugivorous dietary heritage influence contemporary patterns of human ethanol use? *Integr Comp Biol* 44(4):304–314.
57. Wrangham RW, Conklin-Brittain NL, Hunt KD (1998) Dietary response of chimpanzees and cercopithecines to seasonal variation in fruit abundance. I. Antifeedants. *Int J Primatol* 19:949–970.
58. Conklin-Brittain NL, Knot CD, Wrangham RW (2001) The feeding ecology of apes. *The Apes: Challenges for the 21st Century* (Chicago Zoological Society, Brookfield), pp 167–174.
59. Doran-Sheehy D, Mongo P, Lodwick J, Conklin-Brittain NL (2009) Male and female western gorilla diet: Preferred foods, use of fallback resources, and implications for ape versus old world monkey foraging strategies. *Am J Phys Anthropol* 140(4):727–738.
60. Knott CD (2005) Energetic responses to food availability in the great apes: Implications for hominin evolution. *Seasonality in Primates*, eds Brockman D, van Schaik CP (Cambridge Univ Press, Cambridge), pp 351–378.
61. Vogel ER, Haag L, Mitra-Setia T, van Schaik CP, Dominy NJ (2009) Foraging and ranging behavior during a fallback episode: *Hylobates albarbaris* and *Pongo pygmaeus wurmbii* compared. *Am J Phys Anthropol* 140(4):716–726.
62. Chivers DJ (2001) The swinging singing apes: Fighting for food and family in Far-East forests. *The Apes: Challenges for the 21st Century* (Chicago Zoological Society, Brookfield), pp 1–28.
63. MacKinnon JR (1977) A comparative ecology of Asian apes. *Primates* 18:747–772.
64. MacKinnon JR (1974) The behavior and ecology of wild orangutans (*Pongo pygmaeus*). *Anim Behav* 22:3–74.
65. Knott CD (1998) Changes in orangutan caloric intake, energy balance, and ketones in response to fluctuating fruit availability. *Int J Primatol* 19(6):1061–1079.
66. Whitehead SR, Bowers MD (2013) Evidence for the adaptive significance of secondary compounds in vertebrate-dispersed fruits. *Am Nat* 182(5):563–577.
67. Ohashi G (2006) Behavioral repertoire of tool use in the wild chimpanzees at Bossou. *Cognitive Development in Chimpanzees*, eds Matsuzawa T, Tomonaga M, Tanaka M (Springer, Tokyo), pp 439–451.
68. Sterling EJ, Dierenfeld ES, Ashbourne CJ, Feistner ATC (1994) Dietary intake, food composition and nutrient intake in wild and captive populations of *Daubentonia madagascariensis*. *Folia Primatol (Basel)* 62(1-3):115–124.
69. Breslow R, et al. (2003) *State of the Science Report on the Effects of Moderate Drinking* (National Institute on Alcohol Abuse and Alcoholism, Bethesda, MD). Available at pubs.niaaa.nih.gov/publications/ModerateDrinking-03.htm.
70. Prasad AB, Allard MW, Green ED; NISC Comparative Sequencing Program (2008) Confirming the phylogeny of mammals by use of large comparative sequence data sets. *Mol Biol Evol* 25(9):1795–1808.
71. Liu F-GR, et al. (2001) Molecular and morphological supertrees for eutherian (placental) mammals. *Science* 291(5509):1786–1789.
72. Horvath JE, et al. (2008) Development and application of a phylogenomic toolkit: Resolving the evolutionary history of Madagascar's lemurs. *Genome Res* 18(3):489–499.
73. Purvis A (1995) A composite estimate of primate phylogeny. *Philos Trans R Soc Lond B Biol Sci* 348(1326):405–421.
74. Niederhut MS, Gibbons BJ, Perez-Miller S, Hurley TD (2001) Three-dimensional structures of the three human class I alcohol dehydrogenases. *Protein Sci* 10(4):697–706.
75. Yin SJ, Bosron WF, Magnes LJ, Li TK (1984) Human liver alcohol dehydrogenase: Purification and kinetic characterization of the beta 2 beta 2, beta 2 beta 1, alpha beta 2, and beta 2 gamma 1 "Oriental" isoenzymes. *Biochemistry* 23(24):5847–5853.
76. Yin SJ, Chou CF, Lai CL, Lee SL, Han CL (2003) Human class IV alcohol dehydrogenase: Kinetic mechanism, functional roles and medical relevance. *Chem Biol Interact* 143-144:219–227.