

*THE CORRELATION OF CYTOLOGICAL AND GENETICAL
CROSSING-OVER IN ZEA MAYS. A CORROBORATION*

BY HARRIET B. CREIGHTON AND BARBARA McCLINTOCK

BOTANY DEPARTMENT, CORNELL UNIVERSITY

Communicated February 9, 1935

There has recently been some skepticism expressed (Brink and Cooper, 1935)¹ as to the value of the studies on the correlation of cytological and genetical crossing-over in maize published by Creighton and McClintock (1931)⁶ because of the fewness of the data. Since the paper by Stern (1931)⁹ dealing with *Drosophila* and having much more extensive data appeared at practically the same time and yielded the same conclusions, the authors felt it unnecessary to add to the ever-increasing amount of published work merely to record more evidence of the same nature without supplying anything essentially new or advancing. Therefore, confirmatory data which have accumulated since the time the joint paper mentioned above was published have not been considered for a separate publication. However, we now feel forced to add more data merely to counteract any suspicion that the evidence previously presented constituted insufficient proof. This will be done in as brief a form as possible, since a discussion of the method has been given in the paper mentioned above.

Chromosome 9 in maize is characterized by its relative size in the chromosome complement and by the 1:2 ratio in lengths of its two arms. The end of the short arm in some strains possesses a large knob while other strains have a very small knob or no knob. Evidence that the knob or knobless condition of a particular chromosome 9 is inherited with the same precision as a gene has been given in the previous paper and has been confirmed in many additional crosses. The knob, therefore, could be used as one cytological marker for this chromosome. The presence of an interchange between chromosomes 8 and 9 (Burnham, 1930,² 1934³; McClintock, 1930⁷) which broke chromosome 9 at a position on the long arm a short distance away from the spindle fibre attachment region provided the second cytological marker. That the genes *yg*, *c*, *sh*, *wx** lie in the interchanged chromosome which possesses the short arm of chromosome 9 has been shown by McClintock, 1931,⁸ Creighton, 1934,⁵ and Burnham 1934.^{3,4} With reference to the knob and the interchange point, the order of the genes is knob-*yg-c-sh-wx*-interchange with *yg* very close to the knob (Creighton, 1934)⁵ and *wx* close to the spindle fibre attachment region (Burnham, 1934⁴ and unpublished). The standard crossover values for these genes alone are *yg-c* 21%, *c-sh* 3.3%, *sh-wx* 21%. The crossover value of *wx* to the interchange is 13.7% (Burnham, 1934³). That there is very little crossing-over between the knob and *yg* can be seen from the data given below.

A plant with the constitution knob-*Yg-C-Sh-Wx*-interchange was crossed to a plant with the constitution knobless-*yg-c-sh-wx*-normal. The F_1 was backcrossed to knobless-*yg-c-sh-wx*-normal. Two hundred and sixty-one individuals resulting from this backcross were examined cytologically to determine the presence or absence of the knob (knob or knobless in table below) and the presence or absence of the interchange (interchange or normal in table below) in the chromosome carrying these genes contributed by the F_1 parent. Since there are five regions in which a crossover can be detected, the results have been tabulated according to crossovers which occurred in each of these regions. The tabulated results do not represent the total backcross progeny. A higher percentage of *Yg-C-Sh-Wx* and *yg-c-sh-wx* plants were examined cytologically in an effort to obtain crossovers between the knob and *yg*. Likewise, more *Yg* plants were examined cytologically than *yg*, since plants homozygous for *yg* are reduced in vigor and often do not afford sufficient material for cytological examination.

TABLE 1

$\frac{\text{KNOB-}Yg-C-Sh-Wx\text{-INTERCHANGE}}{\text{KNOBLESS-}yg-c-sh-wx\text{-NORMAL}} \times \text{KNOBLESS-}yg-c-sh-wx\text{-NORMAL}$		NUMBER OF INDIVIDUALS
Non-crossovers		
1. Knob- <i>Yg-C-Sh-Wx</i> -interchange		84
2. Knobless- <i>yg-c-sh-wx</i> -normal		45
Crossovers in region 1		
3. Knob- <i>yg-c-sh-wx</i> -normal		3
4. Knobless- <i>Yg-C-Sh-Wx</i> -interchange		1
Crossovers in region 2		
5. Knob- <i>Yg-c-sh-wx</i> -normal		13
6. Knobless- <i>yg-C-Sh-Wx</i> -interchange		11
Crossovers in region 3		
7. Knob- <i>Yg-C-sh-wx</i> -normal		3
8. Knobless- <i>yg-c-Sh-Wx</i> -interchange		3
Crossovers in region 4		
9. Knob- <i>Yg-C-Sh-wx</i> -normal		53
10. Knobless- <i>yg-c-sh-Wx</i> -interchange		18
Crossovers in region 5		
11. Knob- <i>Yg-C-Sh-Wx</i> -normal		16
12. Knobless- <i>yg-c-sh-wx</i> -interchange		3
Double crossover involving regions 2 and 4		
13. Knobless- <i>yg-C-Sh-wx</i> -normal		1
Double crossovers involving regions 4 and 5		
14. Knob- <i>Yg-C-Sh-wx</i> -interchange		5
15. Knobless- <i>yg-c-sh-Wx</i> -normal		2

It is obvious from the data given above that a genetic crossing-over between the genes *Yg-C-Sh-Wx* involves a cytological crossover between the

knob and the interchange point. These data, therefore, supplement those given in our previous publication and indicate the soundness of the conclusions drawn.

* The genes referred to in this paper by symbols are: *yg*, yellow-green plants; *c*, colored aleurone; *sh*, shrunken endosperm; *wx*, waxy endosperm.

¹ Brink, R. A., and Cooper, D. C., *Genetics*, 20, 22-35 (1935).

² Burnham, C. R., these PROCEEDINGS, 16, 269-277 (1930).

³ Burnham, C. R., *Genetics*, 19, 430-447 (1934).

⁴ Burnham, C. R., *Am. Nat.*, 68, 81-82 (1934).

⁵ Creighton, H. B., these PROCEEDINGS, 20, 111-115 (1934).

⁶ Creighton, H. B., and McClintock, B., *Ibid.*, 17, 492-497 (1931).

⁷ McClintock, B., *Ibid.*, 16, 791-796 (1930).

⁸ McClintock, B., *Ibid.*, 17, 485-491 (1931).

⁹ Stern, C., *Biol. Zbl.*, 51, 547-587 (1931).

TWO HEREDITARY TYPES OF HYDROCEPHALUS IN THE HOUSE MOUSE (*MUS MUSCULUS*)

BY FRANK H. CLARK

LABORATORY OF VERTEBRATE GENETICS, UNIVERSITY OF MICHIGAN

Communicated February 2, 1935

Different hereditary genes that independently produce a similar phenotypic effect are very common in plants but have been found only rarely in mammals. Castle and Nachtsheim (1933) have described three forms of short-hair (*rex*) in the rabbit, all of which look alike but depend upon different genes for their expression. They have shown that two of these genes are linked or borne in the same chromosome.

In the house mouse (*Mus musculus*), the gene causing blue dilution is very similar in its phenotypic effect to another that dilutes the coat in much the same manner. This second dilution character was described by J. M. Murray (1931) and called *leaden*. Likewise, a new pink-eye dilution in the house mouse described by Roberts (1931) resembles ordinary pink-eye dilution in its effect on eye and coat color.

The purpose of this paper is to report the results of a test designed to show whether or not two types of hydrocephalus ("water on the brain") which occur in the house mouse are due to the same gene or to different genes. One of these forms of hydrocephalus was first noticed in a flexed-tail strain of mice at Michigan State College. It proved to be a simple Mendelian recessive character (Clark, 1932) and an anatomical study (Clark, 1934) showed that it was a hydrocephalus of the obstructive or