

The discussion in Janet, *Leçons sur les Systèmes d'Équations aux Dérivées Partielles*, Paris 1929, pp. 74-75, is particularly relevant. The other references are: C. Riquier, *Les Systèmes d'Équations aux Dérivées Partielles*, Paris, 1910, and J. M. Thomas, *Differential Systems*, New York, 1937.

⁸ See the author's paper "The General Geometry of Paths," *Ann. Math.*, 29, 143-168 (1928).

⁹ For the definition of covariant differentiation, see formula (9.2) of the paper cited in the preceding footnote.

THE GENERALITY OF FINITE ABSTRACT COMPLEXES

BY WILLIAM W. FLEXNER

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, AND INSTITUTE FOR ADVANCED STUDY

Communicated November 10, 1939

1. Let A be a finite abstract complex as defined by S. Lefschetz¹ following A. W. Tucker, W. Mayer and J. W. Alexander. An *open* subcomplex C of an abstract complex D is a subset of D (order, dimensions and incidences in C determined by those in D) such that $x \in C$ and $y > x$ implies $y \in C$. It will be shown here that for every A there is an open subcomplex B of a simplicial complex such that the following homology groups using integer coefficients are isomorphic:

$$H^q(A) \approx H^{q+\alpha}(B), \quad q \text{ arbitrary}, \quad (1)$$

where $\alpha = 0$ if A has no elements of negative dimension and no zero-dimensional torsion coefficients and otherwise $\alpha > 0$. A result of Steenrod's² shows that relation (1) then holds for any coefficient group. One of the principal uses of an abstract complex being to carry a homology theory, the present result shows that in this respect and for the finite case simplicially realizable complexes are as general as any abstract complexes.

2. Let $\| b_{ij} \|$ be the normal form of the $p, p - 1$ incidence matrix of A , and give each row of this matrix a name, E_i^p , and each column a name E_j^{p-1} . If this is done for each row and column of all the simultaneously reduced incidence matrices of A , the set $\{E\}$ may be made an abstract complex C by defining as incidence relations $[E_i^p : E_j^{p-1}] = b_{ij}$. Obviously

$$H^q(A) \approx H^q(C) \quad q \text{ arbitrary}. \quad (2)$$

Because the normal matrices are diagonal and $FF = 0$ in C , $[E_i^p : E_j^{p-1}] \neq 0$ implies that all other incidence relations involving either E_i^p or E_j^{p-1} are zero.

3. Suppose that for some i $[E_i^p : E_i^{p-1}] = k \neq 0, 1, -1$. This cannot happen in a simplicial complex, so to make C simplicial each such pair

E_i^p, E_i^{p-1} must be replaced by a complex D_i^p with the same homology groups and no incidence relations of absolute value greater than 1. The situation described in the last sentence of No. 2 makes it possible to do this without disturbing the rest of C .

I. The following adaptation to abstract complexes of "Subdivision by Section"³ replaces a complex G by a complex G' with the same homology groups: If $E^p \in G$ and $FE^p = C_1^{p-1} + C_2^{p-1}$ where C_i^{p-1} are $(p-1)$ -chains of G , E^p is replaced by e_1^p, e_2^p, e^{p-1} to form G' , incidences being given by

$$\begin{aligned} Fe^{p-1} &= FC_2^{p-1} \\ Fe_1^p &= C_1^{p-1} + e^{p-1} \\ Fe_2^p &= C_2^{p-1} - e^{p-1} \\ [E^{p+1}: e_i^p] &= [E^{p+1}: E^p], i = 1, 2 \end{aligned}$$

and all other incidences (not involving E^p) are the same in G' as in G .

II. Without loss of generality k may be assumed > 1 and the subscripts i may be omitted from the E 's. Using I replace E^p by g_1^p, e_1^p, e_1^{p-1} so that $Fe_1^{p-1} = 0, Fe_1^p = E^{p-1} + e_1^{p-1}, Fg_1^p = (k-1)E^{p-1} - e_1^{p-1}, [E^{p+1}: g_1^p] = [E^{p+1}: e_1^p] = [E^{p+1}: E^p] = 0$. Then unless $k-1 = 1$ replace g_1^p by g_2^p, e_2^p, e_2^{p-1} , etc., until after $n = k-1$ steps

$$\begin{aligned} Fe_i^{p-1} &= 0 \quad 1 \leq i \leq k \\ Fe_i^p &= E^{p-1} - e_{i-1}^{p-1} + e_i^{p-1} \end{aligned} \tag{3}$$

where $e_0^{p-1} = e_k^{p-1} = 0$ and $e_k^p = g_k^p$, all other incidences involving these elements being zero. The situation is now as follows: k new elements e_i^p and $k-1$ new elements e_i^{p-1} have replaced e^p . Examination of formula 3 shows that the incidence $[E^p: E^{p-1}] = k > 1$ has been replaced by several incidences all of absolute value 1; e_i^p and e_{i+1}^p have the two common faces E^{p-1} and e_i^{p-1} ; e_i^{p-1} is oriented to e_i^p oppositely than to e_{i+1}^p , the last two of which circumstances are incompatible with simpliciality and impose the following changes:

III. Subdivide each e_i^p twice more by means of I, finally getting the set $\{s_i^p, s_i^{p-1}, t_i^p, t_i^{p-1}, u_i^p, e_i^{p-1}\} = D^p, i = 1, 2, \dots, k$ with incidences given by

$$Fe_i^{p-1} = Ft_i^p = Fs_i^{p-1} = 0 \tag{4}$$

$$Fu_i^p = -e_{i-1}^{p-1} + e_i^{p-1} + t_i^{p-1} \tag{5}$$

$$Ft_i^p = -t_i^{p-1} - s_i^{p-1} \tag{6}$$

$$Fs_i^{p-1} = E^{p-1} + s_i^{p-1}, \tag{7}$$

all other incidences involving elements of D^p being zero.

IV. Clearly D^p is incidence-equivalent to an open 2-subcomplex K^2 of a simplicial complex where s_i^p, t_i^p, u_i^p are represented by 2-simplexes and e_i^{p-1} ,

$s_i^p - 1, t_i^p - 1$ by suitably chosen 1-simplexes on the boundaries of the 2-simplexes. K^2 has no vertices and some of the 1-simplexes on boundaries of 2-simplexes are missing so it is simplicially open.

V. In order to preserve dimension as far as possible, a technique for raising the dimension of K^2 is needed. If $\alpha \geq 0$ is an integer let $L^{p+\alpha}$ be the join $K^2 \smile \sigma$ where σ is a new open $(p + \alpha - 3)$ -simplex ($\sigma^{-1} = 1$). If now s_i^p is replaced by $s_i^p \smile \sigma$ and similarly for all simplexes of K^2 the incidence formulas 4-7 remain unchanged, so $H^{q+\alpha}(L^{p+\alpha})$ is isomorphic to the q th homology group of the complex $\{E^p, E^{p-1}\}$.

4. The final complex B is obtained as follows:

a. If the minimum dimension, m , of any element of A is ≥ 0 and A has no zero-dimensional torsion coefficient, let $\alpha = 0$. If $m < 0$ and A has no m -dimensional torsion coefficient, let $\alpha = -m$. If $m \leq 0$ and A has an m -dimensional torsion coefficient let $\alpha = -m + 1$.

b. To each D_i^p (corresponding to a pair E_i^p, E_i^{p-1} such as considered in No. 3) assign its $L_i^{p+\alpha}$. To each E_i^p of C with $FE_i^p = 0$ assign an open $(p + \alpha)$ -simplex $W_i^{p+\alpha}$ such that neither W nor its boundary FW meets any previously assigned simplex. The set $B = \{L_i^{p+\alpha}, W_i^{p+\alpha}\}$ is an open complex of a simplicial complex and

$$H^{q+\alpha}(B) \approx H^q(A). \tag{1}$$

To see the latter notice that B has the same structure (except for dimension if $\alpha > 0$) as C with regard to all elements E_i^p of C which do not have the property $FE_i^p = \neq E_i^{p-1}$. These E_i^p are omitted from representation in B because they have no effect on the homology groups of C . Hence by formula 2,

$$H^{q+\alpha}(B) \approx H^q(C) \approx H^q(A)$$

which gives formula 1.

¹ Lefschetz, S., *Bull. Am. Math. Soc.*, **43**, 345-359 (1937). See there for references to other authors.

² Steenrod, N. E., *Amer. Jour. Math.*, **58**, 675 (1936).

³ Lefschetz, S., "Topology," *Am. Math. Soc. Colloquium Publications*, **12**, 68 (1930). Replace proof there by its group-theoretic analogue.