

## Double keystone bird in a keystone species complex

(sapsucker/community/extinction)

GRETCHEN C. DAILY\*, PAUL R. EHRLICH, AND NICK M. HADDAD

Department of Biological Sciences, Stanford University, Stanford, CA 94305

Contributed by Paul R. Ehrlich, September 22, 1992

**ABSTRACT** Species in a Colorado subalpine ecosystem show subtle interdependences. Red-naped sapsuckers play two distinct keystone roles. They excavate nest cavities in fungus-infected aspens that are required as nest sites by two species of swallows, and they drill sap wells into willows that provide abundant nourishment for themselves, hummingbirds, orange-crowned warblers, chipmunks, and an array of other sap robbers. The swallows thus depend on, and the sap robbers benefit from, a keystone species complex comprised of sapsuckers, willows, aspens, and a heartwood fungus. Disappearance of any element of the complex could cause an unanticipated unraveling of the community.

Early attempts to explain the distribution and numbers of species stressed two types of factors: (i) physical features of the environment (climate, topography, insolation, and habitat patch size and degree of isolation) and (ii) availability of other resources (especially food) (1, 2). More recently, a major effort to characterize evident species interactions, such as those between predators and prey or between competitors for common, limiting resources (3, 4) has also led to an appreciation of the complexity and ubiquity of subtle biological associations. Indirect interrelationships may have profound effects on the structure of ecological communities (5-7); understanding their prevalence and influence is crucial to guiding efforts to slow the loss of biodiversity (8).

Here we report the importance of a complex of keystone species to the distribution and abundance of other species. Keystone species are those whose removal from a community would precipitate a further reduction in species diversity or produce other significant changes in community structure and dynamics. Keystone species may act in a variety of ways (9-20): for example, by preying on dominant species, by causing disturbance, by providing resources in times of scarcity, by pollinating and thereby enhancing the persistence of highly dispersed and rare plants, and (as in this case) by providing key, limiting resources or by supporting other keystone species.

The red-naped sapsucker (*Sphyrapicus nuchalis*, a woodpecker; hereafter "sapsucker") appears to play two keystone roles that help to support different suites of species in a subalpine ecosystem in the Rocky Mountains. One role has already been described. The sapsucker drills sap wells into spruce (*Picea engelmannii*), aspen, and shrubby willow (*Salix* spp.). The willow sap constitutes a major portion of the sapsucker's diet during the summer breeding season and is stolen by over 40 species, including hummingbirds, warblers, chipmunks, squirrels, wasps, and butterflies (21, 22). The sapsuckers thus make available a rich resource to numerous species at times when they are reproducing and storing fat for the winter migration and hibernation. In fact, the breeding distribution of the closely related yellow-bellied sapsucker (*S. varius*) appears to determine the northern limits of the

breeding ranges of the Rubythroat and Rufous hummingbirds (23).

The second keystone role is elucidated by the research reported here. The sapsucker provides nest cavities to secondary cavity-nesting species. Each breeding season, a sapsucker pair creates a new nest cavity in an aspen (*Populus tremuloides*) tree infected with the fungus *Fomes ignarius* (which causes heartwood decay and makes cavity excavation possible) (refs. 24 and 25; G.C.D., unpublished data). The old cavities are utilized by at least seven secondary cavity-nesting bird species in Colorado. In 4 years of monitoring nest cavity excavations in the study region (as part of another project), we have found that the sapsucker makes >1 order of magnitude more cavities than other primary cavity nesters (other woodpecker species) and is thus potentially important to secondary cavity nesting species.

We show the significance of this second keystone role to community structure as follows. First we determine that the sapsucker is restricted to habitat with both aspen and willow in close proximity. Since experimental removal of the sapsucker is not desirable, we use aspen groves without willow as treatments (i.e., sapsuckers "removed") and groves with willow as controls (i.e., sapsuckers present) to test for associated differences in bird-community composition. Breeding-bird censuses confirm that the only significant avifaunal difference between sites is that species of swallows that locally are obligate secondary cavity nesters are restricted to groves with sapsuckers; they are therefore dependent upon a keystone complex comprised of certain species of willow, *Fomes*-infected aspen, and sapsuckers.

### MATERIALS AND METHODS

We conducted this investigation in the vicinity of the Rocky Mountain Biological Laboratory in Gunnison County, Colorado, during the 1991 breeding season. Study areas were distributed along the East and Taylor River drainages at elevations ranging between 2650 m and 3050 m.

We located suitable sapsucker habitat by looking for the unique and prominent signs of well excavation. This damage on aspen trees is easily recognizable for at least 10 years after creation, and thus provides an indication of habitat occupied by the sapsucker that is integrated over time.

To determine the relationship between the occurrence of the sapsucker and the proximity of willow and aspen, we surveyed the prevalence of sapsucker damage in aspen groves located at various distances from shrubby willow. Only willow species with sufficiently thick stems to be used by the sapsucker were considered (*S. subcoerulea*, *S. pseudocordata*, and *S. brachycarpa*). We examined a minimum of 400 randomly selected trees in each aspen grove; the numbers of trees with sapsucker wells and with nest cavities were recorded. Sapsucker wells are very distinctive; in contrast, we could not generally attribute a nest cavity to any particular primary cavity-nesting species.

\*To whom reprint requests should be addressed.



Table 1. Results of the breeding bird censuses in the 5.25-hectare study sites

Species	Adults at census site, no.					
	SMN	SMS	WG	AS	LR	DR
<b>Primary cavity nester</b>						
Red-naped sapsucker ( <i>Sphyrapicus nuchalis</i> )	4	2	2	0	0	0
<b>Obligate secondary cavity nesters</b>						
Tree swallow ( <i>Tachycineta bicolor</i> )	2	4	2	0	0	0
Violet-green swallow ( <i>Tachycineta thalassina</i> )	2	4	4	0	0	0
<b>Facultative secondary cavity nesters</b>						
Mountain bluebird ( <i>Sialia currucoides</i> )	*	*	2	*	0	0
Chickadee ( <i>Parus gambeli</i> and <i>P. atricapillus</i> )	2	2	4	2	*	2
Northern flicker ( <i>Colaptes auratus</i> )	2	2	3	0	2	2
House wren ( <i>Troglodytes aedon</i> )	18	10	11	*	14	12.5
<b>Common noncavity nesters</b>						
Dark-eyed junco ( <i>Junco hyemalis</i> )	14.7	12	10	12	4	12
American robin ( <i>Turdus migratorius</i> )	6	8	2.7	*	2	*
Hermit thrush ( <i>Catharus guttatus</i> )	4	4	2	6	2.7	4
Warbling vireo ( <i>Vireo gilvus</i> )	6	8	8	10	16	6
Yellow-rumped warbler ( <i>Dendroica coronata</i> )	4	6	8	4	2	*
Western wood-pewee ( <i>Contopus sordidulus</i> )	4.7	4	2	2	2	4.7

\*Present but not detected a sufficient number of times to qualify as a breeding resident.

more intricate than previously thought. Indeed, the subtleties in the requirements of the sapsuckers and their dual keystone roles suggests that preserving some species depends upon the distributions of species with which they have no obvious interaction. The disappearance of a single species could precipitate a wholly unanticipated unraveling of community structure.

Henry Gallin, Nicholas Lypps, and Bob Stuplich generously gave us permission to work on private land. We also thank Paul Buck, Roman Dial, and Kia Middleton for assistance with various aspects of the project. We greatly appreciate the advice of Marcus Feldman and Michael Gilpin on the statistical analyses. Thoughtful reviews of the manuscript were given by Susan Alexander, Jared Diamond, Marcus Feldman, Deborah Gordon, J. Pete Myers, Jonathan Roughgarden, and Peter Vitousek. This paper is dedicated to Peter and Helen Bing, with thanks for their friendship and support.

- Andrewartha, H. G. & Birch, L. C. (1954) *The Distribution and Abundance of Animals* (Univ. of Chicago, Chicago).
- MacArthur, R. H. & Wilson, E. O. (1967) *The Theory of Island Biogeography* (Princeton Univ. Press, Princeton).
- Tilman, D. (1987) *Am. Nat.* **129**, 769–774.
- Roughgarden, J. (1989) in *Perspectives in Ecological Theory*, eds. Roughgarden, J., May, R. M. & Levin, S. A. (Princeton Univ. Press, Princeton), pp. 203–226.
- Futuyma, D. (1973) *Am. Nat.* **107**, 443–446.
- Vitousek, P. M. & Walker, L. R. (1989) *Ecol. Monog.* **59**, 247–265.
- Polis, G. A. & Holt, R. D. (1992) *Trends Ecol. Evol.* **7**, 151–156.
- Ehrlich, P. R. & Wilson, E. O. (1991) *Science* **253**, 758–762.
- Paine, R. T. (1966) *Am. Nat.* **100**, 65–75.
- Paine, R. T. (1969) *Am. Nat.* **103**, 91–93.
- Janzen, D. H. (1971) *Science* **171**, 203–205.
- Howe, H. F. (1977) *Ecology* **58**, 539–550.
- Estes, J. A., Smith, N. S. & Palmisano, J. F. (1978) *Ecology* **59**, 822–833.
- Duggins, D. O. (1980) *Ecology* **61**, 447–453.
- Gilbert, L. E. (1980) in *Conservation Biology*, eds. Soule, M. E. & Wilcox, B. A. (Sinauer, Sunderland, MA), pp. 11–34.
- Terborgh, J. (1983) *Five New World Primates* (Princeton Univ. Press, Princeton), p. 60.
- Owen-Smith, N. (1987) *Paleobiology* **13**, 351–362.
- Owen-Smith, N. (1989) *Conserv. Biol.* **3**, 405–412.
- Brown, J. H. & Heske, E. J. (1990) *Science* **250**, 1705–1707.
- Hunter, M. D. (1992) in *Effects of Resource Distribution on Animal-Plant Interactions*, eds. Hunter, M. D., Ohgushi, T. & Price, P. W. (Academic, San Diego), pp. 288–317.
- Tate, J. (1973) *Auk* **90**, 840–856.
- Ehrlich, P. R. & Daily, G. C. (1988) *Am. Birds* **42**, 357–365.
- Miller, R. S. & Nero, R. W. (1983) *Can. J. Zool.* **61**, 1540–1546.
- Basham, J. T. (1958) *Can. J. Bot.* **36**, 491–505.
- Kilham, L. (1971) *Wilson Bull.* **83**, 159–171.
- Williams, A. B. (1936) *Ecol. Monogr.* **6**, 317–408.
- Kendeigh, S. C. (1944) *Ecol. Monogr.* **14**, 67–106.
- Emlen, J. T. (1977) *Auk* **94**, 455–468.
- Short, L. L. (1979) *Wilson Bull.* **91**, 16–28.