

# Prime factors

LIMING GE†

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139-4307

Communicated by Richard V. Kadison, University of Pennsylvania, Philadelphia, PA, August 19, 1996 (received for review August 6, 1996)

**ABSTRACT** We use Voiculescu’s free probability theory to prove the existence of prime factors, hence answering a longstanding problem in the theory of von Neumann algebras.

In a series of papers, Murray and von Neumann (1–5) introduced and studied certain algebras of Hilbert space operators, also known as *rings of operators*. They are now known as “von Neumann algebras.”

A von Neumann algebra is a strong-operator closed self-adjoint subalgebra of the algebra of all bounded linear transformations on a Hilbert space. *Factors* are von Neumann algebras whose centers consist of scalar multiples of the identity. They are the building blocks from which all the von Neumann algebras are built. The most elementary factors, type  $I_n$  factors, are isomorphic to the algebra  $M_n$  of all  $n \times n$  complex matrices. One of the basic constructions with factors (producing other factors) is that of forming the tensor product. For factors of type  $I_n$ ,  $M_p \otimes M_q$  is isomorphic to  $M_{pq}$ . Of course, no such tensor decomposition of  $M_p$  is possible precisely when  $p$  is a prime. The theory of tensor decompositions of factors of type  $I_n$  is little more than the theory of factoring integers into their prime components.

Murray and von Neumann (1) classified factors by means of a relative dimension function. *Finite factors* are those for which this dimension function has a finite range. For finite factors, this dimension function gives rise to a (unique, when normalized) tracial state. In general, a von Neumann algebra admitting a faithful normal trace is said to be *finite*. Infinite-dimensional finite factors are called factors of type  $II_1$ . They are “continuous” matrix algebras. Murray and von Neumann (2) spoke of “continuous dimensionality” in their factors of type  $II_1$ . In a parallel manner, when we study tensor products of factors of type  $II_1$  and their tensor-product decompositions, we may speak of decomposition into “continuous primes.” Factors of type  $II_1$  tensored with one another (as von Neumann algebras) produce, again, factors of type  $II_1$ . Each factor of type  $II_1$  may be decomposed as the tensor product of  $M_n$  and a factor of type  $II_1$  for each  $n$  in  $\mathbb{N}$ . In a sense, the “discrete primes” (that is, 2, 3, 5, . . .) are not significant in the theory of decomposition into “continuous primes.” The first question of the theory of such decompositions has to be that of the *existence* of a continuous prime: Is there a factor of type  $II_1$  that is not (isomorphic to) the tensor product of two factors of type  $II_1$ ? This problem and some related problems, concerning the basic structure of factors, have been asked and studied by many people (see, e.g., refs. 6 and 7, pp. 4.4.12 and 4.4.45). Popa (6) proves that there are prime factors of type  $II_1$  with a nonseparable predual. The separable case remained open. In this paper, we shall answer this question affirmatively.

We describe below, briefly, a basic construction of factors of type  $II_1$  by using regular representations of discrete groups. Our main result states that certain factors arising from free groups are prime. An outline of the proof follows the statement. We end with some open questions.

## Main Result

There are two main classes of examples of von Neumann algebras introduced by Murray and von Neumann (1, 3). One is obtained from the “group-measure space construction”; the other is based on regular representations of a (discrete) group  $G$  (with unit  $e$ ). The second class is the one needed in this paper. A brief description of that class follows.

The Hilbert space  $\mathcal{H}$  is  $l^2(G)$  (with its usual inner product). We assume that  $G$  is countable so that  $\mathcal{H}$  is separable. For each  $g$  in  $G$ , let  $L_g$  denote the left translation of functions in  $l^2(G)$  by  $g^{-1}$ . Then  $g \rightarrow L_g$  is a faithful unitary representation of  $G$  on  $\mathcal{H}$ . Let  $\mathcal{L}_G$  be the von Neumann algebra generated by  $\{L_g; g \in G\}$ . Similarly, let  $\mathcal{R}_g$  be the right translation by  $g$  on  $l^2(G)$  and  $\mathcal{R}_G$  be the von Neumann algebra generated by  $\{R_g; g \in G\}$ . Then the commutant  $\mathcal{L}'_G$  of  $\mathcal{L}_G$  is equal to  $\mathcal{R}_G$  and  $\mathcal{R}'_G = \mathcal{L}_G$ . The function  $u_g$  that is 1 at the group element  $g$  and 0 elsewhere is a cyclic trace vector for  $\mathcal{L}_G$  (and  $\mathcal{R}_G$ ). In general,  $\mathcal{L}_G$  and  $\mathcal{R}_G$  are finite von Neumann algebras. They are factors (of type  $II_1$ ) precisely when each conjugacy class in  $G$  (other than that of  $e$ ) is infinite. In this case, we say that  $G$  is an *infinite conjugacy class* (i.c.c.) group.

Specific examples of such  $II_1$  factors result from choosing for  $G$  any of the free groups  $F_n$  on  $n$  generators ( $n \geq 2$ ), or the permutation group  $\Pi$  of integers  $\mathbb{Z}$  (consisting of those permutations that leave fixed all but a finite subset of  $\mathbb{Z}$ ). Murray and von Neumann (3) prove that  $\mathcal{L}_{F_n}$  and  $\mathcal{L}_\Pi$  are not  $*$  isomorphic to each other (a deep result). A factor is *hyperfinite* if it is the ultraweak closure of the ascending union of a family of finite-dimensional self-adjoint subalgebras. In fact,  $\mathcal{L}_\Pi$  is the *unique* hyperfinite factor of type  $II_1$ ; it is contained in any factor of type  $II_1$ ; and the tensor product of the hyperfinite  $II_1$  factor with itself is  $*$  isomorphic to itself. Now we state our main theorem.

**MAIN THEOREM.** *The free group factor  $\mathcal{L}_{F_n}$  associated with the free group on  $n$  ( $\geq 2$ ) generators is prime, i.e., it is not isomorphic to the tensor product of any two factors of type  $II_1$ .*

We prove this result with the aid of Voiculescu’s free probability theory (8–10) (especially, his recently introduced concept of free entropy) and some geometrical methods for estimating free entropies. We refer to ref. 10 for the basics of free probability theory.

Let  $\mathcal{M}$  be a von Neumann algebra with a normal faithful trace  $\tau$ ,  $X_1, \dots, X_n$  be self-adjoint elements in  $\mathcal{M}$ . As analogues of classical entropy and of Fisher’s information measure, Voiculescu (8) introduced free entropy  $\chi(X_1, \dots, X_n)$ . Roughly speaking,  $\chi(X_1, \dots, X_n)$  is the limit of certain normalized measurement of all self-adjoint matrices that approximate  $X_1, \dots, X_n$  in joint distributions as the dimension of the matrices tends to infinity. We list some properties of free entropy in the following lemma.

**LEMMA 1.** *Let  $X_1, \dots, X_n$ ,  $n \geq 1$ , be self-adjoint elements in  $\mathcal{M}$  (with trace  $\tau$ ),  $C$  be  $\tau(X_1^2 + \dots + X_n^2)^{1/2}$  and  $R_0$  be  $\max\{\|X_j\|; j = 1, \dots, n\}$ . Then*

$$(i) \text{ (ref. 8; p. 2.2) } \chi(X_1, \dots, X_n) \leq n/2 \log(2\pi e C^2 n^{-1});$$

†e-mail: liming@math.mit.edu.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. §1734 solely to indicate this fact.

(ii) (ref. 8, p. 4.5)  $\chi(X_1) = \iint \log|s - t| d\mu_1(s) d\mu_1(t) + 3/4 + 1/2 \log 2\pi$ , where  $\mu_1$  is the (measure on the spectrum of  $X_1$  corresponding to the) distribution of  $X_1$ ;

(iii) (ref. 8; p. 5.4)  $\chi(X_1, \dots, X_n) = \chi(X_1) + \dots + \chi(X_n)$  when  $X_1, \dots, X_n$  are free random variables.

From the above lemma, we know that there are self-adjoint elements  $X_1, \dots, X_n$  in  $\mathcal{L}_{F_n}$  with finite free entropy such that  $X_1, \dots, X_n$  generate  $\mathcal{L}_{F_n}$  as a von Neumann algebra. In the following, we shall prove that if  $\mathcal{M}$  is not prime and  $X_1, \dots, X_n, n \geq 2$ , generate  $\mathcal{M}$  as a von Neumann algebra, then  $\chi(X_1, \dots, X_n) = -\infty$ . Hence,  $\mathcal{L}_{F_n}$  is prime. In fact, we prove a slightly stronger result in the following lemma.

**LEMMA 2.** Let  $\mathcal{M}$  be a factor of type  $II_1$ ,  $\mathcal{R}_1$  and  $\mathcal{R}_2$  be mutually commuting hyperfinite subfactors of  $\mathcal{M}$ . Let  $P_1, P_2, \dots$  and  $Q_1, Q_2, \dots$  be projections in  $\mathcal{M}$  with trace  $1/2$  that generate  $\mathcal{M}$  as a von Neumann algebra. Suppose that  $P_1, P_2, \dots$  commute with  $\mathcal{R}_1, Q_1, Q_2, \dots$  commute with  $\mathcal{R}_2$ . If  $X_1, \dots, X_n$  are self-adjoint elements in  $\mathcal{M}$  that generate  $\mathcal{M}$  as a von Neumann algebra, then  $\chi(X_1, \dots, X_n) = -\infty$ .

We give an outline of the proof here. The detailed argument will appear elsewhere.

From the assumptions in Lemma 2, we know that, for any positive  $\omega$ , there are projections  $P_1, \dots, P_p$  and  $Q_1, \dots, Q_q$  in  $\mathcal{M}$ ,  $p, q \in \mathbb{N}$ , and self-adjoint polynomials  $\varphi_1, \dots, \varphi_n$  in the noncommutative  $*$  polynomial ring  $\mathbb{C}\langle x_1, \dots, x_{p+q} \rangle$  such that

$$\|X_j - \varphi_j(P_1, \dots, P_p, Q_1, \dots, Q_q)\|_2 < \omega, \quad j = 1, \dots, n,$$

where  $\|\cdot\|_2$  is the trace norm ( $\|X\|_2^2 = \tau(X^*X)$ ,  $X \in \mathcal{M}$ ).

From the definition of free entropy (8), we shall estimate certain measurement of all (finite-dimensional) self-adjoint matrices that approximate  $X_1, \dots, X_n$  in joint distributions. For technical reasons, we use the notion of “modified” free entropy (9). More precisely, we estimate the free entropy of  $X_1, \dots, X_n$  in the presence of  $P_1, \dots, P_p$  and  $Q_1, \dots, Q_q$ . When self-adjoint elements  $A_1, \dots, A_n$  in  $M_k$  approximate  $X_1, \dots, X_n$  in joint distributions, there are projections  $E_1, \dots, E_p$  and  $F_1, \dots, F_q$  in  $M_k$  as well, corresponding to elements in certain Grassmann manifolds, that approximate projections  $P_1, \dots, P_p$  and  $Q_1, \dots, Q_q$  (in  $\mathcal{M}$ ) in joint distributions. At the same time,  $A_j$  are close (in trace-norm) to  $\varphi_j(E_1, \dots, E_p, F_1, \dots, F_q)$ .

From this observation, we are able to reduce the estimate of the free entropy of  $X_1, \dots, X_n$  to the volume estimate of the image of the cartesian product of the Grassmann manifolds under maps given by (noncommutative) polynomials  $\varphi_1, \dots, \varphi_n$ . Let  $k$  be the degree of the matrices that approximate  $X_j$ 's,  $D$  be an upper bound of the first derivatives of  $\varphi_j$ 's in the domain of the cartesian product of the Grassmann manifolds and  $d$  be the dimension of the manifolds. By using Szarek's results (11) on nets in unitary groups and Grassmann manifolds, we have that

$$\begin{aligned} \chi(X_1, \dots, X_n) &\leq \limsup_{k \rightarrow \infty} \left\{ k^{-2} \log \left( \left( \frac{CD\sqrt{p+q}}{\omega} \right)^{d(p+q)} \left( \frac{2aC}{\omega} \right)^{k^2} \pi^{\frac{1}{2}nk^2} \right. \right. \\ &\quad \left. \left. \times \Gamma \left( 1 + \frac{1}{2}nk^2 \right)^{-1} (3\omega\sqrt{nk})^{nk^2} \right) + \frac{n}{2} \log k \right\} \\ &\leq \frac{d(p+q)}{k^2} \log \left( \frac{CD\sqrt{p+q}}{\omega} \right) + \log \left( \frac{2aC}{\omega} \right) + \frac{n}{2} \log(n\pi) \\ &\quad + n \log 3 + n \log \omega + \limsup_{k \rightarrow \infty} \left( n \log k - k^{-2} \log \Gamma \left( 1 + \frac{nk^2}{2} \right) \right), \end{aligned}$$

where  $C$  is a universal constant,  $a = \max\{\|X_j\|_2 + 1 : 1 \leq j \leq n\}$ , and  $\Gamma(\cdot)$  is the classical  $\Gamma$ -function.

The assumptions that  $P_1, P_2, \dots$  commute with  $\mathcal{R}_1$  and  $Q_1, Q_2, \dots$  commute with  $\mathcal{R}_2$  give restrictions on the dimensions of the Grassmann manifolds. Hence, we can choose the dimension  $d$  so that  $d(p+q)/k^2 \leq \omega$  and  $d(p+q)/k^2 \log(CD\sqrt{p+q}) \leq \log(2C)$ . From Stirling's formula for the  $\Gamma$ -function, we have

$$\begin{aligned} \chi(X_1, \dots, X_n) &\leq \log(2C) - \omega \log \omega + \log(2aC) - \log \omega \\ &\quad + \frac{n}{2} \log(n\pi) + n \log 3 + n \log \omega \\ &\quad + \lim_{k \rightarrow \infty} \left( n \log k - \frac{n}{2} \log \left( \frac{nk^2}{2} \right) + \frac{n}{2} \right) \\ &\leq \log(4aC^2) + \frac{n}{2} \log(n\pi) + n \log 3 + \frac{n}{2} + (n-1-\omega) \log \omega. \end{aligned}$$

Choosing  $\omega$  arbitrarily small, we have  $\chi(X_1, \dots, X_n) = -\infty$ .

Using classical von Neumann algebra techniques, one can show that if  $\mathcal{M}$  is the tensor product of two factors of type  $II_1$  (i.e., not prime), then  $\mathcal{M}$  satisfies all the hypotheses of Lemma 2.

### Open Problems

Some questions about decompositions into continuous primes, analogous to simple facts about (discrete-)prime-factor decomposition, and about continuous primes, themselves, come instantly to mind:

1. Are there infinitely many (nonisomorphic) prime factors of type  $II_1$ ?
2. Is  $\mathcal{L}_{F_2} \otimes \mathcal{L}_{F_2} * \mathcal{L}_{F_2}$  isomorphic to  $\mathcal{L}_{F_2} \bar{\otimes} \mathcal{L}_{F_2} \bar{\otimes} \mathcal{L}_{F_2}$  (“uniqueness” of prime decomposition)?
3. With  $\mathcal{M}$  a factor of type  $II_1$ , let  $p(\mathcal{M})$  be the set of integers  $n$  for which there are prime factors  $\mathcal{M}_1, \dots, \mathcal{M}_n, n \in \mathbb{N}$ , of type  $II_1$  such that  $\mathcal{M} \cong \mathcal{M}_1 \otimes \dots \otimes \mathcal{M}_n$ . If there is no such  $n$ , let  $p(\mathcal{M}) = \{\infty\}$ . Does  $p(\mathcal{M})$  contain only one number?

Finally, we propose the project of classifying all von Neumann subalgebras of free group factors as an analogue of Connes's classification of von Neumann subalgebras of the hyperfinite  $II_1$  factor (12), and ask a question suggested by Lemma 2. Is the relative commutant of a nonatomic injective (or abelian) von Neumann subalgebra of  $\mathcal{L}_{F_n}$  in  $\mathcal{L}_{F_n}$  always injective?

This work was supported by the National Science Foundation.

1. Murray, F. J. & von Neumann, J. (1936) *Ann. Math.* **37**, 116–229.
2. Murray, F. J. & von Neumann, J. (1937) *Trans. Am. Math. Soc.* **41**, 208–248.
3. Murray, F. J. & von Neumann, J. (1943) *Ann. Math.* **44**, 716–808.
4. von Neumann, J. (1930) *Math. Ann.* **102**, 370–427.
5. von Neumann, J. (1940) *Ann. Math.* **41**, 94–161.
6. Popa, S. (1983) *J. Oper. Theory* **9**, 253–268.
7. Sakai, S. (1971) *C\*-Algebras and W\*-Algebras* (Springer, New York).
8. Voiculescu, D. (1994) *Invent. Math.* **118**, 411–440.
9. Voiculescu, D. (1996) *Geom. Funct. Anal.* **6**, 172–199.
10. Voiculescu, D., Dykema, K. & Nica, A. (1992) *Free Random Variables*, Centre de Recherches Mathématiques Université de Montréal Monograph Series (Am. Math. Soc., Providence, RI), Vol. 1.
11. Szarek, S. J. (1981) in *Proceedings of Research Workshop on Banach Space Theory*, ed. Lin, B.-L. (Univ. of Iowa, Iowa City), pp. 169–185.
12. Connes, A. (1976) *Ann. Math.* **104**, 73–115.