

# Genomewide insertional mutagenesis in *Streptomyces coelicolor* reveals additional genes involved in morphological differentiation

Amy M. Gehring\*, Justin R. Nodwell\*<sup>†</sup>, Stephen M. Beverley<sup>‡</sup>, and Richard Losick\*<sup>§</sup>

\*Department of Molecular and Cellular Biology, Harvard University, 16 Divinity Avenue, Cambridge, MA 02138; and <sup>†</sup>Department of Molecular Microbiology, Washington University Medical School, St. Louis, MO 63110

Contributed by Richard Losick, February 10, 2000

The filamentous soil bacterium *Streptomyces coelicolor* undergoes a complex cycle of morphological differentiation involving the formation of an aerial mycelium and the production of pigmented antibiotics. We have developed a procedure for generating insertional mutants of *S. coelicolor* based on *in vitro* transposition of a plasmid library of cloned *S. coelicolor* DNAs. The insertionally mutated library was introduced into *S. coelicolor*, and transposon insertions were recovered at widely scattered locations around the chromosome. Many of the insertions revealed previously uncharacterized genes, and several caused novel mutant phenotypes, such as altered pigment production, enhanced antibiotic sensitivity, delayed or impaired formation of aerial hyphae, and a block in spore formation. The sporulation mutant harbored an insertion in one of three adjacent genes that are apparently unique to *Streptomyces* but are each represented by at least 20 paralogs at dispersed locations in the chromosome. Individual members of the three families often are found grouped together in a characteristic arrangement, suggesting that they have a common function.

**S**treptomyces are filamentous soil bacteria that undergo a complex cycle of morphological differentiation (1–3). The cycle begins with a spore, which germinates to give rise to a branching network of multinucleoid hyphae known as the substrate mycelium. Morphological differentiation commences with the formation of specialized aerial hyphae, which grow into the air away from the surface of the colony. Later these aerial hyphae undergo septation into uninucleoid compartments and metamorphose into chains of gray-pigmented spores. The onset of differentiation frequently is associated with the production of pigments, antibiotics, and other secondary metabolites (3, 4).

Relatively few morphological mutants have been described in *Streptomyces*, and even fewer have been characterized at the molecular level. One reason for slow progress is that cloning genes identified by chemically induced mutations is tedious in *Streptomyces*. As a means of identifying genes involved in morphological differentiation and other aspects of *Streptomyces* biology, we set out to develop a method for generating insertional mutants of *Streptomyces coelicolor* A3(2), the most intensively studied of the *Streptomyces* (5). Although several transposon systems have been described for use in *Streptomyces* (6–9), none have been used successfully to generate large numbers of random insertions, and in only two cases have insertion mutants (10, 11) been recovered in genomewide screens in which the mutant phenotype could be attributed to the transposon insertion.

As a fresh approach to the challenge of generating insertional mutants in *Streptomyces* on a genomewide basis, we took advantage of three recent technical advances: improvements in DNA-mediated transformation of *Streptomyces* (12), the emerging availability of a genome sequence for *S. coelicolor* (5), and the development of efficient systems for carrying out transposition *in vitro*. Biochemical systems now have been described for carrying out transposition *in vitro* with high efficiency for the transposable elements Tn7 (13), Ty1 (14), Tn5 (15), *mariner* (16,

17) Tn552 (18), and Mu (19). Several of these systems have been used to carry out insertional mutagenesis of bacteria that are naturally competent, such as *Haemophilus influenzae* (20–22) and *Streptococcus pneumoniae* (21).

Here we describe the introduction into *S. coelicolor* of plasmid libraries of *S. coelicolor* DNA that had been subjected to insertional mutagenesis with derivatives of *mariner* or Tn5. After selection for a transposon-borne drug-resistance gene, we screened for colonies in which the transposon had been incorporated into the corresponding region of homology with the chromosome by double (marker-replacement) recombination. Sequence analysis showed that transposon insertions had been recovered at scattered locations around the *S. coelicolor* chromosome and revealed a wide spectrum of heretofore unidentified genes. Two such insertions blocked the production of the pigmented antibiotic actinorhodin and enhanced apramycin sensitivity, respectively. Other insertions impaired the process of aerial mycelium formation, thereby identifying previously unrecognized genes that seem to play a role in morphological differentiation. One insertion, which resulted in a block in sporulation, was located in one of three adjacent genes that define three previously unrecognized families of paralogous genes in the *S. coelicolor* chromosome.

## Materials and Methods

**Strains and Growth Conditions.** *S. coelicolor* strains were M145 (prototrophic, SCP1<sup>+</sup> SCP2<sup>+</sup>) (23) or J1501 (*hisA1 uraA1 strA1* SCP1<sup>+</sup> SCP2<sup>+</sup>) (24). Strains were grown on solid R2YE or minimal medium with 1% glucose (wt/vol) or in liquid yeast extract-malt extract (YEME) medium (23) at 30°C with, as indicated, 25 µg/ml apramycin sulfate or 200 µg/ml spectinomycin dihydrochloride (Sigma).

**Transposons.** Transposon derivatives were created that harbored *aac(3)IV*, an apramycin resistance gene (25) (hereafter designated *apr*), from pOJ427 (Eli Lilly). The *apr* gene was cloned into a unique *Xba*I site located between the 3' and 5' inverted repeats of the transposable element *mariner* (*Mos1*) in pELHY6–0 (S. Goyard and S.M.B., unpublished work), yielding pELApr. Likewise, *apr* was cloned between the *Bam*HI and *Xba*I sites of pMOD<MCS> (Epicentre Technologies, Madison, WI) to give pMODApr. Linear transposon DNA, Tn5*apr*, was liberated from pMODApr by digestion with *Pvu*II.

Abbreviation: Apr<sup>R</sup>, apramycin-resistant.

<sup>†</sup>Present address: Department of Biochemistry, McMaster University, 1200 Main Street West, Hamilton, ON, L8N 3Z5 Canada.

<sup>§</sup>To whom reprint requests should be addressed. E-mail: losick@envoy.mcb.harvard.edu.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

Article published online before print: *Proc. Natl. Acad. Sci. USA*, 10.1073/pnas.170059797. Article and publication date are at www.pnas.org/cgi/doi/10.1073/pnas.170059797

**Table 1. Tn5apr insertion sites**

Strain	Disrupted sequence (5'→3')*	Cosmid†
AG24	1084 CGGCACGCCGCCGACGACGCCCG 1061	SC8B7
AG137	1665 CGGAGCGCGGTGAGATCCTGCGCG 1642	SCE134
AG259	11872 GCGCACACCGACGCCCTGTCCGG 11895	St9A4
AG534	16012 GCCGACGAGCTGGGCCCGCGCGC 16035	SCI8
AG643	11275 CGCGTCACCGCAGATCTGGGCGCC 11298	SCH69
AG671	12027 GGGTGTAGTACAGGTCGATGTGGT 12004	SCD17
AG848	GTCGACACCTATGCTGCGGTCATC	?
AG860	19659 GCGGACGCCCTACCGGCAGACCGAG 19636	SCF42
AG1057	CGAAGCCTGCCTCAACCTGACTCA	?
AG1060	20511 GAGCACCGTGTGAGGCTGGCGCG 20488	SCAH10
AG1061	1895 TCCCGCCCTCAAGAGGCAGGCTC 1872	SCH17
AG1069	TGGACCAGCAAGTGCCCATGGAGT	?
AG1225	4955 CTTGGAACGTGCGGGTCGAACAGG 4932	SCI11
AG1329	19008 GCACGCACTCTCGCCCCACCGGGC 18985	SCF43
AG1338	6054 GAGGACGGACACGACCCAGCGCAG 6031	SCD25
AG1346	28787 GCGGCGCGCACCGCTCCGGGCGCC 28810	2St10A7
AG1359	15322 TGGGCAACCACCGCGCCGAGA 15299	2StG58

\*Transposon insertion sites are shown in bold along with flanking DNA sequence. Numbers denote the bp location of the sequence in the indicated *S. coelicolor* cosmid if known.


†Cosmid sequence information is available in the European Molecular Biology Laboratory/GenBank/DNA Data Base in Japan databases (only those labeled SC) and/or from the *S. coelicolor* Sequencing Group at the Sanger Centre at [ftp://ftp.sanger.ac.uk/pub/S\\_coelicolor/sequences](ftp://ftp.sanger.ac.uk/pub/S_coelicolor/sequences).

**Plasmid Libraries.** Plasmid libraries of *S. coelicolor* DNA were constructed in the vectors pSpec and pSpecoriT. To create pSpec, the  $\Omega$  fragment from pHP45 $\Omega$  (26), which contains the spectinomycin resistance gene *aadA* (hereafter designated *spec*), was first cloned into the *Hind*III site of pBluescript II SK(+) (Stratagene). The  $\Omega$ -containing *Bss*HII fragment isolated from this plasmid was ligated to the *ColE1* origin [amplified by PCR from pBluescript II SK(+)], and the  $\Omega$ -containing *Hind*III fragment was replaced by *spec* (amplified by PCR from HP45 $\Omega$ ). To create the conjugative vector pSpecoriT, the *oriT* sequence from pOJ446 (27) was cloned into the *Pst*I site of pSpec. To create plasmid libraries, genomic DNA was isolated from *S. coelicolor* M145 (28), digested with *Sau*3A1, and cloned into the *Bam*HI site of pSpec or pSpecoriT.

**Insertional Mutagenesis.** Transposition reactions were carried out by using cloned libraries of *S. coelicolor* DNA (above) and pELApr as described by Tosi and Beverley (17) or Tn5apr as described by the manufacturer (Epicentre Technologies). DNA from the reaction mixtures was used to transform *Escherichia coli* with selection for transformants on LB plates containing spectinomycin and apramycin. Tn5apr gave a higher transposition efficiency ( $\approx 1$  per 2,000 hybrid plasmids) and was used for most of the insertional mutagenesis experiments in this study.

The transposon-mutated hybrid plasmids were introduced into *S. coelicolor* by protoplast transformation or by conjugation from *E. coli*. Unmethylated preparations of the transposon-disrupted pSpec hybrid plasmids isolated from *E. coli* SCS110 (Stratagene) (0.5–2  $\mu$ g) were transformed into M145 protoplasts (12, 23), plated on R2YE, and apramycin-resistant (*Apr*<sup>R</sup>) transformants ( $\approx 10^3$  per experiment) selected by overlay with apramycin after 16 h. Tn5-disrupted pSpecoriT hybrid plasmids were introduced into *S. coelicolor* M145 by conjugation (29) from *E. coli* strain ET12567(pUB307) transformed with the hybrid plasmids. Exconjugants were selected with apramycin (29).

The *S. coelicolor* *Apr*<sup>R</sup> transformants or exconjugants were screened for mutant phenotypes. Colonies of interest were restreaked on R2YE plates containing apramycin. Next, colonies of *Apr*<sup>R</sup> cells were replica-plated on R2YE plates containing



**Fig. 1.** Chromosomal locations of transposon insertions. Black arrows indicate insertions that arose from double recombination and exhibited no conspicuous phenotype, red arrows indicate insertions from double recombination that were associated with a mutant phenotype, and green arrows indicate integrants in which the transposon-containing plasmid had inserted into the chromosome by single recombination. The map positions of insertions 848, 1057, and 1069 await completion of the genome sequence. Known genes are indicated inside the chromosome (31). Circles represent the ends of the linear *S. coelicolor* chromosome, and the diamond represents the origin of replication.


spectinomycin to identify those that were *Apr*<sup>R</sup> and spectinomycin-sensitive (about 5% of the total). To test for linkage between the transposon insertion and the mutation causing the mutant phenotype, genomic DNA was prepared (28), alkali-denatured (up to 25  $\mu$ g; ref. 12), and used to transform protoplasts of the parent strain. Transformants were selected with apramycin and then scored for their phenotype. For about 20–25% of strains examined, the transposon insertion cotransformed with the mutant phenotype. Genomic DNA transformation also was used to transfer *mariner*-58 from the J1501 background in which it was isolated to M145 to give AGJ58.

**Identification of *S. coelicolor* Sequences Flanking Transposon Insertions.** Genomic DNA from the insertion strains was digested with *Sac*II or *Not*I, ligated to pBluescript II SK(+), and the ligated DNA was used to transform *E. coli* followed by selection with apramycin. Plasmid DNA from the *Apr*<sup>R</sup> transformants was isolated, and flanking DNA was sequenced by using primers designed to anneal to opposite ends of *apr*.

## Results and Discussion

**Strategy for Generating Insertional Mutants.** Plasmid libraries of *S. coelicolor* DNA were subjected to transposition *in vitro* by using derivatives of the transposons Tn5 (Tn5apr) or *mariner*. The products of the reactions were introduced into *E. coli* followed by selection for both vector-borne (*spec*) and transposon-borne (*apr*) drug resistance genes, thereby enriching for transformants that harbored hybrid plasmids with a transposon insertion. Next, the transposon-bearing hybrid plasmids were introduced into *S. coelicolor* by protoplast transformation or by conjugation from *E. coli* followed by selection for *Apr*<sup>R</sup>. Finally, we screened the *Apr*<sup>R</sup> cells for those that were spectinomycin-sensitive. This distinguished integrants that arose from single, reciprocal (Campbell-like) recombination in which the entire transposon-bearing, hybrid plasmid had inserted into the chromosome from





**Fig. 2.** *S. coelicolor* insertional mutants. Shown are color photographs of colonies of the wild-type parent strain M145 and the indicated insertional mutants that had been grown on solid R2YE medium. (A) Images from the top of the plates. (B) Corresponding images from the back of the plates.

integrants that arose from double (marker-replacement) recombination in which only the transposon had been inserted.


**Chromosomal Locations of Transposon Insertions.** We characterized 18 such Apr<sup>R</sup>-spectinomycin-sensitive integrants, 17 of which were generated with Tn5<sub>apr</sub> and one with *mariner*. The site of insertion of each transposon was determined by sequencing DNA flanking the element. For strain AGJ58, which contains *mariner*, we observed the expected insertion into, and duplication of, a TA dinucleotide (base pairs 7762–7763 of SC4H8). Tn5 insertion causes duplication of a 9-bp target sequence that exhibits base preferences at certain positions (30). The Tn5<sub>apr</sub> insertions characterized were indeed flanked by a duplication of a 9-bp stretch of chromosomal DNA (Table 1). Sequences flanking the transposon for each mutant strain were used to search the Sanger Centre database ([ftp://ftp.sanger.ac.uk/pub/S\\_coelicolor/sequences](ftp://ftp.sanger.ac.uk/pub/S_coelicolor/sequences)) to position the transposon on the *S. coelicolor* chromosome (Fig. 1). A perfect match to the database was found for sequences flanking 15 of the 18 transposon insertions, which were located at scattered sites around the chromosome.

At least six insertions that did not cause a detectable phenotype were located within ORFs. These were insertions in the unknown genes *SCE134.01c*, *SCF42.21c*, and *SCH69.11c*, the putative oxidoreductase genes *SCI8.16* and *SCD17.13*, and the putative ABC transporter gene *SCAH10.23*. Five other insertions did cause a mutant phenotype and are described below.

**An Insertion Mutation Altering Pigment Production.** One of the insertion mutants, strain AG1338, exhibited a striking pigment production phenotype (Fig. 2). AG1338 failed to produce the blue pigment (actinorhodin) that is characteristic of the wild-type strain M145. The substrate mycelium of AG1338 initially exhibited a light orange color and then turned brown as growth proceeded. AG1338 was sporulation-proficient (indeed, sporulation may have been faster than in the wild type), but the spores developed a darker color than that characteristic of the wild type. *S. coelicolor* spores are normally gray, but *whiE*, which is responsible for the spore pigment, can give rise to a related brown pigment (32, 33, 46). Conceivably, AG1338 overproduces the *whiE*-related brown pigment at the expense of actinorhodin.

The Tn5<sub>apr</sub> insertion in AG1338 was located within a previously uncharacterized ORF, *SCD25.05*, which encodes a putative integral membrane protein of unknown function. Because *Streptomyces* is known to exhibit a significant spontaneous mutation rate (34, 35, 47), it was important to determine whether the phenotype of AG1338 was caused by the transposon insertion or by an unlinked spontaneous mutation. To investigate this, we carried out transformation with chromosomal DNA (12) to ask whether the transposon insertion would cotransform with the mutation, causing brown pigment production. Eleven of 11 of the Apr<sup>R</sup> transformants exhibited a phenotype identical to that of the parent strain AG1338.


**An Insertion Causing Increased Apramycin Sensitivity.** Strain AG1225 grew normally on R2YE (rich) medium (Fig. 2) with or



**Fig. 3.** Phase-contrast micrographs of the wild-type strain M145 (A), a *whi1* strain bearing the mutation *whi-77* (B), and the insertional mutant AG1057 (C). Slides were prepared by taking an impression of the surface of colonies grown for 4 days on solid R2YE medium. Visualization was with an Olympus BX60 microscope at  $\times 100$  magnification.

without apramycin and on minimal medium without drug, but grew slowly on minimal medium supplemented with apramycin (25  $\mu\text{g/ml}$ ) (data not shown). The apramycin sensitivity mutation was shown to be 100% (43 of 43 transformants) linked to the *Tn5apr* insertion by cotransformation experiments with chromosomal DNA. This insertion was located to an intergenic region 36 bp downstream of gene *SCJ11.08c* in a large direct repeat that flanks this gene. The *SCJ11.08c* gene product is 98% identical to the product of a previously described *S. lividans* gene (ORF2) found at the junction with the terminal inverted repeat at one end of the linear chromosome of this organism (36). The product of *SCJ11.08c* also shows strong similarity to integral membrane transport proteins including chloramphenicol resistance protein homologs and metabolite transporters as determined by a BLAST search of the protein databases (37). Perhaps the *SCJ11.08c* gene product contributes to *Apr<sup>R</sup>*, and the *Tn5apr-1225* insertion destabilizes the *SCJ11.08c* transcript, reducing levels of the putative efflux transporter and thereby allowing an inhibitory concentration of apramycin to accumulate in the cell when grown on minimal medium. Conceivably and alternatively, insertion into the direct repeat adjacent to *SCJ11.08c* might interfere with expression of the transposon-borne *apr* gene.

**An Insertion Causing Formation of Abnormal Aerial Hyphae.** Strain AG24 formed an abnormal aerial mycelium that was purple in color and became purplish-pink as development proceeded (Fig. 2). The aerial hyphae of AG24 were abnormally short, and the mutant colonies exhibited cracks over their surface. Light microscopy revealed that AG24 produced spores, but some were aberrantly sized (larger or smaller than the wild type), and their quantity was significantly reduced as compared with the M145 parent (data not shown). The *Tn5apr* insertion in AG24 was 100% (46 of 46 transformants) linked to the mutation causing abnormal aerial hyphae formation in cotransformation experiments with chromosomal DNA. The insertion was located within




**Fig. 4.** Chromosomal locations of members of the 283-codon ORF (red), 63-codon ORF (blue), and 141-codon ORF (green) families of paralogous genes. Clusters of adjacent ORFs are designated by end-to-end blue, red, and green lines. Counterclockwise from the left end of the chromosome are: SCF42.12c-14; 2StG1 base pairs 14464–13952, 14680–15540, 15550–15792; SC1G2.07c, 08c; SC1G2.14c, 15c; SC4A7.09, 10; SC121.16, 17; SCC57a.09c; SCC57a.10c, 11; SCE6.01c, 02, base pairs 1594–1800; SCE6.06; SCE94 base pairs 13886–13695, 16c, base pairs 14791–15471; SCE9.28c; SCE9.30, 31c; SCD66.13, 14; SCD6.19, 20; SCD31.02c-04; StD77 base pairs 31404–31135, 32227–31388; St9E12 base pairs 9058–8852, 9870–9055; SC9B2.15c-17; SC2H4.17c, SCAH10.01c, 02; SC5C7.22c; SC4G2.02c-04; St5F1 base pairs 32071–32946; St2H2 base pairs 12405–11959, 12579–13451, 13448–13717. ORFs designated by cosmid base pair numbers have not yet been annotated by the Sanger Centre Sequencing Group. SCF42.13, SCE9.31c, and St2H2 base pairs 13448–13717 are included as members of the 63-codon family but are more weakly similar than are other family members. Only paralogs of the 141-codon ORF that are associated with a 283-codon and/or a 63-codon family member are shown.

the ORF *SC8B7.01c/SC5H4.01*, the inferred product of which is annotated as containing two possible transmembrane domains and a possible ATP/GTP binding site (P-loop).

**An Insertion Causing Delayed Aerial Mycelium Formation.** Mutant strain AGJ58 showed an approximately 1-day delay in aerial mycelium formation, and the aerial mycelium exhibited a pinkish-gray color (Fig. 2). Transformation experiments with chromosomal DNA showed that the *mariner* insertion in AGJ58 was 100% (50 of 50 transformants) linked to the mutation causing delayed aerial mycelium formation. The insertion was located in an intergenic region between the ORF *SC4H8.12c* and a pair of downstream, partially overlapping ORFs, *SC4H8.11c* and *SC4H8.10c*. As determined by a BLAST search of the protein databases (37), the inferred product of the upstream ORF *SC4H8.12c* exhibits strong similarity to bacterial hydroxylases. *SC4H8.11c* and *SC4H8.10c* are similar to a pair of genes, *Rv1065* and *Rv1066*, found in *Mycobacterium tuberculosis*. The *SC4H8.11c* gene product shows weak similarity to cysteine dioxygenases, whereas the *SC4H8.10c* product is weakly similar to putative thiosulfate sulfurtransferases. Interestingly, the activity of a cysteine dioxygenase is believed to play a role in the transition from the mycelial to the yeast phase in the dimorphic pathogenic fungal species *Histoplasma capsulatum* (38–40).

**An Insertion Mutation Blocking Spore Formation at an Early Stage.** Strain AG1057 exhibited a so-called white (*whi*) mutant phenotype (1), that is, it produced a normal-looking aerial mycelium but was markedly impaired in spore formation and the produc-





**Fig. 5.** Alignment of conserved amino acid sequences for inferred protein products of 283-codon paralogs (A), 63-codon paralogs (B), and 141-codon paralogs (C). The top line in each alignment corresponds to the ORF identified by the *Tn5apr-1057* insertion. The other examples correspond to cases identified in Fig. 4 in which a 283-codon family member is flanked downstream by a paralog of the 63-codon ORF and upstream by an oppositely oriented paralog of the 141-codon ORF. All eight proteins are similar along their lengths, but only a subset of the conserved regions are shown.

tion of spore-associated gray pigment (Fig. 2). As shown in Fig. 3, at a time (4 days) when the aerial hyphae of the parent strain M145 had sporulated, the aerial hyphae of AG1057 remained undifferentiated, appearing as straight, branched, or somewhat wavy filaments. Cotransformation experiments with chromosomal DNA demonstrated 100% (12 of 12 transformants) linkage between the *Tn5apr* insertion in AG1057 and the mutation causing the *whi* mutant phenotype.

DNA flanking the *Tn5apr* insertion in AG1057 was sequenced, but no match was found in the Sanger Centre database. There was, however, a perfect match to a portion of a 3.8-kb sequence that had been deposited in GenBank (accession number AF106004). The *Tn5apr* insertion was located within a 283-codon ORF in the deposited sequence. BLAST analysis (37) of this ORF failed to reveal strong similarities to sequences outside of the genus *Streptomyces*. Remarkably, however, 21 paralogs were detected within the current *S. coelicolor* database, and these were located at widely scattered sites on the chromosome (Fig. 4). The 283-codon ORF is just upstream of, and in the same orientation as, an ORF of 63 codons and downstream of an oppositely oriented ORF of 141 codons. The 63-codon ORF and the 141-codon ORF each were also represented by at least 20 paralogs at dispersed locations in the *S. coelicolor* genome (Fig. 4). Members of all three families of paralogs exhibit several blocks of strikingly conserved amino acid coding sequences along their lengths, examples of which are shown in Fig. 5. We also note the presence of a potential helix–turn–helix motif near the N terminus of the 283-codon ORF and each of its paralogs as revealed by PFAM analysis (41). Strikingly, 16 of the 21 paralogs of the 283-codon ORF were located immediately

upstream of a paralog of the 63-codon ORF, and in eight of these 16 cases a paralog of the 141-codon ORF also was found upstream of the 283-codon paralog in the opposite orientation. We conclude that *Tn5apr-1057* defines three novel families of paralogous genes and that members of each family frequently are found adjacent to each other in a characteristic arrangement.

Because the Sanger sequencing project has not yet included the 283-codon ORF, the location of *Tn5apr-1057* is not known. DNA corresponding to GenBank accession number AF106004 was assigned as the *whiJ* locus and was previously mapped by DNA hybridization to the overlap of cosmids D39 and D82 (31). But the sequences of D39 and D82 are now known (Sanger Centre) and evidently do not contain the AF106004 sequence. Moreover, the phenotype of the *Tn5apr-1057* insertion mutant was much more severe than that of a *whiJ* mutant (*whi-77*; refs. 1 and 42) as shown in Fig. 3. Conceivably, *whi-77* is a leaky allele of the 283-codon ORF. Alternatively, *whi-77* could be allelic to the 63-codon ORF, the 141-codon ORF or another ORF in the 3.8-kb segment of DNA. It will be interesting to see where *Tn5apr-1057* (and *whiJ*) is located on the *S. coelicolor* chromosome once the Sanger sequencing project is completed.

**Insertional Mutants Arising from Single Recombination.** Transposon integration also can arise by single, reciprocal recombination in which the entire transposon-bearing, hybrid plasmid is incorporated into the chromosome. Such Apr<sup>R</sup>–spectinomycin-resistant integrants could exhibit a mutant phenotype if the *S. coelicolor* insert carried by the plasmid is internal to a transcription unit (43). Two mutant strains that arose from integration of a hybrid plasmid were AG29 and AG59. Strain AG29 was auxotrophic

and strain AG59 failed to produce the red pigment undecylprodigiosin. In both cases, the vector-borne drug resistance gene was 100% linked to the mutation responsible for the mutant phenotype in cotransformation experiments with chromosomal DNA. DNA sequence analysis showed that AG29 harbored a hybrid plasmid insert (base pairs 13638–18321 of cosmid SC4G6; Fig. 1) within the histidine biosynthesis operon (44). Consistent with this, the mutant was able to grow on minimal medium that had been supplemented with histidine (not shown). Likewise, AG59 harbored a hybrid plasmid insert within a cluster of (*red*) genes (base pairs 17370–22882 of cosmid SC3F7; Fig. 1) responsible for undecylprodigiosin biosynthesis (31, 45).

**Summary.** We have succeeded in introducing insertions of the transposons Tn5 or *mariner* at widely scattered locations around the *S. coelicolor* chromosome. This has revealed multiple novel genes associated with defects in secondary metabolism, drug resistance, and morphological differentiation. In each case cotransformation with chromosomal DNA showed that the transposon insertion was linked to the mutation causing the mutant phenotype. Specifically, we identified an insertion in a previously uncharacterized gene that blocked actinorhodin production and resulted instead in the production of a brown pigment. We identified an insertion that caused apramycin sensitivity and is

located adjacent to a gene whose inferred product is similar to antibiotic efflux pumps. We recovered two insertions in or near uncharacterized genes that delayed or impaired aerial mycelium formation. We also identified an insertion that blocked the conversion of aerial hyphae into chains of spores, defining a previously uncharacterized *whi* gene. Interestingly, the *whi* gene and two flanking ORFs are each members of novel families of paralogous genes. Individual members from each family frequently are found clustered together in a characteristic arrangement at dispersed locations in the chromosome. It will be interesting to determine whether the three gene families have related functions and whether any of the additional gene clusters are involved in morphological differentiation. The stage is now set to use insertional mutagenesis on a large scale for discovering genes involved in morphological differentiation, secondary metabolism, and other aspects of *Streptomyces* biology.

We thank S. Goyard for pELHY6–0, Z. Li for pELApr, and S. Goyard and J. Guzova for transposition tests of pELApr. We thank M. Buttner, K. Chater, D. Hopwood, J. McCormick, L. Shapiro, and J. Westpheling for helpful comments on the manuscript. This work was supported by a grant (to R.L.) from the National Science Foundation (MCB-9727234) and National Institutes of Health Grant AI29646 (to S.M.B.). A.M.G. is supported by the Cancer Research Fund of the Damon Runyon-Walter Winchell Foundation Fellowship, DRG-1524.

- Chater, K. F. (1998) *Microbiology* **144**, 1465–1478.
- Kelemen, G. H. & Buttner, M. J. (1998) *Curr. Opin. Microbiol.* **1**, 656–662.
- Champness, W. (2000) in *Prokaryotic Development*, eds. Brun, Y. V. & Shinkets, L. J. (Am. Soc. Microbiol., Washington, DC), pp. 11–31.
- Bibb, M. (1996) *Microbiology* **142**, 1335–1344.
- Hopwood, D. A. (1999) *Microbiology* **145**, 2183–2202.
- Olson, E. R. & Chung, S.-T. (1988) *J. Bacteriol.* **170**, 1955–1957.
- Baltz, R. H., Hahn, D. R., McHenney, M. A. & Solenberg, P. J. (1992) *Gene* **115**, 61–65.
- Volff, J.-N. & Altenbuchner, J. (1997) *Gene* **194**, 81–86.
- Herron, P. R., Evans, M. C. & Dyson, P. J. (1999) *FEMS Microbiol. Lett.* **171**, 215–221.
- Ikeda, H., Takada, Y., Pang, C.-H., Tanaka, H. & Omura, S. (1993) *J. Bacteriol.* **175**, 2077–2082.
- McHenney, M. A., Hosted, T. J., Dehoff, B. S., Rostek, P. R., Jr. & Baltz, R. H. (1998) *J. Bacteriol.* **180**, 143–151.
- Oh, S.-H. & Chater, K. F. (1997) *J. Bacteriol.* **179**, 122–127.
- Bainton, R. J., Kubo, K. M., Feng, J. & Craig, N. L. (1993) *Cell* **72**, 931–943.
- Devine, S. E. & Boeke, J. D. (1994) *Nucleic Acids Res.* **22**, 3765–3772.
- Goryshin, I. Y. & Reznikoff, W. S. (1998) *J. Biol. Chem.* **273**, 7367–7374.
- Lampe, D. J., Grant, T. E. & Robertson, H. M. (1998) *Genetics* **149**, 179–187.
- Tosi, L. R. O. & Beverley, S. M. (2000) *Nucleic Acids Res.* **28**, 784–790.
- Griffin, T. J., IV, Parsons, L., Leschziner, A. E., DeVost, J., Derbyshire, K. M. & Grindley, N. D. F. (1999) *Nucleic Acids Res.* **27**, 3859–3865.
- Haapa, S., Taira, S., Heikkinen, E. & Savilahti, H. (1999) *Nucleic Acids Res.* **27**, 2777–2784.
- Gwinn, M. L., Stellwagen, A. E., Craig, N. L., Tomb, J.-F. & Smith, H. O. (1997) *J. Bacteriol.* **179**, 7315–7320.
- Akerley, B. J., Rubin, E. J., Camilli, A., Lampe, D. J., Robertson, H. M. & Mekalanos, J. J. (1998) *Proc. Natl. Acad. Sci. USA* **95**, 8927–8932.
- Reich, K. A., Chovan, L. & Hessler, P. (1999) *J. Bacteriol.* **181**, 4961–4968.
- Hopwood, D. A., Bibb, M. J., Chater, K. F., Kieser, T., Bruton, C. J., Kieser, H. M., Lydiate, D. J., Smith, C. P., Ward, J. M. & Schrepf, H. (1985) *Genetic Manipulation of Streptomyces: A Laboratory Manual* (The John Innes Foundation, Norwich, U.K.).
- Chater, K. F., Bruton, C. J., King, A. A. & Suarez, J. E. (1982) *Gene* **19**, 21–32.
- Braü, B., Pilz, U. & Piepersberg, W. (1984) *Mol. Gen. Genet.* **193**, 179–187.
- Prentki, P. & Krisch, H. M. (1984) *Gene* **29**, 303–313.
- Bierman, M., Logan, R., O'Brien, K., Seno, E. T., Rao, R. N. & Schonher, B. E. (1992) *Gene* **116**, 43–49.
- Pospiech, A. & Neumann, B. (1995) *Trends Genet.* **11**, 217–218.
- Flett, F., Mersinias, V. & Smith, C. P. (1997) *FEMS Microbiol. Lett.* **155**, 223–229.
- Goryshin, I. Y., Miller, J. A., Kil, Y. V., Lanzov, V. A. & Reznikoff, W. S. (1998) *Proc. Natl. Acad. Sci. USA* **95**, 10716–10721.
- Redenbach, M., Kieser, H. M., Denapaite, D., Eichner, A., Cullum, J., Kinashi, H. & Hopwood, D. A. (1996) *Mol. Microbiol.* **21**, 77–96.
- Davis, N. K. & Chater, K. F. (1990) *Mol. Microbiol.* **4**, 1679–1691.
- Horinouchi, S. & Beppu, T. (1985) *J. Bacteriol.* **162**, 406–412.
- Leblond, P. & Decaris, B. (1994) *FEMS Microbiol. Lett.* **123**, 225–232.
- Volff, J.-N. & Altenbuchner, J. (1998) *Mol. Microbiol.* **27**, 239–246.
- Volff, J.-N., Viell, P. & Altenbuchner, J. (1997) *Mol. Gen. Genet.* **253**, 761–765.
- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D. J. (1997) *Nucleic Acids Res.* **25**, 3389–3402.
- Maresca, B., Lambowitz, A. M., Kumar, V. B., Grant, G. A., Kobayashi, G. S. & Medoff, G. (1981) *Proc. Natl. Acad. Sci. USA* **78**, 4596–4600.
- Sacco, M., Maresca, B., Kumar, B. V., Kobayashi, G. S. & Medoff, G. (1981) *J. Bacteriol.* **146**, 117–120.
- Kumar, V., Maresca, B., Sacco, M., Goewert, R., Kobayashi, G. S. & Medoff, G. (1983) *Biochemistry* **22**, 762–768.
- Bateman, A., Birney, E., Durbin, R., Eddy, S. R., Howe, K. L. & Sonnhammer, E. L. L. (2000) *Nucleic Acids Res.* **28**, 263–266.
- Chater, K. F. & Merrick, M. J. (1976) in *Second International Symposium on the Genetics of Industrial Microorganisms*, ed. MacDonald, K. D. (Academic, London), pp. 583–593.
- Chater, K. F. & Bruton, C. J. (1983) *Gene* **26**, 67–78.
- Limauro, D., Avitabile, A., Cappellano, C., Puglia, A. M. & Bruni, C. B. (1990) *Gene* **90**, 31–41.
- Malpartida, F., Niemi, J., Navarrete, R. & Hopwood, D. A. (1990) *Gene* **93**, 91–99.
- Yu, T.-W. & Hopwood, D. A. (1995) *Microbiology* **141**, 2779–2791.
- Nodwell, J. R., McGovern, K. & Losick, R. (1996) *Mol. Microbiol.* **22**, 881–893.