

# The human pathogen *Pseudomonas aeruginosa* utilizes conserved virulence pathways to infect the social amoeba *Dictyostelium discoideum*

Stefan Pukatzki<sup>\*†</sup>, Richard H. Kessin<sup>‡</sup>, and John J. Mekalanos<sup>\*</sup>

<sup>\*</sup>Department of Microbiology and Molecular Genetics, Harvard Medical School, 200 Longwood Avenue, Boston, MA 02115; and <sup>‡</sup>Department of Anatomy and Cell Biology, College of Physicians and Surgeons, Columbia University, 630 West 168th Street, New York, NY 10032

Contributed by John J. Mekalanos, December 27, 2001

**Genetically accessible host models are useful for studying microbial pathogenesis because they offer the means to identify novel strategies that pathogens use to evade immune mechanisms, cause cellular injury, and induce disease. We have developed conditions under which the human pathogen *Pseudomonas aeruginosa* infects *Dictyostelium discoideum*, a genetically tractable eukaryotic organism. When *D. discoideum* is plated on nutrient agar plates with different *P. aeruginosa* strains, the bacteria form lawns on these plates with amoebae embedded in them. Virulent *P. aeruginosa* strains kill these amoebae and leave an intact bacterial lawn. A number of *P. aeruginosa* mutants have been identified that are avirulent in this assay. Amoebae feed on these bacteria and form plaques in their bacterial lawns. One avirulent mutant strain carries an insertional mutation in the *lasR* gene. LasR is a transcription factor that controls a number of virulence genes in a density-dependent fashion. Another class of avirulent *P. aeruginosa* mutants is defective in type III secretion. One mutant lacks the PscJ protein, a structural component of the secretion apparatus, suggesting that cytotoxins are injected into the *D. discoideum* cell. One of these cytotoxins is ExoU, and *exoU* mutants are avirulent toward *D. discoideum*. Complementation of the *lasR* and *exoU* mutations restores virulence. Therefore, *P. aeruginosa* uses conserved virulence pathways to kill *D. discoideum*.**

slime mold | type III secretion | host–pathogen interactions | LasR | ExoU

**P***seudomonas aeruginosa* is an opportunistic pathogen that causes life-threatening infections in individuals with compromised immune systems, such as cancer patients undergoing chemotherapy or patients with cystic fibrosis. Immunocompromised individuals are at high risk of becoming infected in a hospital setting where *P. aeruginosa* causes a variety of nosocomial infections, including pneumonia, urinary tract infections, surgical wound infections, and blood stream infections (for review, see ref. 1).

Cystic fibrosis patients, who carry mutations in both alleles of the cystic fibrosis transmembrane conductance regulator, are at high risk of developing chronic lung infections at some stage in their lives. Initially, *P. aeruginosa* colonizes the airways with other pathogens, like *Haemophilis influenzae* and *Staphylococcus aureus*. After a variable period, chronic lung disease develops in which the bacterial population consists almost exclusively of mucoid *P. aeruginosa* in the form of biofilms (2). Biofilms are structured communities of bacteria embedded in a polysaccharide matrix, and growth in this form renders *P. aeruginosa* innately resistant to antimicrobial treatment (3).

The fact that *P. aeruginosa* is resistant to treatment with antibiotics demands alternative strategies to treat infections with this pathogen. One such approach is to interfere with the interaction between *P. aeruginosa* and its host. Several experimental systems involving the plant *Arabidopsis thaliana*, the nematode *Caenorhabditis elegans*, the fruit fly *Drosophila melanogaster*, and mice have been used to study such host–pathogen interactions (for review, see ref. 4). To study the interaction of

*P. aeruginosa* and a host cell, we suggest the genetically tractable organism *Dictyostelium discoideum*. *D. discoideum* is a complex eukaryote that lives part of its life cycle as a unicellular amoeba. Free-living amoebae feed on bacteria by a phagocytic mechanism that resembles that of mammalian macrophages (for review, see ref. 5). Using *Dictyostelium* as a genetic model host takes advantage of the organism's ability to integrate complicated activities in a single cell, which includes responses to pathogens. It is this unicellular lifestyle that enables us to study the interaction between a host and a pathogen directly with only these two organisms present. *D. discoideum* is well suited for studying the host biology of pathogens, because it offers a variety of different genetic manipulations. Because *D. discoideum* is haploid, mutations are not masked by additional alleles, so that mutations, including recessive ones, are readily identified. The small size of the genome ( $\approx 34$  Mb) and the small amount of noncoding sequence allow us to readily saturate the genome with mutations. The entire genomic sequence will be known soon, and this will accelerate the identification of genes that emerge from any genetic screen for resistance to *P. aeruginosa*. *D. discoideum* has already been shown to be useful in studying the virulence mechanisms of the intracellular pathogen *Legionella pneumophila* (6).

A simple plating assay has been developed to study the interaction between *P. aeruginosa* and *D. discoideum*. *P. aeruginosa* formed lawns on these plates with amoebae embedded in them. Virulent *P. aeruginosa* strains kill these amoebae. If a mutation causes a particular strain to be avirulent toward *D. discoideum*, the amoebae feed on these bacteria and form plaques in bacterial lawns after a few days. We have identified a number of *P. aeruginosa* mutants that are avirulent in this assay. Analysis of these avirulent mutants suggests that *P. aeruginosa* utilizes at least two major virulence pathways, namely quorum-sensing mediated virulence and type III secretion of virulence factors, to infect amoebae. We propose that *D. discoideum* is a genetically tractable system that can be used to study the host biology of *P. aeruginosa*.

## Materials and Methods

**Strains and Culture Conditions.** *Dictyostelium* strain AX3 was used in all experiments (7). AX3 was grown in liquid HL/5 cultures or in lawns of *Klebsiella aerogenes* on SM/5 plates, as described by Sussman (8). In some of the experiments, low-nutrient plates were used to limit bacterial growth. SM/25, SM/50, and SM/100-plates contained 1/25, 1/50, or 1/100 of a SM-stock solution [1% glucose/1% Bacto peptone (Difco)/0.1% Bacto yeast extract (Difco)/4.2 mM MgSO<sub>4</sub>], respectively. Solutions

<sup>†</sup>To whom reprint requests should be addressed. E-mail: stefan.pukatzki@hms.harvard.edu.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

were buffered to pH 6.5 by the addition of 13.2 mM  $\text{KH}_2\text{PO}_4$  and 6.9 mM  $\text{K}_2\text{HPO}_4$ .

The *lasR*-mutant 12A1, the *phenA/phenB*-mutant  $\Delta\text{phenA}/\text{phenB}$ , the *TnphoA*-mutant 8C12, as well as their parent strain UCB-PP PA14, were kind gifts from Fred Ausubel (Massachusetts General Hospital, Boston) (9–11). Strains SUP16 and SUP17 were generated by introducing plasmid pUCP18 (12) and plasmid pLasR-PUH (see below) into *lasR*-mutant 12A1. A *rhlA* mutant of PA14 was constructed with knockout-plasmid pEX100T-*rhlA*::Gm by using the site-specific insertional mutagenesis strategy of Schweizer and Hoang (13, 14).

The *exoU*-mutant PA103::*exoU* (mut 8) (15), the *pscJ*-mutant PA103::*pscJ* (mut N) (15), and their isogenic parent PA103 (16) were gifts from Joanne Engel (University of California, San Francisco). The *exoU* mutation was complemented by introducing plasmid pAH807 (pLAFRSK1 with a 3.8-kb genomic fragment containing *exoU*) (17); the resulting strain was also a kind gift from Joanne Engel.

**Creation of pLasR-PUH.** To complement the *lasR* mutation in *P. aeruginosa* strain 12A1, we designed a plasmid that provides high expression levels of *lasR*. The *lasR* gene was recovered from plasmid pKDT17 (18) as a 0.8-kb *EcoRI/HindIII* fragment and subcloned in front of the *lac* promoter of *EcoRI/HindIII*-digested pUCP18 (12).

**Plate Killing Assay.** Bacteria were grown in Luria broth for 16 h, pelleted by centrifugation ( $1,600 \times g$ ; 15 min), washed once, and resuspended in SorC (16.7 mM  $\text{Na}_2\text{H/KH}_2\text{PO}_4/50 \mu\text{M CaCl}_2$ , pH 6.0) at a final optical density of 5.5 at 600 nm. *D. discoideum* cells from midlogarithmic cultures were collected by centrifugation ( $1,000 \times g$ ; 4 min), washed once with SorC (8), and added to the bacterial suspensions at a final concentration of  $5 \times 10^2$  cells/ml suspension; 0.2 ml of this mixture was plated out on SM/5 plates and allowed to dry under a sterile flow of air. Plates were incubated for 3–5 days and examined for plaques formed by *Dictyostelium* amoebae.

**Elastase and Pyocyanin Assays.** Elastolytic activity was determined as described by Gambello and Iglewski (19). Pyocyanin levels in bacterial cultures were determined as described by Essar *et al.* (20).

**Gentamicin Protection Assay.** To test whether *P. aeruginosa* is internalized and digested by *D. discoideum*, a modification of the gentamicin protection assay was used (21). *D. discoideum* cells from midlogarithmic cultures were collected by centrifugation ( $1,000 \times g$ ; 4 min), washed once with fresh HL/5 (8), and resuspended in HL/5 at a concentration of  $10^6$  cells/ml. Aliquots of 1.0 ml were added to 24-well tissue culture dishes (Falcon) and incubated for 60 min at 22°C to allow cells to adhere to the plastic surface. Bacteria were grown in Luria broth for 16 h, pelleted by centrifugation ( $1,600 \times g$ ; 15 min), and resuspended in HL/5. Bacteria were added to the tissue culture wells at a final optical density of 0.1 at 600 nm (multiplicity of infection = 100). Infection was initiated by centrifugation ( $750 \times g$ ; 10 min). Thirty minutes after the initiation of infection, cells were washed twice and resuspended in SorC (16.7 mM  $\text{Na}_2\text{H/KH}_2\text{PO}_4/50 \mu\text{M CaCl}_2$ , pH 6.0) containing 400  $\mu\text{g/ml}$  of gentamicin to kill all bacteria that were not ingested by the amoebae. To determine the number of viable bacteria inside the amoebae at various time points, cells were washed twice with SorC and lysed in SorC containing 0.05% Triton X-100. The lysates were diluted and plated on Luria broth agar plates. After an incubation at 37°C for 16 h, the colony-forming units were counted.



**Fig. 1.** *D. discoideum* cells do not form plaques in lawns of *P. aeruginosa* strain PA14. *D. discoideum* cells (AX3) were plated on SM/5 with *K. aerogenes* (Left) and *P. aeruginosa* strains PA14 (Right) at a density of  $\approx 100$  *D. discoideum* cells/plate. After 3 days, *D. discoideum* plaques appeared on plates with *K. aerogenes*. No plaques were formed on plates with the virulent *P. aeruginosa* strain.


## Results

***D. discoideum* Is Susceptible to Infection with *P. aeruginosa*.** To test whether *P. aeruginosa* could infect *D. discoideum*, the pathogenicity of *P. aeruginosa* toward *D. discoideum* cells was determined by using a simple plating assay. *D. discoideum* cells were plated on nutrient agar plates with nonpathogenic *Klebsiella aerogenes* or virulent *P. aeruginosa* strain PA14. Over the course of a few days, the bacteria formed lawns on these plates with amoebae embedded in them. As shown in Fig. 1, *D. discoideum* feeds on *K. aerogenes* and forms plaques that become readily apparent after a few days. *D. discoideum* does not form any plaques on the lawns of *P. aeruginosa* strain PA14. This experiment does not distinguish between killing and growth inhibition, but we have shown that the bacteria kill the amoebae (see below). *P. aeruginosa* therefore has the capacity to kill *D. discoideum* under these conditions.

**Virulence Factors Controlled by Quorum Sensing Kill *D. discoideum*.** A number of virulence factors of *P. aeruginosa* are produced only when bacteria are present at high titers. This “quorum-sensing mechanism” is thought to be an important regulator of pathogenesis, because it guarantees that certain virulence factors are made only when *P. aeruginosa* has reached a crucial density and can launch a successful infection (22).

We chose to investigate the role of the quorum-sensing mechanism by testing the virulence of quorum-sensing mutants of *P. aeruginosa* strain PA14 (11). PA14 is a multihost pathogen that has been used widely for studying the molecular basis of biofilm formation (9–11, 23). As shown in Fig. 2 *Top*, *D. discoideum* is unable to form plaques in lawns of parental strain PA14 but forms plaques in lawns of the isogenic *lasR*-mutant 12A1. Mutant strain 12A1 carries a *TnphoA* transposon in the ORF of *lasR* (9). *lasR* encodes the main transcriptional regulator of the *las* quorum-sensing pathway, and the *lasR*-mutant 12A1 has been shown to be avirulent toward *C. elegans* (9).

To test whether the ability of *Dictyostelium* to form plaques on strain 12A1 is due to the removal of LasR, we carried out a plasmid complementation experiment. We constructed a plasmid, pLasR-PUH, which carries *lasR* fused to the *lacZ* promoter. Plasmid pLasR-PUH and vector control pUCP18 were introduced into *lasR*-mutant 12A1, resulting in strains SUP17 and SUP16, respectively. To ensure that the introduction of pLasR-PUH restores LasR function, we determined the activity of the LasR-controlled *lasB* gene product, elastase. As shown in Fig. 2 *Middle*, *lasR*-mutant 12A1 and vector control strain SUP16 show low elastase activity, compared with wild-type PA14. The complemented *lasR* strain SUP17 shows elastase levels comparable



**Fig. 2.** Complementation of the *lasR* mutation restores LasR function and virulence but does not affect synthesis of the pigment pyocyanin. (Top) *D. discoideum* cells (AX3) were plated on nutrient agar (SM/5) with *P. aeruginosa* strain PA14, the isogenic *lasR*-mutant 12A1, complemented strains SUP16 ( $\Delta lasR$  with pUCP18) and SUP17 ( $\Delta lasR$  with plasR-PUH), and the pyocyanin-deficient strain  $\Delta phenA/\Delta phenB$ . Plates were incubated at 22°C for 5 days, and virulence was assessed by determining the number of *Dictyostelium* plaques. (Middle) Elastase activity of supernatants from overnight cultures were assayed by using elastin Congo red (ECR) as the substrate. Data represent the means of duplicate elastase assays, and activity is expressed as the activity relative to wild-type PA14 (=100%)  $\pm$  SD<sup>n-1</sup>. (Bottom) The ability of each strain to secrete the pigment pyocyanin was determined by assaying the amount of pyocyanin present in overnight cultures of each strain. Pyocyanin was extracted from supernatants and assayed as described in (20). Data represent the means of duplicate assays, and amounts of secreted pyocyanin are expressed as micrograms/milliliter culture medium  $\pm$  SD<sup>n-1</sup>.

to PA14, indicating that the function of LasR is restored in this strain. We then determined the virulence of these strains in our plaque assay. *Dictyostelium* formed plaques in lawns of SUP16, but only a few petite plaques could be identified in lawns of SUP17 (see Fig. 2 Top). Complementation of *lasR* therefore restores virulence that is comparable to wild-type strain PA14. For various reasons, elastase is unlikely to be a virulence factor in this host–pathogen system (see Discussion).

Work with *C. elegans* as a model host showed that *P. aeruginosa* secretes a pigment, pyocyanin, that exerts its toxic effect on eukaryotic cells through reactive oxygen species (10). Because pyocyanin synthesis is controlled by quorum sensing, we investigated whether *P. aeruginosa*-induced killing of *Dictyostelium* was mediated by pyocyanin. First, we determined the amount of pyocyanin that was secreted by PA14 and its mutants during stationary phase. Pyocyanin ( $\approx 6\text{--}7$   $\mu\text{g/ml}$ ) were found in cultures of wild-type PA14 (see Fig. 2 Bottom). The isogenic *lasR*-mutant 12A1 produced about one-third of this amount of pyocyanin. Reduced pyocyanin production is expected in a *lasR* mutant because LasR directly controls the expression of RhIR, a transcription factor that is involved in pyocyanin production (24). Overexpression of *lasR* from plasmid pLasR-PUH did not restore pyocyanin production in strain SUP17. We speculate that this is because high levels of intracellular LasR inhibit the activity of the transcription factor RhIR at a posttranslational level (25). These results showed that virulent strain SUP17 and avirulent strains 12A1 and SUP16 release similar amounts of pyocyanin into the culture medium. Furthermore, a PA14 mutant that is unable to secrete pyocyanin because of a deletion in the *phenA* and *phenB* genes is still virulent in our plaque assay (see Fig. 2 Top) (10). The addition of purified pyocyanin to axenic cultures of *D. discoideum* at a concentration of 7  $\mu\text{g/ml}$ , the highest amount of pyocyanin we ever found in *P. aeruginosa* supernatants did not affect the viability of growing amoebae


(data not shown). Pyocyanin is, therefore, unlikely to be responsible for killing *D. discoideum* amoebae.

Another class of virulence factors that is controlled by the *lasR* quorum-sensing mechanism are the rhamnolipids. Rhamnolipids are biosurfactants that lyse host cells by disrupting their cell membranes (26). We investigated whether the avirulent phenotype of the *lasR*-mutant 12A1 was due to a reduction in rhamnolipid secretion. We measured the amounts of extracellular rhamnolipids from *P. aeruginosa* cultures in stationary phase and found no significant differences among the *lasR*-mutant 12A1, the plasmid-complemented strain SUP17, and wild-type PA14 (data not shown). To determine the role of rhamnolipids in the plaque assay, we tested the virulence of a PA14 mutant defective in the *rhlA* gene. *RhlA* encodes a subunit of rhamnosyltransferase 1, an essential enzyme in the synthesis of rhamnolipids, and mutations in *rhlA* result in the inability to secrete any rhamnolipids (13). We constructed a *rhlA* mutant of *P. aeruginosa* PA14 by using the site-specific insertional mutagenesis strategy of Schweizer and Hoang (14). As expected, the resulting mutant, PA14 $\Delta rhlA$ , showed a defect in rhamnolipid synthesis (data not shown). We then tested the virulence of this strain in our plaque assay. *D. discoideum* amoebae were plated on nutrient agar plates in combination with *K. aerogenes* or *P. aeruginosa* strains PA14 and PA14 $\Delta rhlA$ . *D. discoideum* formed plaques in lawns of *K. aerogenes* but was unable to do so in lawns of PA14 and PA14 $\Delta rhlA$ . Therefore, rhamnolipids do not play a role in the virulence of *P. aeruginosa* on *D. discoideum* (data not shown).

#### Plaque Formation on Strain 12A1 Is Not Due to Increased Phagocytosis.


*D. discoideum* may form plaques in lawns of the *lasR*-mutant 12A1 but not in lawns of its isogenic parent PA14, because amoebae phagocytose strain 12A1 more efficiently. For example, strain 12A1 might be taken up at an increased rate if the loss of LasR had affected a virulence mechanism that inhibits phagocytosis by the host cell. To determine whether this is the case, we applied a modified gentamicin protection protocol to measure the uptake rates of various *P. aeruginosa* strains, including strains 12A1 and PA14 (21). *D. discoideum* and *P. aeruginosa* were cocultured for 30 min, and any bacteria that were not taken up by the amoebae were killed by the addition of gentamicin. The fate of intracellular bacteria was monitored over a period of 4 h. As shown in Fig. 3, avirulent *K. aerogenes* is readily phagocytosed by the amoebae. Once internalized, *K. aerogenes* is digested by the endosomal pathway (27). The virulent *P. aeruginosa* strain PA14 is phagocytosed at a much lower rate ( $\approx 100$ -fold less compared with *K. aerogenes*), but once internalized, PA14 is also rapidly digested. The same is true for another virulent strain, SUP17, which carries the *lasR* gene *in trans* on a high-copy plasmid. Uptake and digestion rates are comparable to those of PA14. The avirulent *lasR*-mutant 12A1, although ingested at an even lower rate, is degraded at rates comparable to those of PA14 and SUP17. Therefore, the ability of *D. discoideum* to form plaques in lawns of strain 12A1 is not because of increased phagocytosis of this particular mutant. The low phagocytosis rate of strain 12A1 is reflected in reduced plaque size. *D. discoideum* forms plaques in lawns of 12A1 that are approximately one-tenth the size of those formed in lawns of *K. aerogenes* (data not shown). The gentamicin protection experiment demonstrates that *D. discoideum* has the capacity to phagocytose all *P. aeruginosa* strains tested. Therefore, the ability of *D. discoideum* to form plaques in lawns of 12A1 is because of the loss of a virulence mechanism that does not involve the inhibition of phagocytosis.

When exposed to low titers of bacterial cells in a liquid medium, *D. discoideum* amoebae ingested all *P. aeruginosa* strains by phagocytosis in our gentamicin-protection experiment. We asked whether the same is true on plates when bacteria




**Fig. 3.** Uptake of *P. aeruginosa* by *D. discoideum*. *D. discoideum* cells were placed in tissue culture wells at  $10^6$  cells/ml. Cells were infected with bacteria at a multiplicity of infection of approximately 100:1. Cultures were incubated at 22°C for 30 min, at which time gentamicin was added to kill all extracellular bacteria. Amoebae were collected at indicated time points, lysed, and plated on nutrient agar plates to determine the colony-forming units (cfu) and the number of internalized bacteria per well is expressed as cfu/ml  $\pm$  SD<sup>n-1</sup>.

were plated at low titers. Bacteria and amoebae were plated on agar plates with decreasing nutrient content to control the thickness of the bacterial layer, which is proportional to the nutrient concentration in the agar. *D. discoideum* plaques were scored after 4 days (see Fig. 4). On SM/25 and SM/50 plates, which contain one-fifth and one-tenth of the nutrient concentration of SM/5, respectively, plaques emerged in lawns of the



**Fig. 4.** *D. discoideum* forms plaques in thin lawns of *P. aeruginosa* PA14. *D. discoideum* cells (AX3) were plated with *P. aeruginosa* strains PA14 (wild type) and PA14 mutants 8C12 and 12A1 ( $\Delta$ lasR) on nutrient agar plates with decreasing nutrient content. Agar plates contained 1/25 (SM/25), 1/50 (SM/50), and 1/100 (SM/100) of the nutrient content in SM plates. After 4 days at 22°C, *D. discoideum* formed plaques in lawns of 12A1 on SM/25 and plates with lower nutrient content, 8C12 and PA14 allowed plaque formation only on SM/100 plates.




**Fig. 5.** Complementation of the *exoU* mutation restores virulence. *D. discoideum* cells (AX3) were plated with different bacterial associates on nutrient agar plates (SM/5) at a density of approximately 40 amoebae/plate. Emerging *D. discoideum* plaques on each plate were scored after 5 days. Virulence is expressed as the average number of emerging plaques per plate, and error bars represent the standard deviation of five replicates. The virulence of PA103, as well as isogenic mutants PA103::pscJ ( $\Delta$ pscJ), PA103::exoU ( $\Delta$ exoU), PA103::exoU+pLAFRSK1 ( $\Delta$ exoU with plasmid control pLAFRSK1), and PA103::exoU+pAH807 [ $\Delta$ exoU with plasmid pAH807 (*exoU*+/*spcU*+)] was tested in this assay. As a reference for plating efficiency, AX3 was also cultured with *K. aerogenes*.

*lasR*-mutant 12A1. No plaques could be seen in lawns of wild-type PA14 and the isogenic mutant 8C12 on SM/25 and SM/50 plates. Strain 8C12 is identical to 12A1, except that the *TnphoA* transposon is in another gene, which does not affect virulence toward amoebae (10). When nutrient content was reduced 20-fold compared with SM/5, plaques emerged in all lawns, including PA14. Therefore, *D. discoideum* amoebae are able to feed on virulent *P. aeruginosa* when bacteria are present at low abundance. The plaques formed in lawns of PA14 and 8C12 are smaller in size compared with those in lawns of 12A1. Reduced plaque size might be a result of residual virulence that PA14 and 8C12 exhibit toward *D. discoideum* on low-nutrient plates (see Fig. 4).

**The Acute Cytotoxin ExoU Kills Dictyostelium Amoebae.** *P. aeruginosa* utilizes virulence factors that are injected into the host cell through a “type III secretion apparatus” (for review, see ref. 28). This apparatus is a multiprotein complex that is embedded in the outer membrane of the bacterium and that inserts a pilus-like structure into the host cell. This structure allows cytotoxins to be transferred from the bacterium to the host cell. Four toxins that are injected in this way, ExoS, ExoT, ExoU, and ExoY, have been identified (15, 29, 30).

We examined the role of type III secretion of toxins in the observed virulence of *P. aeruginosa* toward *D. discoideum* by testing the virulence of the clinical *P. aeruginosa* isolate, PA103. Previous work demonstrated that PA103 uses type III secretion to kill macrophages and epithelial cells *in vitro* (31). When amoebae were plated on nutrient agar plates with lawns of PA103, no *Dictyostelium* plaques emerged (see Figs. 5 and 6A). To address the possibility that the amoebae simply cannot eat PA103 and therefore cannot grow but remain alive, we monitored the fate of amoebae in lawns of virulent PA103 by recovering amoebae from these lawns after an incubation period of 24 h. Recovered amoebae were plated with *K. aerogenes* on nutrient agar plates supplemented with 200  $\mu$ g/ml of kanamycin that kill *P. aeruginosa* but not *K. aerogenes*. Only about 2.5% of the amoebae that were plated with PA103 could be recovered



**Fig. 6.** Expression of ExoU in *P. aeruginosa* PA103 prohibits plaque formation by *D. discoideum*. *D. discoideum* cells (AX3) were plated on SM/5 with *P. aeruginosa* strains PA103 (A), PA103::*exoU* ( $\Delta$ *exoU*) (B), and PA103::*exoU* + pAH807 ( $\Delta$ *exoU* with plasmid pAH807 [*exoU*<sup>+</sup>/*spcU*<sup>+</sup>]) (C) at a density of  $\approx$ 100 *D. discoideum* cells/plate. Plaques were scored after 4 days of incubation at 22°C.

from these lawns (data not shown), whereas more than 95% of amoebae that were plated on agar plates without a bacterial associate survived such starvation treatment. Thus, the loss of amoebae in lawns of PA103 is because of killing rather than starvation. The same is true for PA14 (data not shown).

To test whether killing is because of the translocation of toxins, we examined the virulence of a mutant of PA103 that carries a mutation in the *pscJ* gene and is therefore defective in type III secretion (15). *pscJ* exhibits 83% similarity to the type III secretion apparatus component YscJ from *Yersinia enterocolitica* and is essential for the export of toxins (32). The mutation in *pscJ* prohibits export of the four known cytotoxins by *P. aeruginosa* strain PA103 (15, 33). *Dictyostelium* amoebae formed plaques in lawns of the *pscJ* mutant, suggesting that during infections with wild-type strain PA103, cytotoxins are injected into the *D. discoideum* host cells (see Fig. 5).

Knowing that a type III secretion apparatus was involved, we asked which virulence factors were responsible for the killing of *Dictyostelium* amoebae. Because strain PA103 does not produce cytotoxins ExoY and ExoS, we determined the virulence of a PA103 mutant that carries a *Tn5*-transposon insertion in the *exoU* gene (15, 30, 34). This *exoU* mutant is less virulent toward *D. discoideum* than PA103, allowing *D. discoideum* to form plaques (see Figs. 5 and 6B).

The *exoU* operon contains the *exoU* and *spcU* genes. *SpcU* (for specific *Pseudomonas* chaperone for ExoU) encodes a chaperone that is necessary for proper translocation of ExoU (35). The transposon insertion at the 5' end of the gene, 125 bp downstream from the translational start site, results in a polar mutation that disrupts both *exoU* and *spcU* in strain PA103 $\Delta$ *exoU*. To confirm that the loss of virulence is exclusively because of the inability of this strain to export active ExoU, we complemented the *exoU* mutation by introduction of plasmid pAH807, which carries both the *exoU* and *spcU* genes (17). If ExoU is the cause of lethality, then complementation of the mutation should prevent growth of *D. discoideum* on lawns of the complemented mutant. *D. discoideum* amoebae could not form plaques in the complemented *exoU* mutant (see Figs. 5 and 6C) but were able to form plaques in lawns of the *exoU* mutant that carries the empty plasmid control (see Fig. 5). Thus, ExoU is responsible for killing *D. discoideum*.

*D. discoideum* is able to form plaques in thin lawns of virulent *P. aeruginosa* strains. Any genetic manipulation of *P. aeruginosa* that results in a growth defect could lead to the formation of thin bacterial lawns on rich-medium plates, allowing *D. discoideum* to form plaques. To determine whether the ability of *D. discoideum* to feed on particular mutants of *P. aeruginosa* strain PA103 was because of reduced growth rates of these strains, we examined the growth properties of the mutant bacteria. Liquid SM/5

medium, the same medium that was used to prepare nutrient agar plates for the plaque assay, was inoculated with various PA103 mutants, and their growth at 22°C was followed over 30 h. The growth properties of strains defective in the *exoU* or *pscJ* genes did not differ from their isogenic parent PA103. Introduction of plasmids pAH807 (*ExoU*<sup>+</sup>/*SpcU*<sup>+</sup>) or pLAFRSKI (plasmid control) into the *exoU* mutant also did not affect growth (data not shown). We can therefore exclude the possibility that the *exoU* and *pscJ* mutants are avirulent, because they do not reach high enough titers on nutrient agar plates to kill *D. discoideum*. Virulence of the complemented *exoU* strain is not because of an increased growth rate.

## Discussion

The results presented here establish a host–pathogen interaction between the human pathogen *P. aeruginosa* and the social amoeba *D. discoideum*. When the two organisms are coincubated on nutrient agar plates, *P. aeruginosa* is capable of killing *D. discoideum*. Defined mutants of *P. aeruginosa*, however, allow *D. discoideum* to feed on these bacteria and to form plaques in the bacterial lawns. These mutants are impaired in two conserved virulence pathways: quorum-sensing-mediated virulence and type III secretion of cytotoxins.

Quorum sensing is thought to be an important determinant of *P. aeruginosa* virulence, because it guarantees that the organism expresses virulence factors only when it is present at high numbers. A mutant in the *lasR* gene, which codes for the transcriptional activator of the *lasR* quorum-sensing mechanism, is less virulent toward *D. discoideum* and allows plaque formation. Complementation of the *lasR* mutation restores virulence, as *D. discoideum* amoebae are not able to form plaques in lawns of this complemented strain. Thus, *P. aeruginosa* predation of *D. discoideum* depends on LasR, which likely regulates expression of a downstream effector gene through the quorum-sensing pathway. This LasR-controlled virulence factor remains to be identified, but we considered rhamnolipids, phenazines, and elastase as possible candidates. We directly examined the contribution of rhamnolipids and phenazines to *D. discoideum* killing but found that these factors were not involved. Furthermore, addition of purified pyocyanin did not affect the viability of axenically growing amoebae. The fact that *D. discoideum* is resistant to pyocyanin at concentrations that kill mammalian cells is not surprising. Pyocyanin is a phenazine that applies oxidative stress to eukaryotic cells. *D. discoideum* is a soil organism that normally encounters a variety of different oxidative radicals and has developed a wide range of mechanisms to deactivate such radicals (5). We are currently examining the role of elastase in the *P. aeruginosa*–*D. discoideum* interaction.

Besides quorum-sensing mediated virulence, *P. aeruginosa* utilizes a type III secretion mechanism to kill *D. discoideum*. Cytotoxin ExoU contributes to the pathogenicity of *P. aeruginosa* PA103 toward *D. discoideum*, because PA103 mutants that are defective in type III secretion, or production of cytotoxin ExoU are avirulent toward *D. discoideum*. ExoU is a 71-kDa protein that, once translocated into a host cell, causes lysis of the host cell within a few hours (17, 31, 36, 37). Isogenic mutants that do not produce or secrete ExoU are defective in virulence in a mouse model of pneumonia (17, 36). The mechanism by which ExoU induces rapid lysis is not yet known. The use of *D. discoideum* as a host model may lead to the identification of cellular host components that are targeted by ExoU.

Most *P. aeruginosa* isolates from cystic fibrosis patients lack the *exoU* gene or are unable to secrete the toxin, whereas a higher proportion of strains isolated from corneal infections are reported to contain ExoU (38–40). This finding suggests that *P. aeruginosa* strains that use ExoU as a virulence factor are readily cleared from the lungs of cystic fibrosis patients. The ExoU cytotoxin may induce an unusually strong host response that

results in rapid clearing from the lungs. Identification of the host targets of ExoU may reveal a signaling pathway that can be stimulated with existing or novel drugs. Increasing the host response in such a manner may help cystic fibrosis patients to clear infections with *P. aeruginosa* that lack ExoU.

The work presented here suggests that *D. discoideum* represents a permissive host for *P. aeruginosa* infections. Using *D. discoideum* as a genetically tractable model host takes advantage of the organism's ability to integrate complicated activities in a single cell, which includes responses to pathogens. A number of different genetic manipulations, including insertional mutagenesis, chemical mutagenesis, complementation, and cloning, are available to identify mutations that confer resistance to *Pseudomonas* infections (for a review of available methods, see ref. 5). For example, insertional mutagenesis by restriction-enzyme-mediated integration (REMI) may introduce null mutations that result in the genetic removal of proteins from cells (41). If such a protein is a host component that is targeted by a particular virulence factor, its removal may confer resistance to the patho-

gen. Identification of the affected gene is straightforward because the gene can be readily cloned because of the insertion of a selectable plasmid during the REMI procedure. With such a genetic approach, two major virulence pathways, quorum-sensing-mediated virulence and type III secretion of cytotoxins, can be studied in more detail from the host's perspective.

We are grateful to Herbert Ennis, Jakob Franke, Felipe B. Manalo, Grant P. Otto, Mary Wu, and members of the Mekalanos laboratory for helpful discussions, and to Su Chiang for critically reading the manuscript. We thank Lynne Garrity-Ryan (University of California, San Francisco) and Joanne Engel for *P. aeruginosa* strain PA103 and its isogenic mutants, as well as Fred Ausubel for *P. aeruginosa* strain PA14 and its isogenic mutants. We thank Bradley Britigan (University of Iowa) and Barry Schniepp (Jeneil Biosurfactant Co., Saukville, WI) for their kind gifts of purified pyocyanin and rhamnolipids, respectively. We also thank Urs Ochsner (University of Colorado Health Sciences Center) for knockout plasmid pEX100T-rhLA::Gm. This work was supported by National Institutes of Health Grant GM33136-20 (to R.H.K.) and Grant AI26289 (to J.J.M.), as well as by a grant from the Cystic Fibrosis Foundation (to S.U.P.).

- Deretic, V. (2000) in *Persistent Bacterial Infections*, eds. Nataro, J. P., Blaser, M. J. & Cunningham-Rundles, S. (Am. Soc. Microbiol., Washington, DC), pp. 305-326.
- Govan, J. R. & Deretic, V. (1996) *Microbiol. Rev.* **60**, 539-574.
- Singh, P. K., Schaefer, A. L., Parsek, M. R., Moninger, T. O., Welsh, M. J. & Greenberg, E. P. (2000) *Nature (London)* **407**, 762-764.
- Mahajan-Miklos, S., Rahme, L. G. & Ausubel, F. M. (2000) *Mol. Microbiol.* **37**, 981-988.
- Kessin, R. H. (2001) *Dictyostelium—Evolution, Cell Biology, and the Development of Multicellularity* (Cambridge Univ. Press, Cambridge, U.K.).
- Solomon, J. M., Rupper, A., Cardelli, J. A. & Isberg, R. R. (2000) *Infect. Immun.* **68**, 2939-2947.
- Loomis, W. F. (1971) *Exp. Cell Res.* **64**, 484-486.
- Sussman, M. (1987) *Methods Cell Biol.* **28**, 9-29.
- Tan, M. W., Rahme, L. G., Sternberg, J. A., Tompkins, R. G. & Ausubel, F. M. (1999) *Proc. Natl. Acad. Sci. USA* **96**, 2408-2413.
- Mahajan-Miklos, S., Tan, M. W., Rahme, L. G. & Ausubel, F. M. (1999) *Cell* **96**, 47-56.
- Rahme, L. G., Stevens, E. J., Wolfort, S. F., Shao, J., Tompkins, R. G. & Ausubel, F. M. (1995) *Science* **268**, 1899-1902.
- Schweizer, H. P. (1991) *Gene* **97**, 109-121.
- Rahim, R., Ochsner, U. A., Olvera, C., Graninger, M., Messner, P., Lam, J. S. & Soberon-Chavez, G. (2001) *Mol. Microbiol.* **40**, 708-718.
- Schweizer, H. P. & Hoang, T. T. (1995) *Gene* **158**, 15-22.
- Kang, P. J., Hauser, A. R., Apodaca, G., Fleiszig, S. M., Wiener-Kronish, J., Mostov, K. & Engel, J. N. (1997) *Mol. Microbiol.* **24**, 1249-1262.
- Apodaca, G., Bomsel, M., Lindstedt, R., Engel, J., Frank, D., Mostov, K. E. & Wiener-Kronish, J. (1995) *Infect. Immun.* **63**, 1541-1551.
- Hauser, A. R., Kang, P. J. & Engel, J. N. (1998) *Mol. Microbiol.* **27**, 807-818.
- Pearson, J. P., Gray, K. M., Passador, L., Tucker, K. D., Eberhard, A., Iglewski, B. H. & Greenberg, E. P. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 197-201.
- Gambello, M. J. & Iglewski, B. H. (1991) *J. Bacteriol.* **173**, 3000-3009.
- Essar, D. W., Eberly, L., Hadero, A. & Crawford, I. P. (1990) *J. Bacteriol.* **172**, 884-900.
- Mounier, J., Vasselon, T., Helliou, R., Lesourd, M. & Sansonetti, P. J. (1992) *Infect. Immun.* **60**, 237-248.
- Fuqua, C., Winans, S. C. & Greenberg, E. P. (1996) *Annu. Rev. Microbiol.* **50**, 727-551.
- O'Toole, G. A. & Kolter, R. (1998) *Mol. Microbiol.* **30**, 295-304.
- Ochsner, U. A., Koch, A. K., Fiechter, A. & Reiser, J. (1994) *J. Bacteriol.* **176**, 2044-2054.
- Pesci, E. C., Pearson, J. P., Seed, P. C. & Iglewski, B. H. (1997) *J. Bacteriol.* **179**, 3127-3132.
- Van Delden, C. & Iglewski, B. H. (1998) *Emerg. Infect. Dis.* **4**, 551-560.
- Cornillon, S., Pech, E., Benghezal, M., Ravanel, K., Gaynor, E., Letourneur, F., Bruckert, F. & Cosson, P. (2000) *J. Biol. Chem.* **275**, 34287-34292.
- Galan, J. E. & Collmer, A. (1999) *Science* **284**, 1322-1328.
- McGuffee, E. M., Frank, D. W., Vincent, T. S. & Olson, J. C. (1998) *Infect. Immun.* **66**, 2607-2613.
- Yahr, T. L., Vallis, A. J., Hancock, M. K., Barbieri, J. T. & Frank, D. W. (1998) *Proc. Natl. Acad. Sci. USA* **95**, 13899-13904.
- Hauser, A. R. & Engel, J. N. (1999) *Infect. Immun.* **67**, 5530-5537.
- Michiels, T., Vanooteghem, J. C., Lambert de Rouvroit, C., China, B., Gustin, A., Boudry, P. & Cornelis, G. R. (1991) *J. Bacteriol.* **173**, 4994-5009.
- Hauser, A. R., Fleiszig, S., Kang, P. J., Mostov, K. & Engel, J. N. (1998) *Infect. Immun.* **66**, 1413-1420.
- Yahr, T. L., Mende-Mueller, L. M., Friese, M. B. & Frank, D. W. (1997) *J. Bacteriol.* **179**, 7165-7168.
- Finck-Barbancon, V., Yahr, T. L. & Frank, D. W. (1998) *J. Bacteriol.* **180**, 6224-6231.
- Finck-Barbancon, V., Goranson, J., Zhu, L., Sawa, T., Wiener-Kronish, J. P., Fleiszig, S. M., Wu, C., Mende-Mueller, L. & Frank, D. W. (1997) *Mol. Microbiol.* **25**, 547-557.
- Evans, D. J., Frank, D. W., Finck-Barbancon, V., Wu, C. & Fleiszig, S. M. (1998) *Infect. Immun.* **66**, 1453-1459.
- Dacheux, D., Toussaint, B., Richard, M., Brochier, G., Croize, J. & Attree, I. (2000) *Infect. Immun.* **68**, 2916-2924.
- Feltman, H., Schulert, G., Khan, S., Jain, M., Peterson, L. & Hauser, A. R. (2001) *Microbiology* **147**, 2659-2669.
- Lomholt, J. A., Poulsen, K. & Kilian, M. (2001) *Infect. Immun.* **69**, 6284-6295.
- Kuspa, A. & Loomis, W. F. (1992) *Proc. Natl. Acad. Sci. USA* **89**, 8803-8807.