

The origin of malignant malaria

Stephen M. Rich^{a,1}, Fabian H. Leendertz^b, Guang Xu^a, Matthew LeBreton^c, Cyrille F. Djoko^{c,d}, Makoah N. Aminake^d, Eric E. Takang^c, Joseph L. D. Diffo^c, Brian L. Pike^c, Benjamin M. Rosenthal^e, Pierre Formenty^f, Christophe Boesch^g, Francisco J. Ayala^{h,1}, and Nathan D. Wolfe^{c,i,1}

^aLaboratory of Medical Zoology, Division of Entomology (PSIS), University of Massachusetts, Amherst, MA 01003; ^bDepartment Emerging Zoonoses, Robert Koch Institute, Nordufer 20, D-13353 Berlin, Germany; ^cGlobal Viral Forecasting Initiative, San Francisco, CA 94104; ^dBiotechnology Centre, University of Yaounde I, Yaounde, Cameroon; ^eAnimal Parasitic Diseases Laboratory, Agricultural Research Service, US Department of Agriculture, Beltsville, MD 20705; ^fEbola Taï Forest Project, World Health Organization (WHO), WHO Office in Abidjan, Côte d'Ivoire; ^gDepartment of Primatology, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, D-04103 Leipzig, Germany; ^hDepartment of Ecology and Evolutionary Biology, University of California, Irvine, CA, 92697; and ⁱProgram in Human Biology, Stanford University, Stanford, CA 94305

Contributed by Francisco J. Ayala, July 13, 2009 (sent for review June 29, 2009)

Plasmodium falciparum, the causative agent of malignant malaria, is among the most severe human infectious diseases. The closest known relative of *P. falciparum* is a chimpanzee parasite, *Plasmodium reichenowi*, of which one single isolate was previously known. The co-speciation hypothesis suggests that both parasites evolved separately from a common ancestor over the last 5–7 million years, in parallel with the divergence of their hosts, the hominin and chimpanzee lineages. Genetic analysis of eight new isolates of *P. reichenowi*, from wild and wild-born captive chimpanzees in Cameroon and Côte d'Ivoire, shows that *P. reichenowi* is a geographically widespread and genetically diverse chimpanzee parasite. The genetic lineage comprising the totality of global *P. falciparum* is fully included within the much broader genetic diversity of *P. reichenowi*. This finding is inconsistent with the co-speciation hypothesis. Phylogenetic analysis indicates that all extant *P. falciparum* populations originated from *P. reichenowi*, likely by a single host transfer, which may have occurred as early as 2–3 million years ago, or as recently as 10,000 years ago. The evolutionary history of this relationship may be explained by two critical genetic mutations. First, inactivation of the *CMAH* gene in the human lineage rendered human ancestors unable to generate the sialic acid Neu5Gc from its precursor Neu5Ac, and likely made humans resistant to *P. reichenowi*. More recently, mutations in the dominant invasion receptor EBA 175 in the *P. falciparum* lineage provided the parasite with preference for the overabundant Neu5Ac precursor, accounting for its extreme human pathogenicity.

chimpanzees | human evolution | *Plasmodium falciparum* | *Plasmodium reichenowi* | zoonosis

Malaria counts among the worst scourges of humankind, accounting for some 500 million clinical cases per year and more than one million deaths, mostly children (1). It amounts to an immeasurable health burden and inhibits economic prosperity in numerous tropical countries, most extensively in Africa. *Plasmodium falciparum* is the most virulent among the four *Plasmodium* species parasitic to humans, accounting for ≈85% of all malaria cases, and nearly all of the mortality. The extreme pathogenicity of *P. falciparum* has suggested that it is a recent human parasite, acquired by transfer from a nonhuman host (2). Some early molecular phylogenies seemed to be consistent with this hypothesis, because they showed *P. falciparum* to be more closely related to *Plasmodium gallinaceum*, a chicken parasite, than to any of the other human parasite species (3). A considered possibility was that *P. falciparum* evolved from an avian parasite following a horizontal host transfer, perhaps in association with the Neolithic domestication of the chicken. It was, however, shown by Escalante and Ayala (4) and Escalante et al. (5) that the closest relative of *P. falciparum* is *P. reichenowi*, a malaria parasite isolated from a captive chimpanzee that had not been included in earlier studies. These authors showed that *P. falciparum* and *P. reichenowi* form an independent clade distinct from other malaria parasites, including the other three human malaria

parasites, which appear to have originated in Old World monkeys (4, 5). The close phylogenetic relationship between *P. falciparum* and *P. reichenowi*, their distinctness from the other human malaria parasites, and their remoteness from bird or lizard parasites was soon confirmed by other studies (6–8).

Three mutually exclusive hypotheses could account for the relationship between *P. falciparum* and *P. reichenowi*. (i) Co-speciation: *P. falciparum* and *P. reichenowi* evolved from a common ancestor parasite, independently in their respective hosts, humans and chimpanzees, over the last 5–7 million years; (ii) Human origin: *P. reichenowi* evolved from an introduction of *P. falciparum* into chimpanzee hosts; or (iii) Chimpanzee origin: *P. falciparum* evolved from the introduction of the chimpanzee *P. reichenowi* into the human lineage (Fig. 1).

These hypotheses can readily be tested by determining the phylogenetic relationships among the two parasite species and by comparing their levels of genetic polymorphism, particularly with respect to silent nucleotides, which are expected to accumulate as a function of the rate of mutation and the time elapsed since the origin of a species. If co-speciation occurred, *P. falciparum* and *P. reichenowi* should form distinct, sister clades. If hypotheses ii or iii are correct, the level of neutral polymorphism should be greater in the ancestral parasite than in the derived species. In the absence of the molecular characterization of multiple strains of *P. reichenowi*, of which only one strain was available, Escalante and Ayala (4) and Escalante et al. (5) favored the co-speciation hypothesis, as soon did other investigators (6–8). The two alternative transfer hypotheses were largely ignored because of the availability of only one known isolate of *P. reichenowi* (from Cameroon, described in 1917 and 1920, ref. 9). However, Martin et al. (10) suggested a mechanism compatible with hypothesis iii on the basis of differential expression of host sialic acid (Sia) ligands and differences in parasite receptor Sia-binding preferences.

Rich, Ayala, and collaborators soon demonstrated that *P. falciparum* has very low levels of neutral polymorphism (11–13), a result that was subsequently confirmed by other investigators (14–18). The scarcity of neutral polymorphisms in *P. falciparum* was interpreted to be the result of a recent world expansion of the species, which was estimated to have happened a few

Author contributions: S.M.R., F.H.L., and N.D.W. designed research; S.M.R., F.H.L., G.X., M.L., C.F.D., M.N.A., E.E.T., J.L.D.D., and N.D.W. performed research; F.H.L., C.B., and N.D.W. contributed new reagents/analytic tools; S.M.R., G.X., and B.M.R. analyzed data; and S.M.R., B.L.P., B.M.R., P.F., F.J.A., and N.D.W. wrote the paper.

The authors declare no conflict of interest.

Freely available online through the PNAS open access option.

Data deposition: The genetic sequences reported in this paper have been deposited in the GenBank database. Accession numbers are listed in Tables S1–S3.

¹To whom correspondence may be addressed. E-mail: smrich@psis.umass.edu, ndwolfe@stanford.edu, or fjayala@uci.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0907740106/DCSupplemental.




Fig. 1. Alternative hypotheses for the origin of *P. falciparum* as a human parasite. Green- and blue-shaded areas correspond to parasites infecting chimpanzee and human hosts, denoted as C and H, respectively. The hypothetical ancestors are denoted as F (for *P. falciparum*), R (for *P. reichenowi*), and RF (for their common ancestor). (Left) Parasites of humans and chimpanzees diverged as their respective hosts did, starting 5–7 million years ago (5, 52). Alternatively, a human parasite lineage could have given rise to the chimpanzee lineage (Center), or vice versa (Right). The present data overwhelmingly support the latter alternative, identifying chimpanzees as the ancestral host for the agent of malignant human malaria.

thousand years ago, rather than millions of years ago (the Malaria Eve hypothesis; ref. 13).

The recent world expansion of *P. falciparum* could have come about in two ways. First, as a consequence of a severe bottleneck, so that all extant populations of the species would have derived from a few surviving strains. An alternative, equally viable explanation would be that *falciparum* malaria had been restricted to a local population (somewhere in tropical Africa), from where it would have expanded through much of the African tropics and beyond as a consequence of recent environmental and vector changes (12, 13, 19–21). These changes would include, among others, the introduction of agriculture in tropical Africa during the late Neolithic and consequent deforestation, creating pools of standing water and other conditions favorable for mosquito breeding; the evolution of anthropophilic *Anopheles* vectors (19, 20); and the gradual warming of the planet that eventually allowed the geographic expansion of the vector–parasite association (11, 13, 22). These three sets of events are timed within the past 10,000 years, which is consistent with the estimated time of the world expansion of *P. falciparum* (13, 21, 22). The current availability of additional isolates of *P. reichenowi* makes it now possible to investigate comparatively these hypotheses.

Results

We have sampled wild and wild-born captive chimpanzees from Côte d’Ivoire ($n = 10$) and Cameroon ($n = 84$). From the tissues collected from these chimpanzees, we have identified eight new isolates of *P. reichenowi* by amplifying three gene fragments that have been extensively used in *Plasmodium* studies: mitochondrial *cytB*, apicoplast *clpC*, and nuclear *18S rRNA*. Five isolates were derived from the 84 Cameroonian chimpanzees and three from the 10 chimpanzees from Côte d’Ivoire. One of the Ivorian chimpanzees, Rafiki, was infected with two strains of *Plasmodium*, characterized by distinct nucleotide sequences in each of the three gene fragments. Because our blood samples were collected on filter paper, we were unable to determine whether the parasites’ gametocytes were crescent-shaped, an attribute exclusively characteristic of *P. falciparum* and *P. reichenowi* (23). However, the diagnostic repeat region from the circumsporozoite protein (*Csp*) gene from five sequenced isolates (Bana, Gabon, Max, Nino, and Loukoum) shows that each gene encodes amino acid repeat motifs that specifically distinguish *P. falciparum* and *P. reichenowi* from all other malarial parasites (Fig. 2) (24), consistent with their designation as *P. reichenowi*.

We have examined the nucleotide polymorphisms among the three gene fragments. As reported previously, nucleotide polymorphisms among *P. falciparum* isolates are extremely scarce (18, 21, 25, 26). Among the *P. reichenowi* isolates, there is considerably greater nucleotide polymorphism than in *P. falciparum*, nearly all of which occurs among silent (synonymous)

Preich (Bana)	KEKNN . KLRQP AGND NVDP NANP NANP NVDP
Preich (Loukoum)	KEKNN . KLRQP AGND NVDP NANP NANP NVDP
Preich (Max)	KEKNN . KLRQP AGND NADP NADP NANP NANP
Preich (CDC1)	KPKHN . KLRQP GND NVDP NANP NANP NVDP NANP
Preich (Gabon)	KPDNN . KLRQP VDD NPDP NANP NANP NANP NANP
Preich (Nino)	KPDNN . KLRQP VDD NPDP NANP NANP NANP NANP
Pfalciparum1	KPKHK . KLRQP GDG NPDP NANP NANP NVDP NANP
Pfalciparum2	KPKHK . KLRQP GDG NPDP NANP NANP NVDP NANP
Pmalariae1	KAVEN . KLRQP PGDDDGAGNDEGN DAGN DAGN DAGN AAGNA
Pmalariae2	KAVEN . KLRQP PGDDDGAGN DAGN DAGN DAGN AAGNA
Pviva1	NPREN . KLRQP GDRADGQPA GDRADGQPA GDRADGQPA
Psimium	NPREN . KLRQP GDRADGQPA GDRADGQPA GDRADGQPA
Pcynomolgi1	KPREN . KLRQP AGNNAAGF AGNNAAGF AGNNAAGF
Pcynomolgi2	KPREN . KLRQP PAGDGA EGDGAPAA PAGDGA PA PAGDGA
Psimiovale	KPHEN . KLRQP VPGANQZGGAAGF GANQZGGAAGF
Pknowlesi1	KPNEN . KLRQP NECQPOAQGDGANAG QPOAQGDGANAG
Pknowlesi2	KPNEN . KLRQP EQPAAGAGCEQPAAGAGCEQPAAGAGG
Pberghai	IERNN . KLRQP P P P P N P N P P P P P N P N
Pyoelii1	KEAQN . KLNQP VVADENVD QGPGA QGPGA QGPGA
Pyoelii2	KEAQN . KLRQP AA VADPNAE VADPNAE VADPNAE
Pgallinaceum	YFRENVNINQP VGGNGGVQPA GNGNGGVQPA GNGNGGVQPA

Fig. 2. Circumsporozoite surface protein (*Csp*) repeat composition of several *Plasmodium* species showing that repeat units of *Csp* genes are diagnostic of species type (24). All characteristic *Csp* amino acid repeats of *P. falciparum*, including NVDP and NANP are also found in *P. reichenowi*. Some *P. reichenowi* alleles show additional repeat types (NADP and NP).

sites in the two amino acid coding regions (*cytB* and *clpC*) (Fig. 3). The number of synonymous differences among sequences is expected to increase when evolutionarily independent lineages are combined. However, the mean pairwise number of nucleotide differences (π) is unaffected when sequences from 133 isolates of *P. falciparum* are added to the nine isolates of *P. reichenowi* (Fig. 3). This observation, coupled with the related observation of much greater diversity among *P. reichenowi* isolates than among *P. falciparum* isolates (even though these are greatly more numerous) is inconsistent with the co-speciation hypothesis and supports the hypothesis that *P. falciparum* evolved from a chimpanzee parasite.

Fig. 4 shows the phylogenetic relationships among the eight new isolates of *P. reichenowi*, plus the one previously reported isolate (CDC1) together with 133 strains of *P. falciparum*, based on the *cytB* gene sequences (see Table S1). We have included in this phylogeny the three other human malaria parasites (*P. vivax*, *P. ovale*, and *P. malariae*), as well as two malaria parasites from African rodents (*P. yoelii* and *P. berghei*), and an avian malaria parasite (*P. gallinaceum*). The *cytB* gene (Fig. 4), which is thought to be best for recovering the deeper divergences within




Fig. 3. Pairwise nucleotide polymorphism (π) at three genetic loci—*cytB*, *clpC*, and *18S rRNA*—among *Plasmodium* species (see SI Text). Diversity in malaria parasites is estimated for (i) *P. falciparum* isolates alone, (ii) *P. reichenowi* isolates alone, (iii) *P. falciparum* + *P. reichenowi*, and (iv) *P. falciparum* + *P. reichenowi* + *P. vivax*. The addition of *P. vivax* significantly inflates estimates of nucleotide variation (π), which is consistent with the evolutionary independence of *P. vivax*. The addition to *P. reichenowi* of the nucleotide polymorphism found in a very large sample of 133 *P. falciparum* isolates does not substantially increase the polymorphism found in *P. reichenowi* alone.

How and when did the host transfer occur? A hypothesis proposed in the past was that the ancestors of *P. falciparum* would have been transferred from another host to humans as our Neolithic ancestors transitioned from hunter-gatherers to agriculturalists some 10,000 years ago (22, 31). This proposal was based on anthropological information about the history of our species, but also on the estimated age of hemoglobin mutants that render humans resistant to malaria infection. The Malaria Eve hypothesis, based on *P. falciparum*'s very low levels of neutral polymorphisms, is consistent with this hypothesis (13, 21), which is also supported by the recent evolution of anthropophilic vectors and by climatic considerations (19, 20, 22). Coluzzi and colleagues have shown that the rapid incipient speciation of the principal African vectors of *P. falciparum* was driven by postagricultural human conditions, which were "a key influence on the origin of the modern *P. falciparum* from an ancestral, less pathogenic, taxon" (20).

Other investigators subsequently have sought to determine the time of the most recent common ancestor of extant *P. falciparum* populations. A time-period consistent with a postagricultural introduction remains most probable (18, 25). Indirect corroboration of this hypothesis lies within the human genome. Specific genetic mechanisms that have given humans resistance to malaria, such as G6PD deficiency and other hemoglobinopathies, appear to have arisen within a time-frame consistent with a postagricultural origin (16, 17).

Our results confirm that the extant populations of *P. falciparum* have originated in the recent past, because of their very low genetic diversity, but our results do not tell us when the transfer to humans may have occurred. This transfer must have happened much earlier, allowing for the differentiation of *P. falciparum* from *P. reichenowi*, which may have originally consisted of nothing more than a change in binding specificity. At present, the divergence between the two species is not large, although much larger than the divergence among the extant *P. falciparum*. It seems likely that considerable time, on the order of many tens or hundreds of thousands of years, may have elapsed from the time of the host transfer to the time when genetic changes in the parasite and/or the human lineages made possible the rapid expansion of *P. falciparum*.

Martin et al. (10) have suggested that the classic failure of experimental cross-infection of *P. falciparum* into chimpanzees and of *P. reichenowi* into humans can be explained by a human-specific mutation event. Whereas the *P. falciparum* merozoite can use multiple pathways to invade erythrocytes (32), the dominant invasion receptor appears to be the erythrocyte-binding-like (EBA)-175. The underlying polypeptide sequence of its primary target molecule in erythrocytes is the Sia-capped N-terminal domain of the major erythrocyte glycoprotein glycoporphin-A (GYPA), which itself is undergoing rapid evolution (33), presumably due to selection pressure from the parasite.

Several million years after divergence from the chimpanzee lineage, a major biochemical change in Sia biology occurred in the human ancestral lineage (34, 35). The most common Old World primate Sias are *N*-glycolylneuraminic acid (Neu5Gc) and its metabolic precursor, *N*-acetylneuraminic acid (Neu5Ac), which differ by a single oxygen atom. A mutation in the *CMAH* gene makes humans unable to produce Neu5Gc from Neu5Ac, which accumulates in great excess on human erythrocytes. Chimpanzees carry both Sias, with Neu5Gc being dominant (10, 36). *P. reichenowi* EBA-175 has a marked preference for Neu5Gc, which was suggested to represent the ancestral condition, whereas *P. falciparum* EBA-175 has a preference for Neu5Ac, which accumulates in humans as a consequence of the *CMAH* mutation (10, 36). Martin et al. (10) thus conjectured that the loss of Neu5Gc in the human lineage (by way of the *CMAH* mutation) would have provided our emerging *Homo* ancestors, perhaps as early as 2–3 million years ago, with temporary relief

from *P. reichenowi* malaria (8, 10). Indeed, it is possible that the *CMAH* mutation was driven to fixation by continued selection pressure from the then extant form of *P. reichenowi*. The parasitic malignancy of *P. falciparum* would have come about later, by selective evolution of its EBA-175, which preferentially recognizes the Neu5Ac-rich human erythrocytes (10). It seems also likely that there would have been an intermediate stage, wherein EBA-175 of the *P. reichenowi* ancestor would have relaxed its specificity to accommodate binding of Neu5Ac. The final EBA-175 mutations potentially responsible for the malignancy and rapid expansion of *P. falciparum* may have occurred relatively recently, perhaps ≈5,000–10,000 years ago, which would account for the scarcity of neutral DNA polymorphisms in *P. falciparum*, consistent with a recent worldwide population expansion of this parasite (11, 12, 18, 22, 25).

Our investigations suggest that *P. falciparum* has only once established itself among human hosts. The zoonotic origin of *P. falciparum* elevates interest in the possible ongoing transmission of other malaria parasites of primate origin into the human population (37). The repeated emergence of human malaria parasites from zoonotic reservoirs raises the question of whether ongoing transmission of *P. reichenowi* from chimpanzees to humans may be possible (or vice versa). The fact that this transmission has not happened repeatedly may reflect the difficulty in changing the sialic acid binding specificity of the parasite-binding proteins. In this regard, it is interesting that a major barrier limiting cross-transmission of avian influenza into humans (and vice versa) is due to differences in sialic acid linkage binding specificity. In this case, it is the relative preference of the human and avian virus hemagglutinins for binding α -2-6- and α -2-3-linked sialic acids, respectively, on epithelial cells in target tissues (38–41). This is also another instance wherein a human-specific change in sialic acid biology is relevant to infectious disease transmission, as chimpanzees and other great apes do not express human upper airway epithelial α -2-6-linked sialic acid targets for human influenza viruses (42, 43).

Materials and Methods

Sample Collection and Preparation. In Côte d'Ivoire, tissue and blood samples were collected from ten chimpanzees that had died due to anthrax (44), respiratory disease (45), or other reasons in the research area of Tai National Park between 1998 and 2002 (46). DNA extractions from tissue were performed by using DNAeasy tissue kits (Qiagen). DNA was stored in several aliquots.

In Cameroon, samples were collected from captive animals in three wildlife sanctuaries. Animals were primarily wild-born and brought to the sanctuaries after confiscation by authorities or abandonment by owners. Blood samples were collected during routine health examinations, quarantine, or recaptures by sanctuary staff after escapes. Whole blood was collected via venipuncture into a syringe or EDTA vacutainer and 1 mL was spotted onto Whatman #3 filter paper or Whatman/S&S 903 filter paper. Spots were allowed to dry and were placed in an envelope and sealed in a cliplock bag with silica gel. Dried samples were frozen at -80°C once in the laboratory in the US. Dry Blood Spot filter papers were processed with Epicenter Master Complete DNA and RNA purification kits (Epicenter Technologies) to obtain genomic DNA following the manufacturer protocols. Total DNA was first dissolved in 30 μL molecular grade H_2O .

Gene Fragment Analysis. As noted by Martinsen et al. (47), reliable determination of species phylogeny is best accomplished by analyses of multiple gene fragments. Accordingly, we amplified gene fragments from the three principal genomes of *Plasmodium* parasites, mitochondrial (*cytB*), apicoplast (*clpC*), and nuclear (*18S rDNA*). The 50- μL PCRs used 1 μL stock DNA as template, 1 \times PCR buffer containing 2.5 mM MgCl_2 , 0.2 mM dNTPs, 0.25 μM of each primer, and 1.25 unit TaqDNA polymerase. PCR for *cytB* gene used outer primers DW2 (TAATGCC TAGACGTATTCCTGATTATCCAG) and DW3 (TGCTGATCATACCCTAAAG) (6), and inter primers DW1 (TCAACAATGACTTTATTTGG) and *cytB*1 (GGATCACTACAGTATATCTCC). Amplification for the *clpC* gene used outer primers TFM1421+ (AAAACCTGAATTAGCAAAAATATTA) and TFM1423RC (CGAGCTCCATATAAAGGAT) (48), and inter primers CLPCF1 (TCTAAACAAT-TATTTGGTTCTG) and CLPCRC1 (TTGGACAACCTAAATTACTTG). PCR for *18S*

gene used outer primers 18SF1 (TGTAATTGGAATGGTGGGA) and 18SRC1 (TGATCGTCTTCACTCCCTTAA), and inter primers 18SF2 (TTCCAGCTCCAAT-AGCGTATA) and 18SRC2 (GACATCTGAATACGAATGTCCC).

A touchdown PCR program was used with the annealing temperature starting at 54 °C and decreasing by 3 °C in the first 3 cycles, then 3 °C in the second 3 cycles, followed by 30 cycles at 48 °C. In each cycle, denaturing (at 94 °C) and annealing (at 54–48 °C) were 20 s, and extensions (at 68 °C) were 60 s in duration. An initial 2-min denaturation and a final 6-min extension step were also included. PCR products were separated by electrophoresis in a 1% agarose gel, excised from the gel, and purified by using a QIAquick Gel Extraction Kit (Qiagen). Purified PCR products were cloned by using a TOPO TA cloning for sequencing kit as suggested by the manufacturer, Invitrogen. For each sample, 8–48 clones were picked up and Sanger-sequenced in complementary directions by using T7 and T3 primers. The resulting nucleotide sequences were aligned by using CLUSTALW (49), with corrections made by eye to account for inappropriate alignments (e.g., frame shifts from improperly aligned codon sequences).

Phylogenetic Comparisons. We conducted a phylogenetic analysis of each gene fragment independently. These nucleotide sequences were compared with those of other human parasites (*P. falciparum*, *P. vivax*, *P. malariae*, and *P. ovale*), of African rodents (*P. berghei* and *P. yoelii*), and one isolate of bird malaria associated with chickens (*P. gallinaceum*). The sampling of these genetic loci varies greatly among the taxa, with the data for *cytB* being the most extensive in the public database. For present purposes, we have included only unique haplotypes in our phylogenetic analysis, but the occurrence of multiple representatives of that haplotype are indicated in Table S1 as well as in the resulting phylogenies.

Phylogenetic reconstructions were performed by using maximum parsimony (MP) and maximum likelihood (ML) criteria. MP and ML trees were estimated by using PAUP*4.0b4 (50). Gene-specific evolutionary models were determined empirically for ML analyses by using ModelTest (51). Not surprisingly, the genes from the different genomes adhere to markedly different models of molecular evolution. We used the hierarchical likelihood ratio tests to choose the most suitable model parameters for each genetic locus. The best fit models for *cytB*, *clpC*, and *18S rRNA* are the F81+G, GTR+G, and HKY+I+G models, respectively.

Each gene tree is shown as a phylogram with branch lengths correspondent to the degree of divergence, with the scale shown at the bottom (Fig. 4, Figs. S1 and S2). The topology of these trees is derived by the Neighbor-Joining method using gene-specific maximum likelihood parameters. Bootstrap analyses using 1,000 replicates of the MP and ML methods were also conducted. The bootstrap values for nodes supported by $\geq 50\%$ are shown in the tree for both MP and ML iterative analyses. (Bootstrap support values are not shown if $< 50\%$.) The resulting consensus dendrograms from MP and ML bootstrap analyses vary slightly in the degree to which they reliably resolve polytomies, particularly among the nonhuman primate parasites. However, in every instance there was extremely high support (93–100%) for the monophyly of the *P. reichenowi*–*P. falciparum* clade.

Testing Reciprocal Monophyly. Two means were used to evaluate whether different ancestors could likely have given rise to *P. falciparum* and *P. reiche-*

nowi, based on the distribution of variation now evident in the *cytB*, *clpC*, and *18S* genes. First, trees were searched under the criteria of maximum parsimony and maximum likelihood, with and without the imposition of a topological constraint consistent with the co-speciation hypothesis. The lengths and likelihoods of optimal trees identified in each case were compared. The cost of enforcing reciprocal monophyly, wherein the ancestor of *P. reichenowi* did not also give rise to *P. falciparum*, was judged to be considerable.

More formal statistical consideration of this cost was evaluated by means of Bayesian statistical sampling of the posterior distribution of trees identified under the null and alternative (co-speciation) hypotheses. For *cytB*, the HKY + G model was taken as a prior, which resembles the model identified as optimal by ModelTest (see above) but which also allows different rates of substitution for transitions and transversions. For the *clpC* and *18S* genes, models identified by ModelTest (see above) were taken as priors. Tree topology and model parameters were allowed to vary in 10 million generations of an MCMC algorithm seeking optimal trees constrained to be consistent with the co-speciation hypothesis, or freed from any such constraint. After a “burn-in” period of 10 thousand generations assuming exponential demographic growth and a strict molecular clock, stable estimates of substitution parameters, tree topology, and tree likelihood were obtained. The ratio of the likelihood of each model, given the data, provided a means to evaluate the statistical cost of imposing reciprocal monophyly on each dataset after integrating over other plausible values for model parameters. The “weight of evidence”, representing the ratio of likelihoods, was represented by Bayes factors (28) calculated by using the methods implemented by BEAST v. 1.4.8 (27, 29).

ACKNOWLEDGMENTS. We thank the Cameroon Ministry of Forestry and Wildlife, Ministry of Scientific Research and Innovation, Cameroon Wildlife Aid Fund, and Limbe Wildlife Centre for permits and permission to sample animals at Limbe Wildlife Centre, Mvog Betsi Zoo, and Mfou National Park. We thank Dr. John Kiyang, Dr. Felix Lankster, Babila Talfon, Jean-Michel Takuo, Felix Nkom, Mary-Chantal Bindele, and Mamo Benadette Adzenyuy for Cameroonian field support. We thank the Ministry of the Environment and Forests as well as the Ministry of Research, the directorship of the Tai National Park, and the Swiss Research Center in Abidjan in Côte d'Ivoire. For Ivorian field support, we thank Ilka Herbinger, Rebecca Stumpf, Yasmin Moebius, Cristina Gomes, Tobias Deschner, and Emmanuel Normand. Additional support was provided by the Robert Koch-Institut, the Max-Planck-Society, the Global Viral Forecasting Initiative, Google.org, and The Skoll Foundation. We are extremely grateful to Dr. François Renaud (Centre National de la Recherche Scientifique-International Relief and Development, Montpellier, France) for making available to us the DNA sequences of *Plasmodium gaboni*. Comments or assistance were provided by Norman Johnson, Dr. Mike Dean, Russell Hanson, Charles Wolfe, Dr. David Sintasath, Lisa Krain, Bill Switzer, and Dr. Jared Diamond. We are grateful to Drs. François Renaud, Nora Besansky, and Ajit Varki for helpful comments about the manuscript. Research and experimentation was funded by National Institutes of Health Grants NIH-RO1GM70077 and NIH-RO1GM60759 (to S.M.R.), with additional support from Cummings School of Veterinary Medicine at Tufts University. Support for collection of specimens in Cameroon was provided by the National Institutes of Health Director's Pioneer Award DP1-OD00370 (to N.D.W.), the National Institutes of Health Fogarty International Center International Research Scientist Development Award 5 K01 TW000003–05 (to N.D.W.), and the National Geographic Society Committee for Research and Exploration.

- Snow RW, Guerra CA, Noor AM, Myint HY, Hay SI (2005) The global distribution of clinical episodes of *Plasmodium falciparum* malaria. *Nature* 434:214–217.
- Boyd MF (1949) *Malaria: A comprehensive survey of all aspects of this group of diseases from a global standpoint* (Saunders, Philadelphia), pp 1–47.
- Waters AP, Higgins DG, McCutchan TF (1991) *Plasmodium falciparum* appears to have arisen as a result of lateral transfer between avian and human hosts. *Proc Natl Acad Sci USA* 88:3140–3144.
- Escalante AA, Ayala FJ (1994) Phylogeny of the malarial genus *Plasmodium*, derived from rRNA gene sequences. *Proc Natl Acad Sci USA* 91:11373–11377.
- Escalante AA, Barrio E, Ayala FJ (1995) Evolutionary origin of human and primate malaria: Evidence from the circumsporozoite protein gene. *Mol Biol Evol* 12:616–626.
- Escalante AA, Freeland DE, Collins WE, Lal AA (1998) The evolution of primate malaria parasites based on the gene encoding cytochrome b from the linear mitochondrial genome. *Proc Natl Acad Sci USA* 95:8124–8129.
- Hagner SC, Misof B, Maier WA, Kampen H (2007) Bayesian analysis of new and old malaria parasite DNA sequence data demonstrates the need for more phylogenetic signal to clarify the descent of *Plasmodium falciparum*. *Parasitol Res* 101:493–503.
- Perkins SL, Schall JJ (2002) A molecular phylogeny of malarial parasites recovered from cytochrome b gene sequences. *J Parasitol* 88:972–978.
- Reichenow E (1920) The presence of human malaria parasites in the African apes (Translated from German). *Cent F Bakt I Abt Orig* 85:207.
- Martin MJ, Rayner JC, Gagneux P, Barnwell JW, Varki A (2005) Evolution of human-chimpanzee differences in malaria susceptibility: Relationship to human genetic loss of N-glycolylneuraminic acid. *Proc Natl Acad Sci USA* 102:12819–12824.
- Ayala FJ, Rich SM (2000) Genetic variation and the recent worldwide expansion of *Plasmodium falciparum*. *Gene* 261:161–170.
- Rich SM, Ayala FJ (2000) Population structure and recent evolution of *Plasmodium falciparum*. *Proc Natl Acad Sci USA* 97:6994–7001.
- Rich SM, Licht MC, Hudson RR, Ayala FJ (1998) Malaria's Eve: Evidence of a recent population bottleneck throughout the world populations of *Plasmodium falciparum*. *Proc Natl Acad Sci USA* 95:4425–4430.
- Escalante AA, Cornejo OE, Rojas A, Udhayakumar V, Lal AA (2004) Assessing the effect of natural selection in malaria parasites. *Trends Parasitol* 20:388–395.
- Hartl DL, et al. (2002) The paradoxical population genetics of *Plasmodium falciparum*. *Trends Parasitol* 18:266–272.
- Tishkoff SA, et al. (2001) Haplotype diversity and linkage disequilibrium at human G6PD: Recent origin of alleles that confer malarial resistance. *Science* 293:455–462.
- Tishkoff SA, Verrelli BC (2004) *Infectious Disease and Host-Pathogen Evolution*, ed Dronamraju KR (Cambridge Univ Press, Cambridge, UK), pp 113–140.
- Volkman SK, et al. (2001) Recent origin of *Plasmodium falciparum* from a single progenitor. *Science* 293:482–484.
- Ayala FJ, Coluzzi M (2005) Chromosome speciation: Humans, Drosophila, and mosquitoes. *Proc Natl Acad Sci USA* 102(Suppl 1):6535–6542.
- Coluzzi M (1999) The clay feet of the malaria giant and its African roots: Hypotheses and inferences about origin, spread and control of *Plasmodium falciparum*. *Parassitologia* 41:277–283.
- Rich SM, Ferreira MU, Ayala FJ (2000) The origin of antigenic diversity in *Plasmodium falciparum*. *Parasitol Today* 16:390–396.

22. Ayala FJ, Escalante AA, Rich SM (1999) Evolution of *Plasmodium* and the recent origin of the world populations of *Plasmodium falciparum*. *Parassitologia* 41:55–68.
23. Gilles HM, Warrel DA (1993) *Bruce-Chwatt's Essential Malariology* (Edward Arnold, London), 3rd Ed.
24. McCutchan TF, et al. (1996) Comparison of circumsporozoite proteins from avian and mammalian malarial: Biological and phylogenetic implications. *Proc Natl Acad Sci USA* 93:11889–11894.
25. Joy DA, et al. (2003) Early origin and recent expansion of *Plasmodium falciparum*. *Science* 300:318–321.
26. Su XZ, Mu J, Joy DA (2003) The “Malaria's Eve” hypothesis and the debate concerning the origin of the human malaria parasite *Plasmodium falciparum*. *Microbes Infect* 5:891–896.
27. Drummond AJ, Rambaut A (2007) BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol Biol* 7:214.
28. Kass RE, Raftery AE (1995) Bayes factors. *J Amer Stat Assoc* 90:773–795.
29. Suchard MA, Weiss RE, Sinsheimer JS (2001) Bayesian selection of continuous-time Markov chain evolutionary models. *Mol Biol Evol* 18:1001–1013.
30. Ollomo B, et al. (2009) A new malaria agent in African hominids. *PLoS Pathog* 5:e1000446.
31. Livingstone FB (1958) The distribution of the sickle cell gene in Liberia. *Am J Hum Genet* 10:33–41.
32. Baum J, Pinder M, Conway DJ (2003) Erythrocyte invasion phenotypes of *Plasmodium falciparum* in The Gambia. *Infect Immun* 71:1856–1863.
33. Wang HY, Tang H, Shen CK, Wu CI (2003) Rapidly evolving genes in human. I. The glycoporphins and their possible role in evading malaria parasites. *Mol Biol Evol* 20:1795–1804.
34. Chou HH, et al. (2002) Inactivation of CMP-N-acetylneuraminic acid hydroxylase occurred prior to brain expansion during human evolution. *Proc Natl Acad Sci USA* 99:11736–11741.
35. Hayakawa T, Aki I, Varki A, Satta Y, Takahata N (2006) Fixation of the human-specific CMP-N-acetylneuraminic acid hydroxylase pseudogene and implications of haplotype diversity for human evolution. *Genetics* 172:1139–1146.
36. Varki A, Nelson DL (2007) Genomic comparisons of humans and chimpanzees. *Annu Rev Anthropol* 36:191–209.
37. Cox-Singh J, et al. (2008) *Plasmodium knowlesi* malaria in humans is widely distributed and potentially life threatening. *Clin Infect Dis* 46:165–171.
38. Shinya K, et al. (2006) Avian flu: Influenza virus receptors in the human airway. *Nature* 440:435–436.
39. Stevens J, et al. (2006) Structure and receptor specificity of the hemagglutinin from an H5N1 influenza virus. *Science* 312:404–410.
40. Tumpey TM, et al. (2007) A two-amino acid change in the hemagglutinin of the 1918 influenza virus abolishes transmission. *Science* 315:655–659.
41. Yen HL, et al. (2007) Inefficient transmission of H5N1 influenza viruses in a ferret contact model. *J Virol* 81:6890–6898.
42. Couceiro JN, Paulson JC, Baum LG (1993) Influenza virus strains selectively recognize sialyloligosaccharides on human respiratory epithelium: The role of the host cell in selection of hemagglutinin receptor specificity. *Virus Res* 29:155–165.
43. Gagneux P, et al. (2003) Human-specific regulation of alpha 2–6-linked sialic acids. *J Biol Chem* 278:48245–48250.
44. Leendertz FH, et al. (2004) Anthrax kills wild chimpanzees in a tropical rainforest. *Nature* 430:451–452.
45. Kondgen S, et al. (2008) Pandemic human viruses cause decline of endangered great apes. *Curr Biol* 18:260–264.
46. Leendertz FH, et al. (2006) Pathogens as drivers of population declines: The importance of systematic monitoring in great apes and other threatened mammals. *Biol Cons* 131:325–337.
47. Martinsen ES, Perkins SL, Schall JJ (2008) A three-genome phylogeny of malaria parasites (*Plasmodium* and closely related genera): Evolution of life-history traits and host switches. *Mol Phylogenet Evol* 47:261–273.
48. Rathore D, Wahl AM, Sullivan M, McCutchan TF (2001) A phylogenetic comparison of gene trees constructed from plastid, mitochondrial and genomic DNA of *Plasmodium* species. *Mol Biochem Parasitol* 114:89–94.
49. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL_X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25:4876–4882, 2nd Ed.
50. Swofford D (2003) *Molecular Systematics* (Sinauer Associates, Sunderland, MA).
51. Posada D, Crandall KA (1998) MODELTEST: Testing the model of DNA substitution. *Bioinformatics* 14:817–818.
52. Escalante AA, Ayala FJ (1995) Evolutionary origin of *Plasmodium* and other Apicomplexa based on rRNA genes. *Proc Natl Acad Sci USA* 92:5793–5797.