

Sulforaphane destabilizes the androgen receptor in prostate cancer cells by inactivating histone deacetylase 6

Angela Gibbs¹, Jacob Schwartzman¹, Vivianne Deng, and Joshi Alumkal²

Division of Hematology and Oncology, Oregon Health and Science University, 3181 SW Sam Jackson Park Road, Portland, OR 97239

Communicated by Brian J. Druker, Oregon Health and Science University, Portland, OR, August 11, 2009 (received for review November 10, 2008)

High consumption of cruciferous vegetables is associated with a reduced risk of prostate cancer in epidemiological studies. There is preliminary evidence that sulforaphane, derived from glucoraphanin found in a number of crucifers, may prevent and induce regression of prostate cancer and other malignancies in preclinical models, but the mechanisms that may explain these effects are not fully defined. Recent reports show that sulforaphane may impair prostate cancer growth through inhibition of histone deacetylases, which are up-regulated in cancer. Indeed, one of these enzymes, histone deacetylase 6 (HDAC6), influences the acetylation state of a key androgen receptor (AR) chaperone, HSP90. AR is the central signaling pathway in prostate cancer, and its inhibition is used for both prevention and treatment of this disease. However, it is not known whether the effects of sulforaphane involve suppression of AR. We hypothesized that sulforaphane treatment would lead to hyperacetylation of HSP90 and that this would destabilize AR and attenuate AR signaling. We confirmed this by demonstrating that sulforaphane enhances HSP90 acetylation, thereby inhibiting its association with AR. Moreover, AR is subsequently degraded in the proteasome, which leads to reduced AR target gene expression and reduced AR occupancy at its target genes. Finally, sulforaphane inhibits HDAC6 deacetylase activity, and the effects of sulforaphane on AR protein are abrogated by overexpression of HDAC6 and mimicked by HDAC6 siRNA. The inactivation by sulforaphane of HDAC6-mediated HSP90 deacetylation and consequent attenuation of AR signaling represents a newly defined mechanism that may help explain this agent's effects in prostate cancer.

HSP90 | acetylation | ERG

High consumption of cruciferous vegetables is associated with a lower risk of prostate cancer in epidemiological studies, although the precise constituents that may mediate this observation are unknown (1–4). Sulforaphane, a derivative of glucoraphanin found in crucifers, has heterogeneous biological activities including Phase 2 enzyme induction, cell cycle arrest, and apoptosis, but data on pathways that mediate effects on cell growth, survival, and differentiation are incomplete (5). Of note, several reports show that sulforaphane treatment of prostate cancer cells *in vitro* leads to reduced prostate cancer cell survival, and treatment of xenograft implants or transgenic animal models of prostate cancer with sulforaphane inhibits tumor formation and metastases (6–8).

Recently, sulforaphane was shown to inhibit histone deacetylase (HDAC) proteins, which are up-regulated in cancer (6, 9–11). Despite their name, some HDAC proteins remove acetyl groups from histone proteins whereas others act on non-histone proteins. Histone deacetylation reduces gene expression whereas non-histone protein deacetylation changes protein function (11). Histone deacetylase 6 (HDAC6) is a cytoplasmic non-histone protein deacetylase whose substrates include alpha-tubulin and the HSP90 chaperone protein (12–14). When HDAC6 deacetylates HSP90, this leads to activation of HSP90, enhanced binding of HSP90 to client proteins including the androgen receptor

(AR) protein, and consequently attenuated degradation of client proteins including AR (13). Conversely, HSP90 hyperacetylation after treatment with certain HDAC inhibitors results in dissociation of HSP90 from client proteins such as AR and AR degradation (12, 13, 15, 16).

Indeed, AR, which is activated by androgens, is the most therapeutically relevant target in malignant prostate epithelial cells (17). Both preclinical studies and clinical trials have demonstrated the usefulness and effectiveness of targeting AR with hormonal agents, which act by reducing androgen production or binding to AR for prostate cancer prevention and treatment (17–19). Hormonal agents targeting AR in recurrent prostate cancer are commonly used but are not curative, and resistance mechanisms including AR gene amplification, AR mutations, and up-regulation of pathways that activate AR independent of androgens eventually develop in patients (20). In addition, recent work also demonstrates that androgen levels sufficient for AR activation persist in prostate cancer cells in man despite the use of hormonal agents, and that AR transcript variants, which encode for AR proteins, which are active even in the absence of androgens, exist (21, 22).

Thus, the destabilization and degradation of AR protein represents a rational strategy to interfere with AR signaling and to overcome the aforementioned resistance mechanisms. The present study was designed to test the hypothesis that sulforaphane would inhibit the function of HDAC6, and that this would suppress the stability or function of AR. Sulforaphane's ability to suppress HDAC6 was unknown, and its effects on AR signaling remained poorly characterized. We describe herein experimental results that confirm that sulforaphane destabilizes AR protein and disrupts AR signaling by inactivating HDAC6.

Results

Sulforaphane Treatment Increases HSP90 Acetylation and Leads to Dissociation of AR from HSP90. We treated LNCaP prostate cancer cells with sulforaphane and showed increased enrichment of acetylated lysines on HSP90 compared with vehicle 4 h after treatment (Fig. 1A). In the input samples, HSP90, HDAC6, and alpha-tubulin levels were similar between vehicle and sulforaphane-treated cells; levels of acetylated alpha-tubulin, a HDAC6 target, increased in sulforaphane-treated samples, indicating inhibition of protein deacetylation (Fig. 1A). Similar results were seen in VCaP prostate cancer cells (Fig. S1). Sulforaphane treatment also disrupted the interaction between HSP90 and AR

Author contributions: A.G., J.S., and J.A. designed research; A.G., J.S., V.D., and J.A. performed research; A.G., J.S., V.D., and J.A. analyzed data; and A.G., J.S., and J.A. wrote the paper.

The authors declare no conflict of interest.

¹A.G. and J.S. contributed equally to this work.

²To whom correspondence should be addressed at: Oregon Health and Science University, 3303 SW Bond Avenue, CH14R, Portland, OR 97239. E-mail: alumkalj@ohsu.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0908908106/DCSupplemental.




Fig. 1. Sulforaphane treatment of prostate cancer cells increases HSP90 acetylation and dissociates it from AR. (A and B) Immunoprecipitations were carried out after treatment with sulforaphane (SFN) 20 μ M or vehicle for 4 h followed by a Western blot for (A) HSP90 and (B) AR. Enrichment was quantified for each immunoprecipitation. Inputs were probed with the indicated antibodies by Western blot and were quantified.

(Fig. 1B). The AR immunoprecipitation followed by Western blotting for AR demonstrates similar AR protein levels between vehicle (lane 2) and sulforaphane-treated (lane 6) cells in the setting of decreased HSP90–AR interaction with sulforaphane treatment (lanes 1 and 5). Levels of acetylation of the androgen receptor were similar in both conditions (lanes 3 and 7). In the input samples, levels of HSP90, AR, and alpha-tubulin were similar between vehicle and sulforaphane-treated cells whereas sulforaphane increased levels of acetylated alpha-tubulin (Fig. 1B).

Sulforaphane Treatment Lowers AR Protein Levels. Using later time points, we treated LNCaP cells with sulforaphane and found that AR protein levels decreased at 12 h for the highest doses of sulforaphane (10–20 μ M) (Fig. 2A). In VCaP cells, at 24 h, AR protein levels declined for all sulforaphane dose levels (Fig. 2B). Similar results were seen for both cell lines with the HDAC inhibitor trichostatin A (TSA) and the triterpenoid CDDO-Imidazole that, like sulforaphane, acts via the Nrf2-Keap1 pathway (Fig. S2). AR transcripts were not consistently sup-

pressed by sulforaphane treatment, emphasizing that posttranscriptional mechanisms are involved (Fig. S3). Thus, across both AR-expressing cell lines, sulforaphane treatment reduced AR protein levels.

Sulforaphane Treatment Lowers AR Target Gene Expression. We next determined the functional consequences of AR protein depletion by performing real-time PCR for several AR target genes, *PSA* and the *TMPRSS2-ERG* gene fusion, which is present in VCaP cells (Fig. 3A–C). There was a dose-dependent reduction in gene expression in both cell lines. Similar results were seen with TSA. Thus, there is a high concordance between reduced AR protein levels and reduced AR target gene expression in prostate cancer cells.

Sulforaphane Treatment Reduces AR Occupancy at Its Androgen Response Elements (AREs) and Lowers ERG Protein Levels. To determine whether AR protein depletion from its target genes' AREs was responsible for reduced target gene expression, we performed chromatin immunoprecipitation (ChIP). Sulforaphane treatment reduced enrichment of AR at the *PSA* ARE in both cell lines (Fig. S4A and B). Similarly, AR enrichment was decreased with sulforaphane treatment at the *TMPRSS2* ARE (Fig. S4C). A Western blot with VCaP protein lysates at this same time point confirms near-absent levels of AR and ERG proteins after sulforaphane treatment (Fig. S4D).

Proteasome Inhibitor Treatment Rescues AR Protein from Sulforaphane-Induced Degradation. Because it is known that hyperacetylation of HSP90 inhibits its function and targets its client proteins to the proteasome for posttranslational degradation, we assessed the effects of treatment with sulforaphane and TSA with the proteasome inhibitor MG132 (Fig. 4). Treatment with sulforaphane (lane 2) or TSA (lane 4) in the absence of MG132 reduced AR protein levels versus the vehicle-treated cells (lane 6). However, simultaneous treatment with MG132 and either sulforaphane (lane 1) or TSA (lane 3) restored the AR steady-state level close to that seen in the vehicle-treated control (lane 5).

Sulforaphane Inhibits HDAC6 Enzymatic Function. We determined recombinant HDAC6 enzymatic activity on its tubulin substrate in a cell-free tubulin deacetylase assay (Fig. 5). Incubation of vehicle-treated recombinant HDAC6 with tubulin dimers (lane




Fig. 2. Sulforaphane treatment of prostate cancer cells lowers AR protein levels. (A) Western blot of protein lysates from LNCaP cells treated with vehicle, increasing doses of sulforaphane, or TSA (trichostatin A) at the indicated time points. (B) Western blot of protein lysates from VCaP cells at 24 h. AR and GAPDH levels by Western blot were quantified.




Fig. 3. Sulforaphane treatment of prostate cancer cells reduces AR target gene expression. (A) Real-time PCR of *PSA* expression from LNCaP cells treated with vehicle, increasing doses of sulforaphane, or TSA at the indicated time points. (B and C) Real-time PCR of *PSA* (B) and *TMPRSS2-ERG* (C) gene expression from VCaP cells at 24 h. The vehicle-treated sample was set to 1. *18S* was used as an endogenous control in all assays.

2) reduced levels of acetylated tubulin versus the vehicle-treated tubulin dimers incubated without HDAC6 (lane 1), which indicated that the HDAC6 enzyme was functional. However, incubation of HDAC6 with sulforaphane (lanes 3 and 4) or TSA (lane 5) led to decreased HDAC6 deacetylase activity as evidenced by increased levels of alpha-tubulin compared with the vehicle-treated HDAC6 control (lane 2).

HDAC6 Overexpression Rescues AR and HDAC6 Proteins from Sulforaphane Treatment. We overexpressed empty vector or FLAG-HDAC6 in LNCaP cells followed by treatment with sulforaphane or vehicle (Fig. 6A). Overexpression of HDAC6 led to a detectable anti-FLAG signal and higher levels of HDAC6 versus empty-vector-transfected cells, whereas levels of acetylated alpha-tubulin declined, indicating the presence of functional HDAC6 protein. Surprisingly, in empty-vector-transfected cells, HDAC6 protein levels were reduced with sulforaphane treatment; although sulforaphane does reduce *HDAC6* transcript levels, this does not fully account for the observed HDAC6 protein depletion (Fig. S5A). Concomitant proteasome inhibitor and sulforaphane treatment (lane 4) restored HDAC6 protein levels versus sulforaphane-treated cells (lane 3) whereas TSA had minimal effects on HDAC6 protein levels (lanes 5 and 6) (Fig. S6). Additionally, in empty-vector-transfected cells, sulforaphane treatment lowered AR protein and increased acetylation of alpha-tubulin. However, in HDAC6-overexpressing,

sulforaphane-treated cells, HDAC6 protein expression was restored, AR protein depletion was attenuated (without changing *AR* transcript levels), and alpha-tubulin acetylation was blunted versus the effects seen in empty-vector, sulforaphane-treated cells (Fig. 6A, Fig. S5B).

HDAC6 siRNA Recapitulates Sulforaphane's Effect on AR Protein Levels. We transiently transfected LNCaP cells with either a control or *HDAC6* siRNA. *HDAC6* siRNA depleted cells of HDAC6, which was most pronounced at 72 h (Fig. 6B). At this later time point, depletion of AR protein and acetylation of alpha-tubulin in the HDAC6 knockdown cells were most pronounced versus the control siRNA cells (Fig. 6B).

Discussion

High consumption of cruciferous vegetables is associated with a lower risk of prostate cancer development, but the precise constituents that may mediate this association are unknown. One possible candidate is sulforaphane, derived from glucoraphanin found in crucifers, which has been shown to have effects as a chemopreventive and anticancer agent in multiple preclinical systems through Phase 2 enzyme induction, cell cycle arrest, and apoptosis (1–4, 8, 23–33). However, the molecular mechanisms that




Fig. 4. Proteasome inhibitor treatment of prostate cancer cells rescues AR protein from sulforaphane treatment. LNCaP cancer cells were treated for 24 h with 20 μM sulforaphane with or without 10 μM MG132, 300 nM TSA with or without 10 μM MG132, 10 μM MG132, or vehicle. AR levels and GAPDH levels by Western blot were quantified.




Fig. 5. Sulforaphane treatment inhibits HDAC6. Tubulin dimers were either incubated without recombinant HDAC6 or with recombinant HDAC6 in the presence of vehicle, sulforaphane, or TSA. Levels of HDAC6, acetylated tubulin, and tubulin by Western blot were quantified.




Fig. 6. Ectopic overexpression of HDAC6 attenuates sulforaphane-mediated AR and HDAC6 protein depletion, and HDAC6 siRNA recapitulates the findings seen with sulforaphane. (A) LNCaP cells were transfected with pCDNA3.1 or FLAG-HDAC6. Cells were then treated with either vehicle or 15 μ M sulforaphane. Intensity values for bands by Western blot in the respective vehicle controls were set to 1. (B) LNCaP cells were transfected with siRNA to either the *luciferase* gene (si LUC) or *HDAC6* gene (si HDAC6). Levels of protein expression by Western blot were quantified, and the bands from the *HDAC6* siRNA samples were compared to the *luciferase* control samples from the same time point.

underlie the previously observed antitumor effects of sulforaphane in prostate cancer models have not been fully clarified.

The AR protein is the central, therapeutically relevant pathway in prostate cancer, and hormonal deprivation strategies are commonly used in the treatment of all phases of this disease and work by interfering with androgen production or binding to AR, which is an activating event (17). Thus, interference with the state of AR activation or AR protein levels represents a rational strategy to ablate AR signaling for the prevention and treatment of this disease (17–19).

We show here that sulforaphane treatment increases acetylation of two HDAC6 target proteins: HSP90 and alpha-tubulin. At early time points, in the setting of unchanged HDAC6 protein levels, HSP90 becomes hyperacetylated, and the AR protein dissociates from HSP90, which matches an earlier report with the HDAC inhibitor LAQ824 (Fig. 1, Fig S1) (15). At later time points, after treatment with sulforaphane or the HDAC inhibitor TSA, AR protein levels decline, and this decline is attenuated by inhibition of proteasomal degradation, which suggests that sulforaphane treatment targets AR for proteolytic degradation (Figs. 2 and 4). This same effect has been demonstrated for other HSP90 client proteins including ErbB2, bcr-abl and c-kit after interference with HSP90 function with pharmacological HDAC inhibitors (12, 15, 34).

Although AR protein levels declined without reductions in *AR* transcript levels for most doses of sulforaphane, for the highest doses of sulforaphane (20 μ M in LNCaP cells and 15–20 μ M in VCaP cells) *AR* transcript levels also declined. We found that the HDAC inhibitor TSA also suppressed *AR* transcripts, a response that has been reported by others and with other HDAC inhibitors (Fig. S3) (35, 36). Consequently, the overall effect of sulforaphane and other agents with HDAC inhibitory properties on AR levels may not be solely posttranscriptional. It is clear, however, that a posttranslational effect is a critical control point in light of our observations that AR protein levels are rescued by simultaneous sulforaphane and proteasome inhibitor treatment or overexpression of HDAC6, and that most sulforaphane doses do not lower *AR* transcript levels (Figs. 4 and 6 and Figs. S3 and S5).

We confirmed that the disappearance of AR corresponded to reduced AR binding to and expression of its target genes (Fig. 3, Fig. S4). The reduced enrichment of AR at its target genes by ChIP demonstrates that AR depletion is mediating the reduced AR target gene expression. Notably, we found that levels of *TMPRSS2-ERG*, a fusion of the AR-regulated *TMPRSS2* gene

and the *ERG* transcription factor, which is commonly overexpressed in human prostate cancer, are reduced (37). *ERG* overexpression in normal prostate cells leads to enhanced invasiveness and growth, and *ERG* knockdown in VCaP prostate cancer cells by siRNA leads to decreased invasiveness (38). *ERG* overexpression under an AR-regulated promoter in a transgenic murine model was recently shown by two independent groups to transform prostate cells and to induce formation of prostate cancer precursor lesions highlighting *ERG*'s importance in early stages of prostate tumorigenesis (38, 39). Thus, therapies such as sulforaphane, which deplete cells of AR and ERG protein, may hold promise for the prevention and treatment of prostate cancer.

We focused on HDAC6 as a key candidate for a sulforaphane target because others have shown that HDAC6, specifically, interacts with HSP90 and alpha-tubulin and deacetylates these proteins and that hyperacetylation of HSP90 cripples its chaperone function and leads to client protein dissociation and degradation (12–16). We demonstrated, by using a cell-free system, that incubation of recombinant HDAC6 with sulforaphane inhibits HDAC6 deacetylase activity. In cells, we showed that sulforaphane treatment inhibits HDAC6 function without changing its levels at early time points (4 h) (Figs. 1 and 5). However, at later time points (16 h), we have shown that sulforaphane leads to reduced HDAC6 protein levels (Fig. 6A, Fig. S6). The related compound CDDO-Imidazole also depletes HDAC6 in a dose-dependent manner, which parallels reduced AR protein levels and increased tubulin acetylation (Fig. S2). CDDO-Imidazole was previously shown to deplete Her2Neu protein levels in breast cancer cells through proteasomal degradation, although the effect of this agent on HDAC6 was not explored in that report (40). Given that CDDO-Imidazole, like sulforaphane, acts via the Nrf2 pathway, it is possible that the effect of these agents on HDAC6 protein levels is mediated by activation of gene targets of the Nrf2–Keap1 pathway.

There are several possible mechanisms for the reduced HDAC6 protein levels (beyond reduced *HDAC6* transcript levels) (Fig. 6A, Fig. S5A); however, our data with proteasome inhibitor rescue suggests that proteasomal degradation is principal (Fig. S6). The importance of reduced HDAC6 levels with sulforaphane treatment is highlighted by the fact that overexpres-




Fig. 7. Model of sulforaphane attenuation of AR signaling via HDAC6 inactivation. Normally, HSP90 is deacetylated by HDAC6, which enables it to chaperone client proteins such as AR. HDAC6 also deacetylates alpha-tubulin. With sulforaphane treatment, HDAC6 is inhibited or targeted for protein degradation. This leads to hyperacetylated alpha-tubulin; the functional significance of this remains unclear. The HDAC6 inactivation with sulforaphane also leads to hyperacetylated, inactive HSP90 protein, which dissociates from AR, and AR is then targeted for protein degradation. Consequently, AR binding to its target gene androgen response elements (ARE), including *TMPRSS2-ERG*, is diminished, which reduces their expression.

22. Hu R, et al. (2009) Ligand-independent androgen receptor variants derived from splicing of cryptic exons signify hormone-refractory prostate cancer. *Cancer Res* 69:16–22.
23. Chung FL, Conaway CC, Rao CV, Reddy BS (2000) Chemoprevention of colonic aberrant crypt foci in Fischer rats by sulforaphane and phenethyl isothiocyanate. *Carcinogenesis* 21:2287–2291.
24. Fahey JW, Zhang Y, Talalay P (1997) Broccoli sprouts: An exceptionally rich source of inducers of enzymes that protect against chemical carcinogens. *Proc Natl Acad Sci USA* 94:10367–10372.
25. Zhang Y, Kensler TW, Cho CG, Posner GH, Talalay P (1994) Anticarcinogenic activities of sulforaphane and structurally related synthetic norbornyl isothiocyanates. *Proc Natl Acad Sci USA* 91:3147–3150.
26. Bonnesen C, Eggleston IM, Hayes JD (2001) Dietary indoles and isothiocyanates that are generated from cruciferous vegetables can both stimulate apoptosis and confer protection against DNA damage in human colon cell lines. *Cancer Res* 61:6120–6130.
27. Fimognari C, et al. (2002) Cyclin D3 and p53 mediate sulforaphane-induced cell cycle delay and apoptosis in non-transformed human T lymphocytes. *Cell Mol Life Sci* 59:2004–2012.
28. Fimognari C, et al. (2002) Growth inhibition, cell-cycle arrest and apoptosis in human T-cell leukemia by the isothiocyanate sulforaphane. *Carcinogenesis* 23:581–586.
29. Gamet-Payrastré L, et al. (2000) Sulforaphane, a naturally occurring isothiocyanate, induces cell cycle arrest and apoptosis in HT29 human colon cancer cells. *Cancer Res* 60:1426–1433.
30. Gamet-Payrastré L, et al. (1998) Selective cytostatic and cytotoxic effects of glucosinolates hydrolysis products on human colon cancer cells in vitro. *Anticancer Drugs* 9:141–148.
31. Gingras D, et al. (2004) Induction of medulloblastoma cell apoptosis by sulforaphane, a dietary anticarcinogen from Brassica vegetables. *Cancer Lett* 203:35–43.
32. Singh AV, Xiao D, Lew KL, Dhir R, Singh SV (2004) Sulforaphane induces caspase-mediated apoptosis in cultured PC-3 human prostate cancer cells and retards growth of PC-3 xenografts in vivo. *Carcinogenesis* 25:83–90.
33. Wang L, et al. (2004) Targeting cell cycle machinery as a molecular mechanism of sulforaphane in prostate cancer prevention. *Int J Oncol* 24:187–192.
34. Bauer S, Yu LK, Demetri GD, Fletcher JA (2006) Heat shock protein 90 inhibition in imatinib-resistant gastrointestinal stromal tumor. *Cancer Res* 66:9153–9161.
35. Marrocco DL, et al. (2007) Suberoylanilide hydroxamic acid (vorinostat) represses androgen receptor expression and acts synergistically with an androgen receptor antagonist to inhibit prostate cancer cell proliferation. *Mol Cancer Ther* 6:51–60.
36. Rokhlin OW, et al. (2006) Mechanisms of cell death induced by histone deacetylase inhibitors in androgen receptor-positive prostate cancer cells. *Mol Cancer Res* 4:113–123.
37. Tomlins SA, et al. (2005) Recurrent fusion of TMPRSS2 and ETS transcription factor genes in prostate cancer. *Science* 310:644–648.
38. Tomlins SA, et al. (2008) Role of the TMPRSS2-ERG gene fusion in prostate cancer. *Neoplasia* 10:177–188.
39. Klezovitch O, et al. (2008) A causal role for ERG in neoplastic transformation of prostate epithelium. *Proc Natl Acad Sci USA* 105:2105–2110.
40. Konopleva M, et al. (2006) Synthetic triterpenoid 2-cyano-3,12-dioxooleana-1,9-dien-28-oic acid induces growth arrest in HER2-overexpressing breast cancer cells. *Mol Cancer Ther* 5:317–328.
41. Qian DZ, et al. (2006) Targeting tumor angiogenesis with histone deacetylase inhibitors: The hydroxamic acid derivative LBH589. *Clin Cancer Res* 12:634–642.
42. Flanagan S, Nelson JD, Castner DG, Denisenko O, Bomsztyk K (2008) Microplate-based chromatin immunoprecipitation method, Matrix CHIP: A platform to study signaling of complex genomic events. *Nucleic Acids Res* 36:e17.
43. Wang Q, et al. (2007) A hierarchical network of transcription factors governs androgen receptor-dependent prostate cancer growth. *Mol Cell* 27:380–392.
44. Yoon HG, Wong J (2006) The corepressors silencing mediator of retinoid and thyroid hormone receptor and nuclear receptor corepressor are involved in agonist- and antagonist-regulated transcription by androgen receptor. *Mol Endocrinol* 20:1048–1060.