

# Terpene synthase genes in eukaryotes beyond plants and fungi: Occurrence in social amoebae

Xinlu Chen<sup>a</sup>, Tobias G. Köllner<sup>b</sup>, Qidong Jia<sup>c</sup>, Ayla Norris<sup>c</sup>, Balaji Santhanam<sup>d,e</sup>, Patrick Rabe<sup>f</sup>, Jeroen S. Dickschat<sup>f</sup>, Gad Shaulsky<sup>e</sup>, Jonathan Gershenzon<sup>b</sup>, and Feng Chen<sup>a,c,1</sup>

<sup>a</sup>Department of Plant Sciences, University of Tennessee, Knoxville, TN 37996; <sup>b</sup>Department of Biochemistry, Max Planck Institute for Chemical Ecology, D-07745 Jena, Germany; <sup>c</sup>Graduate School of Genome Science and Technology, University of Tennessee, Knoxville, TN 37996; <sup>d</sup>Graduate Program in Structural Computational Biology and Molecular Biophysics, Baylor College of Medicine, Houston, TX 77030; <sup>e</sup>Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX 77030; and <sup>f</sup>Kekulé-Institute of Organic Chemistry and Biochemistry, University of Bonn, 53121 Bonn, Germany

Edited by Jerrold Meinwald, Cornell University, Ithaca, NY, and approved August 30, 2016 (received for review June 27, 2016)

Terpenes are structurally diverse natural products involved in many ecological interactions. The pivotal enzymes for terpene biosynthesis, terpene synthases (TPSs), had been described only in plants and fungi in the eukaryotic domain. In this report, we systematically analyzed the genome sequences of a broad range of nonplant/nonfungus eukaryotes and identified putative *TPS* genes in six species of amoebae, five of which are multicellular social amoebae from the order of Dictyosteliida. A phylogenetic analysis revealed that amoebal TPSs are evolutionarily more closely related to fungal TPSs than to bacterial TPSs. The social amoeba *Dictyostelium discoideum* was selected for functional study of the identified TPSs. *D. discoideum* grows as a unicellular organism when food is abundant and switches from vegetative growth to multicellular development upon starvation. We found that expression of most *D. discoideum* *TPS* genes was induced during development. Upon heterologous expression, all nine TPSs from *D. discoideum* showed sesquiterpene synthase activities. Some also exhibited monoterpene and/or diterpene synthase activities. Direct measurement of volatile terpenes in cultures of *D. discoideum* revealed essentially no emission at an early stage of development. In contrast, a bouquet of terpenes, dominated by sesquiterpenes including  $\beta$ -barbatene and (*E,E*)- $\alpha$ -farnesene, was detected at the middle and late stages of development, suggesting a development-specific function of volatile terpenes in *D. discoideum*. The patchy distribution of *TPS* genes in the eukaryotic domain and the evidence for *TPS* function in *D. discoideum* indicate that the *TPS* genes mediate lineage-specific adaptations.

terpene synthases | amoebae | volatiles | evolution | chemical ecology

Terpenes constitute a structurally diverse class of natural products. They are synthesized from two universal precursors: isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP), which are supplied by the mevalonate pathway and/or the methylerythritol phosphate pathway (1). From IPP and DMAPP, isoprenyl diphosphates of various chain lengths are produced by the action of isoprenyl diphosphate synthases (IDSs) (2). Among the many metabolic fates of isoprenyl diphosphates (3), they serve as substrates for terpene synthases, which convert isoprenyl diphosphates to different subclasses of terpenes of fascinating structural diversity, such as monoterpenes, sesquiterpenes, and diterpenes (4). The ability of an organism to produce terpenes depends on whether the organism contains terpene synthase genes.

Unlike *IDS* genes, which are ubiquitous in living organisms, the occurrence of terpene synthase genes and, thus, the production of terpenes appear to be lineage-specific. Presently, two general types of terpene synthases are recognized: classic terpene synthases (abbreviated as TPSs) and IDS-type terpene synthases. The majority of terpene synthases that have been characterized so far belongs to the classic TPSs. In prokaryotes, classic *TPS* genes are widely distributed in bacteria (5, 6), whereas none has been observed in archaea. In eukaryotes, classic *TPS* genes had been found only in land plants (7, 8) and fungi (9, 10), whereas

the IDS-type terpene synthases have been identified recently in two species of insects (11, 12). Sequence analysis of these insect genes suggests that they have evolved recently from insect IDSs (12), whereas classic TPSs probably also evolved from IDSs, but anciently (13). *TPS* genes are major contributors to the chemical diversity exhibited by living organisms, so it is important to understand their distribution and evolution.

In the current global tree of eukaryotes, a domain that is composed of diverse organisms, the five supergroups Opisthokonta, Amoebozoa, Excavata, Archaeplastida, and SAR (stramenopiles + alveolates + Rhizaria) are recognized (14, 15). Only the supergroup Archaeplastida, which contains land plants, and Opisthokonta, which contains fungi, are known to contain classic *TPS* genes. It has been accepted that classic *TPS* genes are absent in insects (12), which are in the supergroup of Opisthokonta. The presence/absence of *TPS* genes in other eukaryotes has not been systematically investigated. Terpenes serve diverse functions in the organisms that produce them, including defense against predators and attraction of beneficial organisms (16), which implies that *TPS* genes play a role in evolutionary adaptations. The goals of this study were to systematically search for classic *TPS* genes in nonplant/nonfungus eukaryotes, infer their evolutionary relationship to known TPSs, and understand their biochemical and biological functions.

## Significance

Many living organisms use terpenes for ecological interactions. Terpenes are biosynthesized by terpene synthases (TPSs), but classic *TPS* genes are known to exist only in plants and fungi among the eukaryotes. In this study, *TPS* genes were identified in six species of amoebae with five of them being multicellular social amoebae. Amoebal TPSs showed closer relatedness to fungal TPSs than bacterial TPSs. In the social amoeba *Dictyostelium discoideum*, all nine *TPS* genes encoded active enzymes and most of their terpene products were released as volatiles in a development-specific manner. This study highlights a wider distribution of *TPS* genes in eukaryotes than previously thought and opens a door to studying the function and evolution of *TPS* genes and their products.

Author contributions: X.C., T.G.K., J.S.D., G.S., J.G., and F.C. designed research; X.C., T.G.K., Q.J., A.N., B.S., and P.R. performed research; X.C., T.G.K., Q.J., J.S.D., G.S., J.G., and F.C. analyzed data; and X.C., T.G.K., J.S.D., G.S., J.G., and F.C. wrote the paper.


The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Data deposition: The sequences for the biochemically characterized terpene synthases reported in this paper have been deposited in the GenBank database (accession nos. [KX364374](https://doi.org/10.1073/pnas.1610379113)–[KX364382](https://doi.org/10.1073/pnas.1610379113)).

<sup>1</sup>To whom correspondence should be addressed. Email: [fengc@utk.edu](mailto:fengc@utk.edu).

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1610379113/-DCSupplemental](https://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1610379113/-DCSupplemental).



**Fig. 1.** Distribution of terpene synthase (*TPS*) genes among the major lineages of eukaryotes with sequenced genomes. A total of 168 species (Table S1), which did not include any species from land plants and fungi (Holomycota), were analyzed. The phylogeny of eukaryotes was adapted from Adl et al. (14) and Burki (15) with five supergroups recognized: Opisthokonta, Amoebozoa, Excavata, Archaeplastida, and SAR (stramenopiles + alveolates + Rhizaria). The first number (before the slash) indicates the number of species in certain lineages that were determined to contain *TPS* genes. The second number (after the slash) indicates the total number of species in that lineage that were analyzed. NA, not analyzed. The “+2” indicates that two additional species from Amoebozoa were identified to contain *TPS* genes in the nonredundant database at NCBI.

## Results

**Identification of Terpene Synthase Genes in Nonplant/Nonfungus Eukaryotes.** To determine the occurrence of *TPS* genes in eukaryotes other than plants and fungi, a HMMER search (17) was conducted by using a set of 168 well-annotated genomes (Table S1) of nonplant/nonfungus eukaryotes. Whereas the general absence of *TPS* genes in nonplant/nonfungus eukaryotes was confirmed, *TPS* genes were detected in the two supergroups of Amoebozoa and Excavata (Fig. 1). Among the seven species of Amoebozoa analyzed (Table S1), all three species from the genus *Dictyostelium*, *Dictyostelium discoideum*, *Dictyostelium fasciculatum*, and *Dictyostelium purpureum*, were found to contain *TPS* genes, whereas no *TPS* genes were found in the other four species *Entamoeba histolytica*, *Entamoeba dispar*, *Entamoeba invadens*, and *Acanthamoeba castellanii*. Among the 10 species of Excavata with sequenced genomes (Table S1), *TPS* genes were found only in *Naegleria gruberi* (Fig. 1). A search against the nonredundant (nr) database of National Center for Biotechnology Information (NCBI) ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)) revealed that *TPS* genes are also present in two additional species of eukaryotes, *Polysphondylium pallidum* and *Actyostelium subglobosum*, both of which are classified as Amoebozoa (Fig. 1). Both genera *Polysphondylium* and *Dictyostelium* belong to the family Dictyosteliidae, whereas *A. subglobosum* is a species in the Actyosteliidae family. Both Dictyosteliidae and Actyosteliidae belong to the same order of Dictyosteliida (18). It should be noted that all of the six eukaryotic species that were found to contain *TPS* genes in this study are amoebae (19), with the five species from Dictyosteliida belonging to the unique group of multicellular social amoebae (18).


There are significant variations in the number of *TPS* genes found in the genome of each of the five Dictyosteliida species. Whereas *A. subglobosum* contains a single *TPS* gene, the four species from Dictyosteliidae possess small gene families ranging from three *TPS* genes in *D. fasciculatum* to 21 *TPS* genes in *P. pallidum* (Table S2). The amoeba *N. gruberi* contains seven *TPS* genes. The number of introns for the *TPS* genes from Dictyosteliida ranges between 0 and 3, whereas in contrast, all of the *TPS* genes from *N. gruberi* are intronless (Table S2).

**Identified Eukaryotic Terpene Synthases: Evolutionary Relatedness and Motifs.** To understand the evolutionary relatedness of the identified eukaryotic *TPS*s with known *TPS*s, a phylogenetic tree was constructed that includes, besides the eukaryotic *TPS*s described here, representative bacterial and fungal *TPS*s, and microbial type *TPS*s from the lycophyte *Selaginella moellendorffii* (8). Notably, the *TPS*s from the five species of Dictyosteliida clustered together (clade I), whereas the seven *TPS*s from *N. gruberi* clustered in a separate, but closely related clade (clade II) (Fig. 2). Together, the amoebal *TPS*s showed closer relatedness to fungal *TPS*s than to bacterial *TPS*s (Fig. 2).

*TPS*s contain several highly conserved motifs that are important for catalytic activity including the aspartate-rich “DDxx(x)D/E” motif and the “NDxxSxxxD/E” motif, both of which are involved in complexing metal ions to coordinate the binding of the isoprenyl diphosphate substrate in the active site (20, 21). Both motifs are also highly conserved among all newly identified eukaryotic *TPS*s (Table S3). In addition, the diphosphate sensor that is involved in substrate recognition and critical for catalytic activity (Arginine) (6, 22) was also highly conserved (Table S3).

**Expression Patterns of Individual Terpene Synthase Genes in *Dictyostelium discoideum*.** *D. discoideum* was selected as a model system to explore the function of the newly identified eukaryotic *TPS*s. As a social amoeba, *D. discoideum* has a distinctive life cycle (Fig. S1). It propagates vegetatively as a unicellular organism when food (bacteria in the natural environment) is abundant. Upon starvation, *D. discoideum* transitions into multicellular development in a highly coordinated process that causes individual cells to aggregate and differentiate with formation of a multicellular slug that migrates and finally turns into a fruiting body. This process lasts approximately 24 h (23).

The *D. discoideum* genome contains 11 putative *TPS* genes, 9 of which show a full-length sequence and were designated *DdTPS1* to *DdTPS9*. Analysis of the published gene expression



**Fig. 2.** Phylogenetic reconstruction of newly identified eukaryotic *TPS*s with known *TPS*s. The set of known *TPS*s includes representative *TPS*s from fungi and bacteria. Also included were the microbial terpene synthase-like proteins identified from the plant *Selaginella moellendorffii*. The newly identified eukaryotic *TPS*s include a total of 50 putative full-length *TPS*s identified from six species (five species from Dictyosteliida and *N. gruberi*) (Table S2). *TPS*s are color-coded based on their source. The *TPS*s from Dictyosteliida form clade I and the *TPS*s from *N. gruberi* form clade II.

dataset of *D. discoideum* (24) revealed that each *TPS* gene of *D. discoideum* was expressed at different times during multicellular development (Fig. 3). Specifically, as *D. discoideum* began to starve (0 h), mRNAs of *DdTPS1*, *DdTPS2*, *DdTPS3*, *DdTPS6*, and *DdTPS7* were present at detectable but low levels, whereas mRNAs of *DdTPS4*, *DdTPS5*, *DdTPS8*, and *DdTPS9* were almost undetectable, but in all cases, the mRNA abundance increased during development. During mound formation (approximately 8–12 h), *DdTPS3* attained the highest level of expression among all nine genes, whereas *DdTPS1* levels were still near the limit of detection. At the time of slug formation (approximately 16 h), *DdTPS2* and *DdTPS8* mRNAs reached their highest abundance, whereas the abundance of *DdTPS3* started to decrease. Finally, during culmination (from 18 to 24 h), *DdTPS2*, *DdTPS3*, *DdTPS5*, and *DdTPS8* levels decreased, whereas *DdTPS1*, *DdTPS4*, *DdTPS6*, and *DdTPS9* mRNAs accumulated to higher levels, reaching their peaks at the ultimate stage of mature fruiting body (24 h).


**DdTPS Genes in *Dictyostelium discoideum* Encode Active Terpene-Producing Enzymes.** To further understand the function of *D. discoideum* terpene synthase genes, we characterized the biochemical activities of the enzymes they encode. Full-length cDNAs of *DdTPS1*–*DdTPS9* were cloned and heterologously expressed in *Escherichia coli*. Individual recombinant DdTPS proteins were tested for terpene synthase activity. All nine enzymes were able to accept farnesyl diphosphate (FPP) as a substrate to produce either a single sesquiterpene or a mixture of compounds (Fig. 4). The major products of *DdTPS1*, *DdTPS4*, *DdTPS5*, and *DdTPS7/9* were identified as (*E,E*)- $\alpha$ -farnesene,

(*E*)-nerolidol, (*E*)- $\beta$ -farnesene, and  $\beta$ -barbatene, respectively, whereas *DdTPS2*, *DdTPS3*, *DdTPS6*, and *DdTPS8* produced unidentified sesquiterpenes. *DdTPS1*, *DdTPS2*, *DdTPS3*, and *DdTPS9* were also able to convert geranyl diphosphate (GPP) into different mixtures of monoterpenes (Fig. S2). In addition, diterpene products from geranylgeranyl diphosphate (GGPP) could be observed for *DdTPS1*, *DdTPS2*, *DdTPS3*, *DdTPS4*, *DdTPS5*, and *DdTPS9* (Fig. S3).

***Dictyostelium discoideum* Emits Terpene-Dominated Volatiles During Multicellular Development.** The elaborate temporal regulation of mRNA abundance during different life stages suggests that the *DdTPS* genes may play a role in development. Monoterpenes and sesquiterpenes are generally volatile compounds that can be trapped by use of, e.g., a closed-loop stripping apparatus or solid-phase microextraction (SPME) and analyzed by gas chromatography/mass spectrometry (GC/MS) (25). Based on the expression patterns of individual *DdTPS* genes (Fig. 3) and the *in vitro* biochemical activities of the respective proteins (Fig. 4 and Figs. S2 and S3), we hypothesized that DdTPSs might be involved in producing volatile compounds during development. To test this hypothesis, we performed volatile profiling of *D. discoideum* cultures at 4-h intervals during the 24 h of development (Fig. 5).

Altogether, a total of 15 volatile compounds were detected from developing *D. discoideum* cultures (Fig. 5), including 11 terpenes and four nonterpene volatiles, one of which was identified as 2-phenylethanol. The terpene portion of the volatiles was dominated by nine detectable sesquiterpenes, of which four were identified as (*E,E*)- $\alpha$ -farnesene, calarene, (*E*)-nerolidol, and  $\beta$ -barbatene (Fig. 5). Comparison of the mass spectra of the unidentified sesquiterpenes in the headspace extracts to those obtained enzymatically with the expressed DdTPSs allowed assignment of each of the compounds emitted by *D. discoideum* with confidence to a specific DdTPS. As such, each sesquiterpene in Fig. 5 was labeled with the same peak number as used in Fig. 4. In addition, the monoterpene linalool and one diterpene, which was identical to *in vitro* diterpene product of *DdTPS5* (Fig. S3), were detected in the culture extracts.

The relative abundance of individual volatile terpenes during the 24 h of development was calculated based on three biological replicates (Fig. 5 and Fig. S4). At the beginning of development (0 h), essentially no volatile terpenes were detected, whereas after 4 h of development, the emission of traces of terpenes including (*E*)-nerolidol was detected. The production of terpenes by *D. discoideum* gradually increased during the next hours of development, but some compounds showed an early maximum production, e.g., calarene peaked at 12 h and 16 h, whereas the production of other terpenes such as (*E,E*)- $\alpha$ -farnesene and  $\beta$ -barbatene exhibited a later maximum of production.




**Fig. 3.** Expression patterns of nine terpene synthase genes in *D. discoideum* (*DdTPS1*–*9*). This analysis was based on published RNAseq data (24), which were obtained at seven time points during a complete developmental program in which individual *D. discoideum* cells aggregated and differentiated, forming a multicellular slug that migrated and then formed a fruiting body in a highly coordinated process that lasted approximately 24 h. The expression levels of nine *DdTPS* genes were measured by RPKM (reads per kilobase per million sequenced reads) and then displayed on a  $\log_2(\text{RPKM}+1)$  scale in this line plot. The line plot shows the transcript abundance (y axis;  $\log_2$ -scale) of nine *DdTPS* genes. The cartoons depict various stages during multicellular development: vegetative, individual cells (0 h), streaming (8 h), loose aggregate (10 h), tipped aggregate (14 h), slug (16 h), Mexican hat (20 h), and fruiting bodies (24 h).

## Discussion

This first report of the occurrence of canonical terpene synthase genes in the social amoebae raises questions about the functions of the terpene products in these organisms. The fact that *TPS* gene expression and terpene volatile emission in *D. discoideum* are restricted to specific periods during multicellular development suggests possible roles for these compounds if the unique biology of *D. discoideum* is considered in light of the known functions of volatile terpenes in other organisms.

One possible function of volatile terpenes emitted from *D. discoideum* is to attract other organisms to facilitate spore dispersal, resembling the function of volatile terpenes from the fruiting bodies of fungi (26). Forming fruiting bodies by social amoebae is considered to be an adaptation for spore dispersal (27). This hypothesis was directly supported by experimental studies in which fruiting bodies were shown to increase the rate at which spores are acquired by a model invertebrate *Drosophila*



**Fig. 4.** Sesquiterpene synthase activity of *D. discoideum* terpene synthases. Genes were heterologously expressed in *E. coli*, and crude protein extracts were incubated with the substrate FPP. Enzyme products were collected by using solid-phase microextraction and analyzed by GC/MS. GC traces (Left) and mass spectra of major products (Right) are shown. 1, (*E,E*)- $\alpha$ -farnesene\*; 2, unidentified sesquiterpene hydrocarbon; 3,  $\beta$ -maaliene; 4, aristolene; 5, calarene; 6, unidentified sesquiterpene hydrocarbon; 7, unidentified sesquiterpene hydrocarbon; 8, unidentified sesquiterpene hydrocarbon; 9, (*E*)-nerolidol\*; 10,  $\beta$ -elemene\*; 11, (*E*)- $\beta$ -farnesene\*; 12, unidentified sesquiterpene hydrocarbon; 13,  $\beta$ -barbatene\*; 14, unidentified sesquiterpene; cont, contamination. Compounds marked with asterisks (\*) were identified by using authentic standards. Each assay was repeated at least three times, and a representative GC chromatogram is shown.


*melanogaster* (27). Although the primary vectors for *D. discoideum* spore dispersal are unknown (27), it will be an interesting future subject to identify such vectors in nature and then determine whether volatile terpenes have a role in attracting such vectors to facilitate spore dispersal. Consistent with this hypothesis,  $\beta$ -barbatene (Fig. 5) emitted from the fruiting bodies of the bracket fungus *Fomitopsis pinicola* has been implicated in attracting insects for spore dispersal (26).

Another possible function of *D. discoideum* volatile terpenes is defense. Social amoebae are preyed on by nematodes (28). They have evolved multiple defense mechanisms, which include the synthesis of a protective extracellular matrix called the slime sheath and the formation of protective coats at the surface of spores (28). In addition, individual amoebae protect themselves by secreting compounds that repel nematodes (28). It will be interesting to see whether any of the volatile terpenes emitted from *D. discoideum* serve such a function as well. Consistent with this hypothesis, (*E,E*)- $\alpha$ -farnesene (Fig. 5) emitted from the leaves

of the model plant *Arabidopsis thaliana* has been implicated in defense against insects (29).

The third possibility is that *D. discoideum* terpenes may function as signals to coordinate multicellular development. The roles of terpenes in signaling have been relatively well-studied in plants (30). As volatile compounds, terpenes can signal over a distance in either multicellular organisms or multicellular aggregates. Previous studies showed that ammonia, a volatile by-product of gluconeogenesis, is involved in regulating several stages of *D. discoideum* development, including aggregation (31), slug migration (32), and culmination (33). It is certainly intriguing to ask whether volatile terpenes could have similar functions, with the diversity of chemical structures helping to provide functional specificity.

It is interesting that all of the five species of social amoebae (i.e., all from Dictyosteliida) with sequenced genomes contain *TPS* genes, whereas the three species from the genus *Entamoeba* (Table S1 and Fig. 1), which also belong to the supergroup Amoebozoa, do not contain any *TPS* gene. The genus *Entamoeba*



**Fig. 5.** *D. discoideum* emits diverse volatile terpenes during development. The beginning of multicellular development was set as 0 h. Volatiles were collected once every 4 h. Shown are GC/MS chromatograms from one representative replicate. All of the peaks labeled with a number are sesquiterpenes, and the number corresponds to the number of the peak in Fig. 4. 1, (*E,E*)- $\alpha$ -farnesene\*; 2, unidentified sesquiterpene hydrocarbon; 5, calarene; 6, unidentified sesquiterpene hydrocarbon; 7, unidentified sesquiterpene hydrocarbon; 8, unidentified sesquiterpene hydrocarbon; 9, (*E*)-nerolidol\*; 12, unidentified sesquiterpene hydrocarbon; 13,  $\beta$ -barbatene\*. Compounds marked with asterisks (\*) were identified by using authentic standard. "li" represents the monoterpene linalool, and "di" represents an unidentified diterpene. The letters indicate nonterpene volatiles. "a" represents 2-phenylethanol; "b", "c", and "d" represent unidentified compounds.

are amoebae but not social amoebae, suggesting that *TPS* genes may provide adaptive functions for social amoeba, which share a unique lifestyle. However, *N. gruberi* of the supergroup Excavata, which is also an amoeba having no multicellular development (34), contains *TPS* genes (Fig. 1). Sharing similarity in lifestyle, *N. gruberi* and *Entamoeba* are evolutionarily distantly related. It will be interesting to investigate the biochemical and biological function of *N. gruberi* *TPS* genes and to understand how they may confer a fitness advantage.

To briefly summarize, we have found classic terpene synthase genes in six species of amoebae among a broad range of non-plant/nonfungus eukaryotes. Amoebal TPSs are more closely related to fungal TPSs and the microbial type TPSs from plants than bacterial TPSs (Fig. 2). The social amoeba *D. discoideum* is an organism other than plants, fungi, and bacteria from which classic terpene synthase genes have been functionally characterized (Fig. 4). This study provides insights into the occurrence, function and evolution of *TPS* genes, particularly in eukaryotes, and it is expected to stimulate important future research.

## Materials and Methods

**Sequence Retrieval and Analysis.** A total of 168 species of nonplant/nonfungus eukaryotes with well-annotated genome sequences (Table S1) archived at the KEGG genome database ([www.genome.jp/kegg/catalog/org\\_list.html](http://www.genome.jp/kegg/catalog/org_list.html)) were downloaded as the genome dataset. Another dataset was the non-redundant (nr) protein database from NCBI, which was downloaded on April 19, 2016. Both databases were searched against the Pfam-A database locally by using HMMER 3.0 with an e-value of  $1e^{-2}$ . Sequences with best hits from the following three HMM profiles were identified as putative terpene

synthases: Terpene\_synth\_C (PF03936) and Terpene Synthase N-terminal domain (PF01397), and TRI5 (PF06330). For the search of the nr database, the terpene synthase hits identified from plants, fungi, archaea, and bacteria were removed. For phylogenetic reconstruction, known bacterial and fungal terpene synthases were retrieved from Pfam database (version 27). MAFFT (L-INS-i) was used to build the multiple sequence alignment with 1,000 iterations of improvement. The maximum-likelihood phylogenetic tree was built with RAxML through the CIPRES Science Gateway (<https://www.phylo.org>) by using the LG+G+F amino acid substitution model with 1,000 bootstrap replicates and then rendered by using FigTree (version 1.4.2).

## Cloning of Full-Length cDNA of *DdTPS* Genes of *Dictyostelium discoideum* via RT-PCR.

*D. discoideum* (strain AX4) was obtained from the Dictybase Stock Center ([www.dictybase.org](http://www.dictybase.org)). *D. discoideum* was cocultured with live *Klebsiella pneumoniae* bacteria on SM agar plates (35) by following the protocol described in the Dictybase Stock Center. When slugs formed, *D. discoideum* cells were collected and used for total RNA isolation following the protocol described by ref. 36. Full-length cDNA of each *DdTPS* gene was amplified by RT-PCR using the gene specific primers listed in Table S4. PCR products were cloned into the pEXP-5-CT/TOPO vector (ThermoFisher Scientific) and confirmed by sequencing.

**Terpene Synthase Enzyme Assays.** Heterologous expression of *DdTPS* genes in *E. coli*, recombinant protein preparation, terpene synthase enzyme assays, and terpene product identification using a Hewlett-Packard 6890 gas chromatograph coupled to a Hewlett-Packard 5973 mass spectrometer were performed as described (8). Each expressed protein was assayed at least three times.

***DdTPS* Gene Expression Analysis.** *DdTPS* gene expression was analyzed by using published RNA-seq data from developing *D. discoideum* (24). Transcript abundance was quantified as described in ref. 37. The data are available on

dictyExpress (38). Briefly, *D. discoideum* was grown on a thick lawn of *K. aerogenes* before the food source was removed to initiate the developmental program. The samples were obtained in 4-h intervals throughout the 24-h developmental program.

**Headspace Collection and GC/MS Analysis.** A mixture of *D. discoideum* spores and freshly grown *K. pneumoniae* was spread onto 5M agar plates to initiate *D. discoideum* culture. Under our experimental conditions, *D. discoideum* progressed from spore germination to vegetative growth to the completion of multicellular development in ~48 h. At 24 h, the appearance of the culture plate changed from opaque (from the bacterial lawn) to translucent, indicating the clearing of bacteria. This time point was defined as the start of multicellular development, after which *D. discoideum* progressed through the various described developmental stages in the next 24 h with the eventual formation of fruiting bodies (Fig. S1). SPME combined with GC/MS was used for volatile profiling of the *D. discoideum* cultures. During the 24 h of development, volatiles were collected once every 4 h (Fig. 5). Before each collection, the lid of the culture plate was removed and the plate was left in

the hood for 1 min to dispose of accumulated volatiles. Then the lid was put back on and a SPME fiber coated with 100- $\mu$ m polydimethylsiloxane was inserted into the headspace of the plate to start volatile collection. After 1 h, the SPME fiber was retracted and inserted into the injector port (a splitless injection and injector temperature of 250 °C) of a Shimadzu 17A gas chromatograph coupled to a Shimadzu QP5050A quadrupole mass selective detector for chemical identification. Separation was performed on a Restek Rxi-5Sil MS column (30 m  $\times$  0.25 mm i.d.  $\times$  0.25  $\mu$ m thickness; Restek) with helium as the carrier gas and a temperature program from 60 °C to 300 °C at 5 °C $\cdot$ min<sup>-1</sup>. The experiment was performed with three biological replicates.

**ACKNOWLEDGMENTS.** We thank Wangdan Xiong for her assistance with the preparation of Fig. 3 and Fig. S1 and Natascha Rauch (Max Planck Institute for Chemical Ecology) for her technical assistance. This project was supported by a University of Tennessee AgResearch Innovation Grant (to F.C.) and the Max Planck Society (T.G.K. and J.G.). G.S. and B.S. acknowledge support from National Institutes of Health Grant P01 HD039691.

- Eisenreich W, Bacher A, Arigoni D, Rohdich F (2004) Biosynthesis of isoprenoids via the non-mevalonate pathway. *Cell Mol Life Sci* 61(12):1401–1426.
- Jia Q, Chen F (2016) Catalytic functions of the isoprenyl diphosphate synthase superfamily in plants: A growing repertoire. *Mol Plant* 9(2):189–191.
- Wang KC, Ohnuma S (2000) Isoprenyl diphosphate synthases. *Biochim Biophys Acta* 1529(1-3):33–48.
- Degenhardt J, Köllner TG, Gershenzon J (2009) Monoterpene and sesquiterpene synthases and the origin of terpene skeletal diversity in plants. *Phytochemistry* 70(15-16):1621–1637.
- Yamada Y, et al. (2015) Terpene synthases are widely distributed in bacteria. *Proc Natl Acad Sci USA* 112(3):857–862.
- Dickschat JS (2016) Bacterial terpene cyclases. *Nat Prod Rep* 33(1):87–110.
- Chen F, Tholl D, Bohlmann J, Pichersky E (2011) The family of terpene synthases in plants: A mid-size family of genes for specialized metabolism that is highly diversified throughout the kingdom. *Plant J* 66(1):212–229.
- Li G, et al. (2012) Nonseed plant *Selaginella moellendorffii* [corrected] has both seed plant and microbial types of terpene synthases. *Proc Natl Acad Sci USA* 109(36):14711–14715.
- Schmidt-Dannert C (2015) Biosynthesis of terpenoid natural products in fungi. *Adv Biochem Eng Biotechnol* 148:19–61.
- Keller NP, Turner G, Bennett JW (2005) Fungal secondary metabolism - from biochemistry to genomics. *Nat Rev Microbiol* 3(12):937–947.
- Gilg AB, Tittiger C, Blomquist GJ (2009) Unique animal prenyltransferase with monoterpene synthase activity. *Naturwissenschaften* 96(6):731–735.
- Beran F, et al. (2016) Novel family of terpene synthases evolved from trans-isoprenyl diphosphate synthases in a flea beetle. *Proc Natl Acad Sci USA* 113(11):2922–2927.
- Cao R, et al. (2010) Diterpene cyclases and the nature of the isoprene fold. *Proteins* 78(11):2417–2432.
- Adl SM, et al. (2012) The revised classification of eukaryotes. *J Eukaryot Microbiol* 59(5):429–493.
- Burki F (2014) The eukaryotic tree of life from a global phylogenomic perspective. *Cold Spring Harb Perspect Biol* 6(5):a016147.
- Gershenzon J, Dudareva N (2007) The function of terpene natural products in the natural world. *Nat Chem Biol* 3(7):408–414.
- Finn RD, Clements J, Eddy SR (2011) HMMER web server: Interactive sequence similarity searching. *Nucleic Acids Res* 39(Web Server issue):W29–37.
- Schilde C, Schaap P (2013) The Amoebozoa. *Methods Mol Biol* 983:1–15.
- Pawlowski J, Burki F (2009) Untangling the phylogeny of amoeboid protists. *J Eukaryot Microbiol* 56(1):16–25.
- Starks CM, Back K, Chappell J, Noel JP (1997) Structural basis for cyclic terpene biosynthesis by tobacco 5-epi-aristolochene synthase. *Science* 277(5333):1815–1820.
- Christianson DW (2008) Unearthing the roots of the terpenome. *Curr Opin Chem Biol* 12(2):141–150.
- Baer P, et al. (2014) Induced-fit mechanism in class I terpene cyclases. *Angew Chem Int Ed Engl* 53(29):7652–7656.
- Kessin RH (2001) *Dictyostelium: Evolution, Cell Biology, and the Development of Multicellularity* (Cambridge Univ Press, Cambridge, UK).
- Parikh A, et al. (2010) Conserved developmental transcriptomes in evolutionarily divergent species. *Genome Biol* 11(3):R35.
- Dickschat JS (2014) Capturing volatile natural products by mass spectrometry. *Nat Prod Rep* 31(6):838–861.
- Fäldt J, Jonsell M, Nordlander G, Borg-Karlson A-K (1999) Volatiles of bracket fungi *Fomitopsis pinicola* and *Fomes fomentarius* and their functions as insect attractants. *J Chem Ecol* 25(3):567–590.
- Smith J, Queller DC, Strassmann JE (2014) Fruiting bodies of the social amoeba *Dictyostelium discoideum* increase spore transport by *Drosophila*. *BMC Evol Biol* 14:105.
- Kessin RH, Gundersen GG, Zaydfudim V, Grimson M (1996) How cellular slime molds evade nematodes. *Proc Natl Acad Sci USA* 93(10):4857–4861.
- Huang M, et al. (2010) Variation of herbivore-induced volatile terpenes among *Arabidopsis* ecotypes depends on allelic differences and subcellular targeting of two terpene synthases, TPS02 and TPS03. *Plant Physiol* 153(3):1293–1310.
- Maag D, Erb M, Köllner TG, Gershenzon J (2015) Defensive weapons and defense signals in plants: Some metabolites serve both roles. *BioEssays* 37(2):167–174.
- Bonner JT, Suthers HB, Odell GM (1986) Ammonia orients cell masses and speeds up aggregating cells of slime molds. *Nature* 323(6089):630–632.
- Bonner JT, Chiang A, Lee J, Suthers HB (1988) The possible role of ammonia in phototaxis of migrating slugs of *Dictyostelium discoideum*. *Proc Natl Acad Sci USA* 85(11):3885–3887.
- Feit IN, Sollitto RB (1987) Ammonia is the gas used for the spacing of fruiting bodies in the cellular slime mold, *Dictyostelium discoideum*. *Differentiation* 33(3):193–196.
- Fritz-Laylin LK, et al. (2010) The genome of *Naegleria gruberi* illuminates early eukaryotic versatility. *Cell* 140(5):631–642.
- Sussman M (1966) Biochemical and genetic methods in the study of cellular slime mold development. *Methods Cell Biol* 2:397–410.
- Pilcher KE, Gaudet P, Fey P, Kowal AS, Chisholm RL (2007) A general purpose method for extracting RNA from *Dictyostelium* cells. *Nat Protoc* 2(6):1329–1332.
- Miranda ER, et al. (2013) Transcriptional profiling of *Dictyostelium* with RNA sequencing. *Methods Mol Biol* 983:139–171.
- Rot G, et al. (2009) dictyExpress: A *Dictyostelium discoideum* gene expression database with an explorative data analysis web-based interface. *BMC Bioinformatics* 10:265.