

# Mammal domestication and the symbiotic spectrum

Yaron Dekel<sup>a,b,1</sup>, Yossy Machluf<sup>c</sup>, Rachel Brand<sup>a,b</sup>, Oshrat Noked Partouche<sup>a,d</sup>, Izhar Ben-Shlomo<sup>e</sup>, and Dani Bercovich<sup>d</sup>

In their paper in PNAS, Weissbrod et al. (1) conclude that mice adapted to the temporary sedentary lifestyle of humans during the Late Pleistocene and underwent a domestication process driven by a competitive advantage for commensal mice in the evolving, long-term human settlements. The authors expand the possibility of an early interaction between wild mammals and humans to other wild species, wild boars and wolves, as examples of “following commensal pathways before developing long-term mutualism characteristic of domestication” (1).

We wish to deliberate on both the process and its nature, with regard to these species.

During the Late Pleistocene, wild boars were captured and managed by humans for consumption. Fully domesticated pigs emerged only later, ~9,000 y ago, after the transition to a sedentary agricultural lifestyle (2). Archeological records and genetic evidence suggest that dog domestication took place at least 15,000 y ago, or even as early as 33,000 y ago (3). However, the first domesticated grain crop, wheat, was domesticated 10,000 y ago (4). These two notions suggest that proto-dogs coexisted with humans thousands of years before the agricultural revolution (5). Nevertheless, fossils of domesticated mammals that evolved from wild herbivores were dated to only after the agricultural revolution (6, 7).

Domestication, as an accelerated evolutionary process, resulted from natural selection by a human-induced

environment. The genetic tendency of wolves (8, 9) and wild mice to be less nervous of nearby humans probably enabled them both to get closer initially and to become our pets and pests, respectively, later. The process was driven by human waste, such as bones and skin, which was available in short-term mobile human settlements as well as in more sedentary ones. Similarly, after the agricultural revolution and the appearance of domesticated wheat/barley fields in the Fertile Crescent (10), the less fearful wild herbivores approached the grain fields and later became our sheep, goats, cows, camels, and horses (6, 7). Thus, domestication might have been independent (dogs/mice) or dependent (herbivores) on the agricultural revolution timeline.

It is highly plausible that mice, like dogs, followed humans as fully domesticated animals. Dogs, as sanitation agents, had a relationship with humans that can be described as mutualism whereas mice, on the symbiotic spectrum, were probably parasites, eating humans' food, damaging their belongings, and traveling hidden in their ancient luggage.

In conclusion, adjusting to the nomadic/semisedentary human niche was the first driving force of domestication. Both dogs and mice underwent this process, but they differ from each other in the initial nature of their relationship with humans: mutualism and parasitism, respectively, rather than commensalism.

- 1 Weissbrod L, et al. (2017) Origins of house mice in ecological niches created by settled hunter-gatherers in the Levant 15,000 y ago. *Proc Natl Acad Sci USA* 114:4099–4104.
- 2 Vigne J-D, et al. (2009) Pre-Neolithic wild boar management and introduction to Cyprus more than 11,400 years ago. *Proc Natl Acad Sci USA* 106:16135–16138.
- 3 Larson G, et al. (2012) Rethinking dog domestication by integrating genetics, archeology, and biogeography. *Proc Natl Acad Sci USA* 109:8878–8883.
- 4 King J, et al. (2017) A step change in the transfer of interspecific variation into wheat from *Amblyopyrum muticum*. *Plant Biotechnol J* 15:217–226.
- 5 Freedman AH, et al. (2014) Genome sequencing highlights the dynamic early history of dogs. *PLoS Genet* 10:e1004016.
- 6 Taberlet P, Coissac E, Pansu J, Pompanon F (2011) Conservation genetics of cattle, sheep, and goats. *C R Biol* 334:247–254.
- 7 Larson G, et al. (2007) Ancient DNA, pig domestication, and the spread of the Neolithic into Europe. *Proc Natl Acad Sci USA* 104:15276–15281.
- 8 Trut LN (1999) Early canid domestication: The farm-fox experiment. *Am Sci* 87:160–169.
- 9 Marshall-Pescini S, Virányi Z, Range F (2015) The effect of domestication on inhibitory control: Wolves and dogs compared. *PLoS One* 10:e0118469.
- 10 Hewson CJ (2003) Dogs: A new understanding of canine origin, behavior, and evolution. *Appl Anim Behav Sci* 83:77–78.

<sup>a</sup>Unit of Agrigenomics, Shamir Research Institute, University of Haifa, Kazerim 1290000, Israel; <sup>b</sup>Department of Medical Laboratory Sciences, Zefat Academic College, Zefat 1320611, Israel; <sup>c</sup>Private address, Rehovot 7661509, Israel; <sup>d</sup>Faculty of Biotechnology, Tel-Hai College, Upper Galilee 1220800, Israel; and <sup>e</sup>Unit of Obstetrics and Gynecology, Faculty of Medicine in the Galilee, Bar-Ilan University & Baruch Padeh Medical Center, Poriya M.P. The Lower Galilee 15208, Israel  
 Author contributions: Y.D., Y.M., I.B.-S., and D.B. designed research; R.B. and O.N.P. performed research; and Y.D. and Y.M. wrote the paper.  
 The authors declare no conflict of interest.

<sup>1</sup>To whom correspondence should be addressed. Email: yarond@gri.org.il.