

Molecular hijacking of siroheme for the synthesis of heme and d_1 heme

Shilpa Bali^{a,b}, Andrew D. Lawrence^b, Susana A. Lobo^{b,c}, Lúcia M. Saraiva^c, Bernard T. Golding^d, David J. Palmer^b, Mark J. Howard^b, Stuart J. Ferguson^{a,1}, and Martin J. Warren^{b,1}

^aDepartment of Biochemistry, University of Oxford, South Parks Road, Oxford OX1 3QU, United Kingdom; ^bDepartment of Biosciences, University of Kent, Canterbury, Kent CT2 7NZ, United Kingdom; ^cInstituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa, Avenida da República, Estação Agronómica Nacional, 2780-157 Oeiras, Portugal; and ^dSchool of Chemistry, Newcastle University, Bedson Building, Newcastle upon Tyne NE1 7RU, United Kingdom

Edited by Rowena G. Matthews, University of Michigan, Ann Arbor, MI, and approved August 18, 2011 (received for review May 25, 2011)

Modified tetrapyrroles such as chlorophyll, heme, siroheme, vitamin B₁₂, coenzyme F₄₃₀, and heme d_1 underpin a wide range of essential biological functions in all domains of life, and it is therefore surprising that the syntheses of many of these life pigments remain poorly understood. It is known that the construction of the central molecular framework of modified tetrapyrroles is mediated via a common, core pathway. Herein a further branch of the modified tetrapyrrole biosynthesis pathway is described in denitrifying and sulfate-reducing bacteria as well as the Archaea. This process entails the hijacking of siroheme, the prosthetic group of sulfite and nitrite reductase, and its processing into heme and d_1 heme. The initial step in these transformations involves the decarboxylation of siroheme to give didecarboxysiroheme. For d_1 heme synthesis this intermediate has to undergo the replacement of two propionate side chains with oxygen functionalities and the introduction of a double bond into a further peripheral side chain. For heme synthesis didecarboxysiroheme is converted into Fe-coproporphyrin by oxidative loss of two acetic acid side chains. Fe-coproporphyrin is then transformed into heme by the oxidative decarboxylation of two propionate side chains. The mechanisms of these reactions are discussed and the evolutionary significance of another role for siroheme is examined.

enzymes | metabolic pathway | S-adenosylmethionine

An understanding of biochemical pathways involves a comprehension of the chemical logic underpinning the synthetic process. Biochemical reactions are, in essence, highly controlled and regulated chemical reactions that can also provide insights into transformations for which no laboratory-based chemistry may yet have been described. Such studies are at the heart of chemical biology and are an essential prerequisite for its transformation into synthetic biology. Moreover, from an evolutionary perspective, there are many interesting questions relating to the initial appearance of complex multistep pathways, where theories on retro and patchwork models have been discussed (1, 2). These considerations are pertinent to the synthesis of the modified tetrapyrroles (3), which through their representatives chlorophyll, heme, cobalamin (the biological form of vitamin B₁₂), siroheme, heme d_1 , and coenzyme F₄₃₀ are involved in a broad variety of essential life processes from photosynthesis to methane production.

The tetrapyrrolic architecture that underpins the structural similarity between these various metalloprosthetic groups results from a shared, though branched, biosynthetic pathway (4, 5), where the whole family is derived from the macrocyclic primogenitor, uroporphyrinogen III (Fig. 1A). The pathways to heme and chlorophyll via protoporphyrin have been previously characterized. Similarly, the syntheses of siroheme and cobalamin have also been elucidated, progressing via the bis-methylated uroporphyrinogen III derivative precorrin-2 (Fig. 1A). However, the biosynthesis of coenzyme F₄₃₀ and heme d_1 have not been reported. Likewise, little is known about a proposed alternative

heme biosynthetic route that involves the transformation of precorrin-2 into heme (6). This paper focuses on this unique heme biosynthetic process and reveals a hitherto unsuspected relationship with both the synthesis of siroheme and heme d_1 .

Siroheme, the prosthetic group of sulfite and nitrite reductases, is synthesized from uroporphyrinogen III by bis-methylation to give precorrin-2, dehydrogenation to produce sirohydrochlorin and finally ferrochelation (7) (Fig. 1B). Heme d_1 is only made by denitrifying bacteria with a cytochrome nitrite reductase *cd₁* (*nirS*) for which d_1 acts as an essential cofactor, tailored to meet the mechanistic requirements of the reaction (8). Bacteria with *nirS* always appear to contain, downstream of *nirS*, a set of contiguous genes necessary for heme d_1 biogenesis (9). In *Paracoccus pantotrophus* and *Paracoccus denitrificans*, these genes are organized in the operon *nirECFD-LGHJN* (*nirD* and *nirL* occur as a fused gene in *Paracoccus* species) and are transcribed in the presence of nitric oxide (10). This operon is thought to encode all the enzymes for heme d_1 biogenesis. So far only the function of NirE has been unambiguously determined as an S-adenosylmethionine (AdoMet)-dependent uroporphyrinogen methyltransferase (11, 12), a finding that is consistent with previous labeling studies that demonstrate the synthesis of d_1 heme must proceed via precorrin-2 (Fig. 1A and C) (13).

Heme requires little introduction as a prosthetic group and has a diverse range of biological functions. Its synthesis has been elucidated in eukaryotes and most bacteria, where the classic pathway for its biosynthesis is advanced via an ordered and sequential decarboxylation of the majority of the acidic side chains of uroporphyrinogen III, followed by oxidation and ferrochelation (4) (Fig. 1B). As alluded to earlier, in sulfate-reducing bacteria and Archaea, it is proposed that heme is made via a completely different pathway, herein called the alternative heme biosynthesis route. A combination of biochemistry and bioinformatics approaches has provided some clues as to the mode of operation of this route. Thus, feeding labeled methionine to *Desulfovibrio vulgaris* cultures resulted in the isolation of labeled sirohydrochlorin, 12,18-didecarboxysirohydrochlorin, coproporphyrin III and protoporphyrin IX (6). Similarly, analysis of completely sequenced Archaeal and *Desulfovibrio* genomes suggest that these organisms possess the genes that encode enzymes for the transformation of 5-aminolevulinic acid to uroporphyrinogen but lack those required for the enzymes necessary for the conversion of uroporphyrinogen into protoheme through the classical

Author contributions: S.B., A.D.L., S.A.L., L.M.S., S.J.F., and M.J.W. designed research; S.B., A.D.L., S.A.L., D.J.P., and M.J.H. performed research; S.B., A.D.L., S.A.L., B.T.G., and M.J.H. analyzed data; and S.B., L.M.S., B.T.G., S.J.F., and M.J.W. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence may be addressed. E-mail: m.j.warren@kent.ac.uk or stuart.ferguson@bioch.ox.ac.uk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1108228108/-DCSupplemental.




Fig. 1. Pathways and genes of modified tetrapyrrole biosynthesis. (A) Outline syntheses of modified tetrapyrroles highlighting key intermediates along the branched pathway. Known reaction sequences are shown in black whereas those not yet elucidated are highlighted in magenta. (B) The known transformation of uroporphyrinogen III into siroheme and heme is shown. The numbering system for the tetrapyrrole macrocycle is highlighted on uroporphyrinogen III. (C) Precorrin-2 as the template for both d_1 heme and heme synthesis is shown, with the latter derived via the alternative heme pathway. In both cases the two methyl groups at C2 and C7, highlighted in red, are derived from AdoMet. The enzymes involved in the transformation of precorrin-2 into d_1 heme (Nir enzymes) and heme (Ahb enzymes) are shown. (D) The similarity between some of the Nir proteins implicated in d_1 synthesis and proteins found in sulfate-reducing bacteria and Archaea thought to be associated with the Ahb pathway are indicated by black arrows.

pathway (14–17). Altogether, these observations led to the theory of an alternative heme biosynthesis pathway that diverges from the classical pathway at the uroporphyrinogen step via precorrin-2 (Fig. 1A and C).

Our genome comparison of denitrifying bacteria with some sulfate-reducing bacteria and Archaea that make heme via precorrin-2, revealed the presence of homologues of *nirD*, and *nirJ* (see additionally ref. 17). The presence of these orthologues in sulfate-reducing bacteria and Archaea, which do not need d_1 heme as they do not have *cd₁*, was intriguing because it implied that they may have a role in the alternative heme biosynthesis route. The similarity between the d_1 synthesis genes found in *Paracoccus* and *Desulfovibrio* is shown diagrammatically in Fig. 1D, where the alternative heme biosynthesis genes are given the prefix *ahb*. Taken together, these observations prompted us to investigate the link(s) between the alternate heme and d_1 heme biosynthetic pathways.

Results

We speculated that the biosynthesis of heme and heme d_1 proceeds via the oxidized derivative of precorrin-2, sirohydrochlorin, and didecarboxysirohydrochlorin, at which point the pathways for the two cofactors would diverge. In both cases it was envisaged

that iron would be added at the final stage, as is observed with siroheme and the classic heme pathway. To test the hypothesis that heme d_1 synthesis may proceed via sirohydrochlorin, cell-free extracts of *Escherichia coli*, overproducing either individual *P. pantotrophus* Nir proteins (D–L, G, H, J, F) or combinations of them generated using our link and lock technology (18), were incubated with this metal-free substrate. Strikingly, when lysates of *E. coli*, overproducing *P. pantotrophus* NirDL–G–H, were incubated with sirohydrochlorin an instant change in the appearance of the purple sirohydrochlorin to a blue-green color was observed. The UV-visible spectrum of sirohydrochlorin, which has a Soret band at 378 nm, also changed to a new maximum around 390 nm (Fig. 2A). The shift of 12 nm in the Soret of sirohydrochlorin and the disappearance of its d bands in the region of 590 nm suggested that sirohydrochlorin had undergone a chemical transformation and hence this unique compound was further characterized.

NirDL–G–H Carries an Oxygen-Independent Siroheme Decarboxylase Activity. HPLC analysis of the NirDL–G–H catalyzed conversion of sirohydrochlorin showed the presence of two major peaks, whose retention time and m/z values differed from that of sirohydrochlorin (Fig. 2B). Significantly, the isotopic mass distribu-

gesting that the reaction occurs in a stepwise manner (Fig. 2B). In control reactions where either sirohydrochlorin or siroheme were incubated with the cell extracts of *E. coli* carrying empty expression vector-pET3a (Fig. 2B, traces B and D, respectively), no loss of substrate or formation of didecarboxysiroheme was detected. These results provide compelling evidence that the d_1 heme biosynthesis pathway proceeds via siroheme and didecarboxysiroheme (Fig. 3). Although sequence analysis highlighted a significant sequence identity (35.3%) and similarity (47.1%) between pairs of NirD-L, NirG, and NirH, these proteins are, on the basis of mutagenesis, functionally nonredundant (9, 19).




Fig. 3. Deduced pathway from siroheme to d_1 and heme. (A) The corrected modified tetrapyrrole pathway is shown with siroheme now acting as an intermediate for heme and d_1 heme synthesis. This pathway can be compared to Fig. 1C. (B) The branched pathway from siroheme to d_1 and heme is shown together with the enzymes that are thought to be involved with each particular step. Didecarboxysiroheme is generated from siroheme by the sequential decarboxylation of the side chains attached to C12 and C18. For d_1 synthesis, didecarboxysiroheme is modified by replacement of the propionate side chains at C3 and C8 with oxo groups in a reaction likely catalyzed by NirJ to generate a pre- d_1 intermediate, which is converted into d_1 by the introduction of a double bond on the propionate side chain on C17. For heme synthesis, didecarboxysiroheme is modified by the action of AhbC to remove the acetic acid side chains attached to C2 and C7 to give Fe-coproporphyrin. This intermediate is converted into heme by AhbD through the oxidative decarboxylation of propionate side chains on C3 and C8.




Fig. 5. AhbD catalyzed Fe-coproporphyrin oxidase activity. HPLC traces of observed porphyrin derivatives at 390 nm in reaction containing, (A) Fe-coproporphyrin, cofactor mix, and cell-free extract of *E. coli* harboring empty expression vector, (B) Fe-coproporphyrin, cofactor mix, and cell-free extract of *E. coli* overexpressing *D. desulfuricans* AhbD, and (C) Fe-coproporphyrin, cell-free extract containing *D. desulfuricans* AhbD without AdoMet in the cofactor mix. Cofactor mix consisted of 500 μ M AdoMet and NADPH with 0.3% (vol/vol) Triton X-100 in each assay. Fe-coproporphyrin was used at a final concentration of 25 μ M, and is eluted as two close peaks from the HPLC column but with same m/z value of 708 by MS.

generated to act as an electron sink (Fig. S4). Although we have detected the presence of a monodecarboxylated intermediate, the order of decarboxylation has not been determined.

The biosynthesis of d_1 heme therefore proceeds via siroheme and didecarboxysiroheme, and thereafter via a dioxo intermediate before the introduction of an acrylate function as the final step. We believe that the oxo groups are formed by the action of NirJ, a radical AdoMet enzyme (20), and that the double bond in the propionate side chain attached to C17 may be mediated by NirF, which, being periplasmic, must catalyze the last step. We have shown that the periplasmic NirF is required for d_1 heme production whereas two other periplasmic proteins (NirC and NirN) coded for in the biogenesis operon are not (11, 23). Thus we conclude that NirF catalyzes the final step in d_1 production unless the product of NirF activity is imported to the cytoplasm for further processing, an energetically expensive possibility. At this stage we do not know what reaction is catalyzed by NirF but the dehydrogenation to give the acrylate side chain of the isobacteriochlorin is a possibility to be addressed in future work (23). Our result that siroheme is an actual intermediate during d_1 heme biosynthesis means that a ferrochelatase activity is not coded by the *nir* operon whose product must work in concert with a siroheme synthesis pathway. It also shows that avoidance of reactive radicals and potential toxicity by inserting iron at the end of the biosynthetic pathway cannot be as important as previously thought.

The similarity between the d_1 heme biosynthetic genes *nirD-L*, *G*, and *H* and the putative alternative heme biosynthetic genes *ahbA* and *B* immediately suggested that didecarboxysiroheme could also be an intermediate in the latter pathway. This hypothesis was confirmed by the activity of AhbA and AhbB upon incubation with siroheme, which saw the quantitative conversion into didecarboxysiroheme. Moreover, incubation of siroheme with cell-free extracts of *D. vulgaris* resulted in the appearance not only of didecarboxysiroheme but also Fe-coproporphyrin and heme [and it is also interesting to note that Fe-coproporphyrin is found as a cofactor in the bacterioferritin from *D. desulfuricans* (24)]. Thus, heme synthesis would appear to be mediated by the conversion of siroheme into didecarboxysiroheme, fol-

lowed by oxidative loss of the northern acetic acid side chains and then oxidative decarboxylation of the northern propionate side chains to vinyl groups (Fig. 3).

The loss of the acetic acid side chains at C2 and C7 for the synthesis of Fe-coproporphyrin is catalyzed by AhbC, which is a radical AdoMet family member. To promote this reaction the enzyme is likely to generate adenosyl radicals and use these to abstract hydrogen atoms in turn from the C3 or C8 positions thereby producing the respective substrate radical and deoxyadenosine. The radical can then fragment by homolytic cleavage of the C-C bond connecting the side chain acetic acid group to the macrocycle, generating a double bond and an acetate radical (Fig. S5A). This process is analogous to the proposed fragmentation of glutamate in the reaction catalyzed by coenzyme B₁₂-dependent glutamate mutase, where an acrylate molecule and glycyl radical are formed prior to the rearrangement into 3-methylaspartate (25, 26). However, in the case of the reaction catalyzed by AhbC the enzyme presumably converts the acetate radical to acetate by providing a further electron (and proton), which could come from a second predicted Fe-S center on the protein. The mechanism as described consumes two molecules of AdoMet, each of which generates an adenosyl radical. Alternatively, an initially produced adenosyl radical could be recycled by using the methyl group of deoxyadenosine as a source of a hydrogen atom to convert the acetate radical into acetate.

The final reaction in the alternative heme biosynthesis pathway, catalyzed by AhbD, sees the transformation of Fe-coproporphyrin into heme. Overall, this process is similar to the HemN catalyzed oxidative decarboxylation reaction of coproporphyrinogen III to protoporphyrinogen IX that occurs during the classical heme biosynthesis pathway (27, 28). As with HemN (and AhbC), AhbD is also a radical AdoMet family member. By analogy with HemN, the mechanism of AhbD is likely to involve the abstraction of a hydrogen atom from the beta-position of the propionate side chains attached to C3 and C8. The resulting substrate radical can then further oxidize to a carbon cation, which could lead to the vinyl product by loss of CO₂ (Fig. S5B). Although a monovinyl intermediate has been identified, it is not known if there is a preferential order for the two oxidative decarboxylations.

The identification of the intermediates of both the heme and heme d_1 synthesis pathways highlights some of the ingenious chemistry that takes place. These beguiling reactions include the incorporation of oxo functionalities in place of the northern propionate side chains during d_1 synthesis from didecarboxysiroheme and the oxidative loss of the northern acetate side chain from the same branch-point intermediate during heme synthesis. The sequence similarity between NirJ and AhbC not only reflects the proteins being members of the radical AdoMet family but may also reflect recognition of the same substrate. The assignment of function to these enzymes will now permit a more detailed mechanistic investigation into how these transformations are mediated.

Our research also defines a fresh role for siroheme, previously only thought to act as a prosthetic group for assimilatory sulfite and nitrite reductases, as an intermediate in modified tetrapyrrole biosynthesis (Fig. 3A). From an evolutionary perspective, it has previously been suggested that siroheme may have acted as a primordial heme where the isobacteriochlorin ring would have allowed greater conformational flexibility, prior to the selection of porphyrins in the later more oxidizing environments (29). There is a view that denitrification came early in evolution and it is possible that the d_1 -containing nitrite reductase appeared before both oxygen and the copper nitrite reductase (30). This paper now highlights how heme can be synthesized from siroheme, providing a biosynthetic evolutionary corollary relationship whereby the functional evolution of siroheme into heme is linked by the appearance of a biochemical pathway allowing the physical transformation of one into the other. The seemingly

coincidental molecular hijacking of the cofactor for assimilatory nitrite reductase to form the cofactor for dissimilatory nitrite reductase was unanticipated; we cannot rule out the reverse appearance first of d_1 heme followed by the hijacking of siroheme by the assimilatory nitrite reductase, which might have emerged once oxidized nitrogen increased as atmospheric oxygen increased. The presence of this branch of tetrapyrrole biosynthesis is consistent with a patchwork assembly model of pathway evolution, and provides a hitherto unsuspected relationship between nitrite assimilation and dissimilation.

Methods

DNA Manipulations. DNA manipulations were performed by standard methods. Amplifications of genes from *P. pantotrophus* (GB17), *D. desulfuricans* (G20), and *D. vulgaris* Hildenborough were performed by PCR using KOD DNA polymerase (from *Thermococcus kodakaraensis*) according to supplier's instructions (Novagen). All constructs generated by PCR were confirmed to be correct by sequencing. Plasmids containing *nirED-L*, *nirGH*, and *nirD-LGH* were constructed by inserting the appropriate amplified *P. pantotrophus* genes into pET3a using the link and lock method described in ref. 18. The *nirH* was cloned into pET14b, and thus the recombinant protein contained an N-terminal hexahistidine tag, whereas *nirG* and *nirD-L* were cloned into pET3a and pASKIBA13 plus vectors, respectively. Plasmids containing *ahbA* and *ahbB* from *D. desulfuricans* were obtained by cloning into the vectors pKK2233 and pET14b, respectively, whereas a plasmid harboring *ahbAB* from *D. vulgaris* was obtained by cloning the genes into pET3a vector using the link and lock method described above.

Recombinant Protein Production. Cell growth and protein overexpression. For protein overproduction, the *E. coli* BL21(DE3) strain was transformed with the appropriate plasmid. The resulting strain was grown with aeration at 37 °C in LB medium containing appropriate antibiotics (50 µg/mL ampicillin, 30 µg/L chloramphenicol, to an A_{600} 0.6–0.8 and the protein expression was induced with 0.4 mM IPTG and cultures were transferred to 16 °C cells for overnight expression. Cells were collected by centrifugation $3,500 \times g$ for 15 min at 4 °C (Beckman Coulter, J130). The cell pellets from 1 L of LB broth were resuspended in 20 mL of 50 mM Tris-HCl, pH 8.0, containing 0.5 M NaCl. Cells were disrupted by sonication (Sonics Vibracell Ultrasonic processor). Cell debris and insoluble proteins were removed from the soluble cell lysate by centrifugation $35,000 \times g$ for 20 min at 4 °C. A cell extract of sulfate-reducing bacteria was obtained from *D. vulgaris* Hildenborough frozen cell pellets.

Purification of 5-aminolevulinic acid and succinyl-CoA synthases. Recombinant *Rhodobacter sphaeroides* 5-aminolevulinic acid synthase (HemA) was produced and purified as described in ref. 28. Recombinant *E. coli* succinyl-coenzymeA synthetase (SuccD) was purified as a hexahistidine fusion protein as follows. Cell lysate containing the overexpressed SuccD was applied to the Ni-Sepharose column equilibrated in buffer A (0 mM Tris-HCl, pH 8.0 buffer). The column was washed with 5–10 column volumes of 50 mM imidazole containing buffer A and the proteins were eluted using the 400 mM imidazole containing buffer A. Only Succ (41,400 Da) has a hexahistidine tag but the associated SuccD (30,000 Da), which was not tagged, also eluted.

Purification of multienzyme cocktail for sirohydrochlorin production. For in vitro preparation of sirohydrochlorin a multienzyme method was used as described previously (15) and outlined briefly in *SI Text* and *Fig. S6*.

Production of siroheme. To make siroheme from sirohydrochlorin, a 10-fold excess of FeSO₄ was added to a sirohydrochlorin solution and the mixture was left at room temperature for 4 h before purification on the DEAE column. The siroheme was purified on a small DEAE column to assist NMR studies with the siroheme and the didecarboxysiroheme derivative.

High performance LC-MS analysis of siroheme and siroheme derivatives. The tetrapyrrole intermediates could be purified after separation from the protein by boiling at 80 °C for 10 min and were analyzed by reverse phase chromatography. Samples were resolved on an ACE 5AQ column (2.1 × 150 mm, 5 µm, Advanced Chromatography Technologies) attached to a Agilent 1100 series HPLC equipped with diode array detector and coupled to a microTOF-Q (Bruker) mass spectrometer. The column was developed with a binary gradient at a flow rate of 0.2 mL min⁻¹. Solvent A was 0.1% TFA and solvent B was acetonitrile. The column was equilibrated with 5% B. Following sample injection the concentration of B was increased to 20% over 6 min and then to 30% at 25 min and 100% at 35 min where it was held for 5 min before returning to starting conditions. The total length of each run was 50 min. Alternatively, for incubations starting from Fe-coproporphyrin III a faster linear gradient was employed starting at 20% B and reaching 100% B in 30 min. The observed masses for these various intermediates are given in *SI Text* and shown in *Fig. S7*.

ACKNOWLEDGMENTS. We thank Dr. Michelle Rowe for technical support. This work was supported by Biotechnology and Biological Sciences Research Council Grants BBE0229441 and BB/E024203 (to S.J.F. and M.J.W.) and Wellcome Trust Equipment Grant 091163/Z/10/Z (to M.J.H. and M.J.W.).

- Horowitz NH (1945) On the evolution of biochemical syntheses. *Proc Natl Acad Sci USA* 31:153–157.
- Jensen RA (1976) Enzyme recruitment in evolution of new function. *Annu Rev Microbiol* 30:409–425.
- Holliday GL, et al. (2007) Evolution of enzymes and pathways for the biosynthesis of cofactors. *Nat Prod Rep* 24:972–987.
- Lager G, Reichelt J, Jahn D, Heinz DW (2010) Structure and function of enzymes in heme biosynthesis. *Protein Sci* 19:1137–1161.
- Warren MJ, Raux E, Schubert HL, Escalante-Semerena JC (2002) The biosynthesis of adenosylcobalamin vitamin B₁₂. *Nat Prod Rep* 19:390–412.
- Ishida T, et al. (1998) A primitive pathway of porphyrin biosynthesis and enzymology in *Desulfovibrio vulgaris*. *Proc Natl Acad Sci USA* 95:4853–4858.
- Raux E, et al. (2003) Identification and functional analysis of enzymes required for precorrin-2 dehydrogenation and metal ion insertion in the biosynthesis of sirohaem and cobalamin in *Bacillus megaterium*. *Biochem J* 370:505–516.
- Rinaldo S, et al. (2011) Observation of fast release of NO from ferrous d_1 haem allows formulation of a unified reaction mechanism for cytochrome cd_1 nitrite reductases. *Biochem J* 435:217–225.
- Palmedo G, et al. (1995) Resolution of the *nirD* locus for heme d_1 synthesis of cytochrome cd_1 (respiratory nitrite reductase) from *Pseudomonas stutzeri*. *Eur J Biochem* 232:737–746.
- Zajicek RS, Ferguson SJ (2005) The enigma of *Paracoccus pantotrophus* cytochrome cd_1 activation. *Biochem Soc Trans* 33:147–148.
- Storbeck S, et al. (2009) The *Pseudomonas aeruginosa nirE* gene encodes the S-adenosyl-L-methionine-dependent uroporphyrinogen III methyltransferase required for heme d_1 biosynthesis. *FEBS J* 276:5973–5982.
- Zajicek RS, et al. (2009) d_1 heme biogenesis—assessing the roles of three *nir* gene products. *FEBS J* 276:6399–6411.
- Yapbondoc F, Bondoc LL, Timkovich R, Baker DC, Hebbler A (1990) C-methylation occurs during the biosynthesis of heme- d_1 . *J Biol Chem* 265:13498–13500.
- Cavallaro G, Decaria L, Rosato A (2008) Genome-based analysis of heme biosynthesis and uptake in prokaryotic systems. *J Proteome Res* 7:4946–4954.
- Lobo SAL, Brindley A, Warren MJ, Saraiva LM (2009) Functional characterization of the early steps of tetrapyrrole biosynthesis and modification in *Desulfovibrio vulgaris* Hildenborough. *Biochem J* 420:317–325.
- Panek H, O'Brian MR (2002) A whole genome view of prokaryotic haem biosynthesis. *Microbiology-Sgm* 148:2273–2282.
- Storbeck S, et al. (2010) A novel pathway for the biosynthesis of heme in Archaea: Genome-based bioinformatic predictions and experimental evidence. *Archaea* 2010:175050.
- McGoldrick HM, et al. (2005) Identification and characterization of a novel vitamin B₁₂ (cobalamin) biosynthetic enzyme (CobZ) from *Rhodobacter capsulatus*, containing flavin, heme, and Fe-S cofactors. *J Biol Chem* 280:1086–1094.
- Kawasaki S, Arai H, Kodama T, Igarashi Y (1997) Gene cluster for dissimilatory nitrite reductase (*nir*) from *Pseudomonas aeruginosa*: Sequencing and identification of a locus for heme d_1 biosynthesis. *J Bacteriol* 179:235–242.
- Brindley AA, Zajicek R, Warren MJ, Ferguson SJ, Rigby SE (2010) NirJ, a radical SAM family member of the d_1 heme biogenesis cluster. *FEBS Lett* 584:2461–2466.
- Xiong J, Bauer CE, Pancholy A (2007) Insight into the haem d_1 biosynthesis pathway in heliobacteria through bioinformatics analysis. *Microbiology-Sgm* 153:3548–3562.
- Oglesby-Sherrouse AG, Vasil ML (2010) Characterization of a heme-regulated non-coding RNA encoded by the *prfF* Locus of *Pseudomonas aeruginosa*. *PLoS One* 5:e9930.
- Bali S, Warren MJ, Ferguson SJ (2010) NirF is a periplasmic protein that binds d_1 heme as part of its essential role in d_1 heme biogenesis. *FEBS J* 277:4944–4955.
- Romao CV, et al. (2000) Iron-coproporphyrin III is a natural cofactor in bacterioferritin from the anaerobic bacterium *Desulfovibrio desulfuricans*. *FEBS Lett* 480:213–216.
- Buckel W, Kratky C, Golding BT (2005) Stabilization of methylene radicals by cob(II) alamin in coenzyme B₁₂ dependent mutases. *Chemistry Eur J* 12:352–362.
- Marsh ENG, Patterson DP, Li L (2010) Adenosyl radical: Reagent and catalyst in enzyme reactions. *ChemBioChem* 11:604–621.
- Lager G, Moser J, Heinz DW, Jahn D, Schubert WD (2003) Crystal structure of coproporphyrinogen III oxidase reveals cofactor geometry of Radical SAM enzymes. *EMBO J* 22:6214–6224.
- Lager G, et al. (2006) The substrate radical of *Escherichia coli* oxygen-independent coproporphyrinogen III oxidase HemN. *J Biol Chem* 281:15727–15734.
- Crane BR, Getzoff ED (1996) The relationship between structure and function for the sulfite reductases. *Curr Opin Struct Biol* 6:744–756.
- Ducluzeau AL, et al. (2009) Was nitric oxide the first deep electron sink? *Trends Biochem Sci* 34:9–15.